





\$ 105

No. 38689



Presented
to the
Library of the
Medical Society
of the
County of Kings

LEEDS UNIVERSITY LIBRARY
Special Collections

Bamji Collection

ASH.



30106022949548


ens and
Un19.

4 1919





MILITARY HYGIENE




THE ELEMENTS OF MILITARY HYGIENE

ESPECIALLY ARRANGED FOR OFFICERS
AND MEN OF THE LINE

BY

P. M. ASHBURN

MAJOR, MEDICAL CORPS, U. S. ARMY



BOSTON AND NEW YORK
HOUGHTON MIFFLIN COMPANY
The Riverside Press Cambridge
1909



3147.12°

COPYRIGHT, 1909, BY PERCY M. ASHBURN

ALL RIGHTS RESERVED

M. L. A.

PREFACE

THERE can be little or no doubt that the sanitation of the army would be greatly improved if line officers and enlisted men should become more interested in the subject, and coöperate more freely and intelligently with medical officers in the efforts to promote it.

This book is written in the hope that it may both inform and interest them and so gain for the medical officer the sympathy and coöperation that he always needs, but now too seldom has. It is also hoped that medical officers themselves, especially those of the volunteer service and militia, may find the book useful in their dual capacities of administrators and teachers.




PART I

THE RECRUIT AND HIS
ENVIRONMENT

"Health and a good constitution are better than all gold ; and a strong body than wealth without measure."

ECCLESIASTICUS 30 : 15.



MILITARY HYGIENE

CHAPTER I

THE RECRUIT

ONE of the principal factors in military hygiene is the individual soldier, or, as he comes from civil life, the recruit. If he enters the service in good physical condition, in good health, and with sufficient intelligence to apprehend and apply the instruction he receives in military hygiene, the problem is almost half solved. The final acceptance or rejection of recruits now rests almost entirely with medical officers; but in certain ways line officers exercise a considerable influence in the matter: in the acceptance at the recruiting station, and in requesting special authority to enlist because of special qualifications. For these and other reasons it is important that they should keep well informed as to the requirements necessary and the qualifications desirable in recruits. We will therefore briefly consider some of these.

The age limits for first enlistments are from eighteen to thirty-five years. Preferably men should be chosen between twenty-three and thirty years. Before the age of twenty-three years most **Age** men are immature, their muscles are not hardened, their hearts not so strong, their minds not disciplined by experience, their appetites and emotions not under control; while their susceptibility to impressions from their environment is such as to make it undesirable to limit their associations exclusively to adult unmarried males.

4 THE RECRUIT AND HIS ENVIRONMENT

After the age of thirty years the man's habits of body and mind are fixed, he takes instruction less readily, his body may have begun to deteriorate, particularly if it has not been well cared-for, and, in many instances, he seeks the service as a refuge, after failing in other walks of life. Many good men may be obtained before the age of twenty-three and between the ages of thirty and thirty-five, but in general the case is thought to be as stated.

The size of the recruit is not a matter of the greatest importance, provided that the development is normal and the man strong and active, except
Size as it applies to assigning him to duty. It is obvious that a very large and strong man is better suited for coast-artillery work than for the cavalry, and that a small, wiry man is in that respect better for the cavalry. The proportions of weight and chest measurements to height are published from time to time in general orders, and those proportions usually represent very well those of strong, well-developed men. An increase of weight, if associated with great breadth of shoulder and large chest measurement, is not necessarily harmful, being usually indicative of strength and endurance. The sturdy, stocky Japanese coolie exemplifies this. Obesity, whereby the man is burdened with his own weight, is of course objectionable.¹

¹ The following table copied from the *Medical Record* of September 5, 1908, and compiled from the data of the Mutual Life Insurance Co. of New York, by its Chief Medical Director, is interesting as indicating somewhat "the influence of overweight and underweight on vitality." The weights considered normal by the insurance company are somewhat in excess of those prescribed by the War Department, the increase averaging about 10 pounds at the ages 25-29, and for heights

The chest should be freely mobile, that is, both sides should move freely, symmetrically, and equally, in expanding and contracting. The measured chest expansion should generally correspond with the figures published in the general orders; yet it should be remembered that one man with very ordinary lung capacity may, by use of the shoulder and below 70 inches. A weight 20 per cent below the average is considered "underweight," and 20 per cent above the average, "overweight."

Table showing the Percentage of Deaths in all Classes, and some Individual Diseases, among Overweights and Underweights, and the General Experience of the Mutual Life Insurance Company.

<i>Causes of Death.</i>		<i>Over-weights.</i>	<i>Under-weights.</i>	<i>General Experience.</i>
Class I.	General Diseases — Acute . .	9.67	9.28	8.90
	Typhoid fever	4.00	3.06	3.94
	Malarial fever	1.27	1.21	1.24
	Influenza	1.47	2.04	1.00
Class II.	General Diseases — Chronic .	13.07	24.59	19.56
	Tuberculosis	2.93	16.98	12.42
	Cancer	4.40	5.57	4.18
	Diabetes	3.40	0.65	1.25
Class III.	Diseases of the Nervous System	19.08	12.16	17.44
	Cerebral Congestion and Hemorrhage, Cerebral Softening, Paralysis . .	14.14	8.47	12.32
	General Paralysis and other forms of mental alienation	1.80	0.84	1.30
Class IV.	Diseases of the Circulatory System	16.01	11.69	11.85
	Organic diseases of the heart	12.94	8.54	10.76
Class V.	Diseases of the Respiratory System	8.54	15.78	11.86
	Pneumonia	6.87	12.34	9.03
Class VI.	Diseases of the Digestive System	10.61	8.54	10.19
	Cirrhosis of Liver	3.47	0.65	1.00
Class VII.	Diseases of the Genito-urinary System	12.01	7.42	8.78
	Bright's Disease and Nephritis	11.07	5.30	6.66
Class IX.	Diseases of Skin and Cellular Tissue	1.20	0.47	0.50
Class XI.	Old age	None	2.04	1.50
Class XII.	Violent causes	7.07	5.57	7.42
	Casualties	4.20	3.43	5.21
	Suicides	2.87	2.14	2.20
Class XIII.	Ill-defined	2.60	2.50	3.98

6 THE RECRUIT AND HIS ENVIRONMENT

back muscles and retraction of the abdomen, show an apparent expansion of $3\frac{1}{2}$ inches, while his companions with a capacity as great; may, through ignorance of tricks or of what is desired, show only $1\frac{1}{2}$ inches. It is desirable that the chest be fairly large, to afford ample room for the free working of the heart and lungs, as on such finally depend both strength and endurance. A generally well-developed and strong man will show well-developed and prominent chest muscles.

Certain types of chest frequently indicate a predisposition to, or the actual existence of, disease, and demand careful consideration even when not constituting actual causes for rejection. The long, narrow chest, with prominent or "winged" shoulder-blades, depressed or flat below the collar-bones, and forming an acute angle at the divergence of the ribs below the breast-bone, frequently marks a predisposition to consumption.

A large, barrel-shaped chest, as deep from front to back as transversely, and showing a relatively small movement, is often associated with emphysema and asthma.

A chest more prominent on one side than on the other is apt to be associated with curvature of the spine, or old or chronic pleurisy.

Marked prominence or depression of the breast-bone, especially if associated with "beading," or lumps on the ribs near the breast-bone, frequently results from rickets, and may be accompanied by other bone deformities.

The abdomen should be well muscled, and firm when the applicant is standing. It should move easily and naturally in respiration, should not be pendulous, and should be firm, particularly in

Abdomen

its lower parts, just above the groin. Weakness here will be indicated by a bulging if the patient strains, as at stool, or if he stands on his toes and coughs hard. Such bulging, especially if it be marked or shows a tumor-like swelling, indicates a predisposition to, or the actual existence of, a hernia.

The umbilicus or navel is another frequent site of hernia, and should be noted in that connection, as should any abdominal scars resulting from operations, as for appendicitis. Another, though less frequent, site of hernia is the femoral. This is shown by a fullness or swelling in the lower part of the groin, in the highest part of the thigh rather than in the lowest part of the abdomen.

Large masses or tumors that can be felt through the abdominal wall, or that can be seen to produce inequality or fullness, are causes for rejection.

Distention or prominence of the veins on the surface of the abdomen, or a varicose condition of them, is generally an evidence of impaired circulation or liver disease, and should cause rejection.

Hemorrhoids, or piles, while apparently not connected with the abdomen, are the result of distention of the rectal veins, and may be due to liver disease, to abdominal tumors, to constipation, or other intra-abdominal conditions. They are manifested as tumors, usually of a bluish color, within or about the anus, which vary in size from that of a small pea to that of a man's thumb. Of themselves they are not a cause for rejection unless of large size or producing symptoms, but their presence should always be noted. Internal piles are not visible, and usually make their presence known by burning or irritation in the rectum, or by the passage of

8 THE RECRUIT AND HIS ENVIRONMENT

blood in the stools. In examining for them, note should also be made of other abnormal conditions of or about the anus, such as fissures or cracks in the skin and mucous membrane; fistulas, little openings near the anus from which there may be a slight discharge of pus or fecal matter; or abscesses or boils. Any of these conditions may so disable the man as to unfit him for service.

A history of abdominal trouble, such as severe indigestion, gallstone colic, vomiting of blood, should cause rejection.

The examination of the head is very important, as revealing defects in most of the special senses and as offering important evidence as to the man's character, habits, and abilities. The evidence on the latter points is to be gained largely from the shape of the head and the expression of the features, and, while such evidence is not always reliable, certain heads and faces are so distinctly indicative of stupidity or vice that there should be no hesitation in rejecting their possessors. Then, too, hideous or disgusting deformities should at once cause rejection, for the sake both of the victim and those with whom enlistment would associate him, even though they are not of a character to incapacitate directly. The vision and hearing should be tested in strict accordance with the regulations governing the subject, and failure to meet the requirements is cause for rejection.

In addition to the test for vision, however, the *eyes* should be examined for any signs of inflammation, redness, watering, drooping of the lids, scars or deformities, granulations, styas or boils, undue prominence of the eyeball, inability to move the

eyes in all directions, and discoloration or blurring of the cornea.

The *ears*, in addition to the tests for hearing, should be examined for any discharge, whether watery or purulent, and if present it should be a cause for rejection. **Ears**

The *nose* should be examined as to its freedom from obstruction, by having the applicant close first one side and then the other with his fingers and then breathe and blow through the free side. Incidentally this may call attention to a foul-looking discharge or a foul-smelling breath, either of which may justify rejection. A sunken or much scarred nose is often an indication of syphilis, while a red, bulbous nose, even if not indicating alcoholism, is sometimes an indication of indigestion. It should be noticed that the man habitually breathes through his nose. **Nose**

The *mouth* and *throat* should always be carefully examined. **Mouth**

The condition of the *teeth* is the first point to be noted here. It is desirable that all should be present and good, but there must be at least enough to permit of proper mastication of the food, and for this purpose at least four grinders are demanded, two above and two below, and so disposed as to permit their effective use. Enough of the front teeth should also be present to permit of the biting of food and to preserve the symmetry of the face. At times men are accepted with no upper teeth, but wearing plates. This should only be done by special authority and in special instances. Insufficient or poor teeth are apt to produce digestive disturbances, particularly on campaign, when hard bread may be issued, **Teeth**

10 THE RECRUIT AND HIS ENVIRONMENT

and when the preparation of the food in general may not be as good as in garrison. The examination of the teeth should not only relate to their presence and the presence or absence of cavities, but also to their state of cleanliness. Neglected, filthy teeth and gums may seriously menace the general health, while they certainly predispose to loss of teeth, and, in many instances, indicate general carelessness or filthiness. Sores or ulcerations on the lips, tongue, gums, or inner sides of the cheeks should be carefully noted, as they may indicate syphilis or other general disease, and should at least lead to further and more careful search for syphilis. The same remark applies to milk-white patches, seen especially on the tongue.

In ill-kept mouths the *gums* may be spongy and bleed spontaneously or on pressure, or pus may exude from between them and the teeth; or they may
Gums be greatly retracted and expose the roots of the latter.

In these and other cases the teeth may all be present, yet in such poor condition as to forbid proper mastication, or to threaten their early loss.

The most common and easily detected *throat* trouble is enlargement of the *tonsils*. This condition usually
Throat indicates a liability to frequent attacks of sore throat, and should always excite inquiries as to this, and while not necessarily a cause for rejection, unless very marked, should be noted as a point against desirability. Ulcers, white patches, membranes, and an appearance of acute inflammation in the throat, should cause rejection.

Hoarseness, or loss of *voice*, or cough, should at least delay, and if persistent prevent, acceptance.

Numerous scars on the *tongue*, *lips*, and *cheeks*, unless clearly and certainly accounted for otherwise, should cause rejection as being probably due to bites during epileptic fits, or to syphilitic or other ulceration.

Tongue

Too much importance should not be attached to a coated tongue, but a raw-looking, bright red, or tremulous one should direct careful scrutiny to the general condition and the habits.

Enlarged *lymphatic glands*, or "kernels," are most frequently noticed in the neck or under the jaw. They are frequently indicative of inflammation or ulceration in other nearby regions, especially the mouth or throat, and their presence should lead to careful examination for such conditions; but at times their presence is indicative of a general infection, such as syphilis. Usually, if the enlarged glands are numerous, or the enlargement great, they constitute a proper cause for rejection. When the enlargement is localized at one part of the neck only, it points to an infection entering at a part near it. Thus such a gland at the angle of the jaw may be due to inflammation of a tonsil of the same side; one under and near the jaw to a bad tooth, etc.

Lymphatics

The *scalp* should be carefully examined as to its cleanliness and general care, for the presence of lice, which may be seen as such or may be revealed by the "nits" or eggs, little whitish bodies attached firmly to the hairs; for ringworm, scabbiness or scald-head, patchy baldness, wounds or scars, etc. The presence of vermin is a cause of rejection in two ways, as indicating a lack of personal care, and as a danger to be introduced into crowded barracks.

Scalp

12 THE RECRUIT AND HIS ENVIRONMENT

Superficial suppurative scalp diseases should generally cause rejection, as being contagious or as being caused by scratching excited by lice. Patchy baldness, if the patches are sharply defined and the skin is smooth, clean-looking, and not scarred, is apt to be harmless. If the skin is inflamed and scarred, and stumps of broken hairs are present, it should cause rejection. Irregular and incomplete baldness occurring in patches or tracts, giving at times a "moth-eaten" appearance, is most apt to be due to syphilis, and should cause careful examination for other signs of that disease.

The upper limbs should be examined as to their development and musculature, their free mobility in all joints and in all directions, the presence of
Arms all bones and joints, and a reasonable amount of dexterity and strength. Atrophy or wasting of a group of muscles or a part, inability to bend or make free use of a joint, and similar defects, should be carefully looked for, as they may otherwise be missed. Loss of one or more joints of a finger, swelling and deformity of a joint (baseball finger), or stiff contracted fingers are not uncommon, and, if in the fourth or fifth fingers, are not serious matters, though they should be noted. If in the second or third fingers, the matter is more serious, yet the applicant will usually be able to handle a gun and perform his other duties. Loss of a thumb is a serious defect, and in most instances would justify rejection. All deformities from badly united fractures or other causes should be noted, and should cause rejection unless the applicant can demonstrate his ability to execute the movements necessary in the performance of his duties.

The legs should likewise be examined as to their

development, musculature, size, mobility, etc. The applicant should be required to move all the joints, to jump, hop, and otherwise demonstrate his ability to use his limbs. Marked deformities, such as clubfoot, shortening of one leg, stiffness, marked swelling or marked limitation of motion in a knee or hip, should at once cause rejection. Less marked deformities, such as knock-knee, bow-legs, crooked shins, etc., need not cause rejection unless manifestly interfering with the free use of the limbs, or rendering the subject's appearance ludicrous or unmilitary. Here, again, marked wasting of a part or of a group of muscles will often be found due to a joint lesion, a paralysis, or other nervous trouble, that would serve to disqualify the applicant. In addition to these defects, common to both arms and legs, the lower extremities are subject to other affections rarely or never seen in the upper.

Legs and Feet

Varicose or dilated veins are often seen. They usually show as swollen, bluish, and more or less tortuous vessels beneath the skin, and may be individually as large as a man's finger. They are most commonly seen on the lower legs, but are not very rare in the space behind the knee, and may extend up the inner side and front of the thigh to the groin. If marked, they should cause rejection. If only slight, and in otherwise desirable applicants, they may be passed, but should always be noted. At times they cause the appearance and persistence of very chronic ulcers, which occasionally bleed, pain, or otherwise disable the man. Such ulcers are especially apt to show on the front or to the inner side of the lower half of the leg. They are long in healing, are apt at any time to break open again, and when healed

14 THE RECRUIT AND HIS ENVIRONMENT

usually present brownish, discolored scars. Such ulcers, or such scars if accompanied by varicose veins, should cause rejection. The legs are especially prone to *chronic ulcerations*, which may be caused by syphilis, tuberculosis, leprosy, and other infections. In general these are causes for rejection.

A rough, nodulated, tender shin-bone is not an unusual sign of syphilis.

The *feet* require particularly careful examination, as on their integrity, usefulness, and strength (or what we might call their durability) depends a large part of the man's capacity for "soldiering." The best known and probably the least important of the foot-ills is corns. They are best known because most persons at some time have at least one, and they are one of the least important ills because in most instances they cause not even serious inconvenience, and they usually disappear with the use of properly fitting shoes and ordinarily good care of the feet.

If numerous and painful, however, or if soft corns, situated between the toes, they may justify rejection.

Sweating feet, which soften, blister, redden, and burn, may cause rejection, as they are not apt to stand much marching.

Stinking feet may justify rejection in most instances, as constituting a nuisance in barracks.

Bunion, an inflammatory and deforming joint-affection, usually at the base of the great toe, will, if very marked, disqualify.

Hammer-toe, a condition in which a toe is flexed and the end presses on the floor, while a knuckle projects above, is very apt to cause lameness and may justify rejection.

Overlapping toes may disqualify in a somewhat similar manner.

Deformities of the arch of the foot are very important. It may be too high, so that the weight is borne on the ball of the foot and the heel, the intermediate parts remaining clear of the ground. This is a relatively rare condition, but may disqualify. A common condition is the opposite, or *flat foot*, wherein the whole length and breadth of the foot touches the ground and the arch is almost or entirely obliterated. Such a foot in a white man is usually a poor marching foot, and, unless the applicant is a desirable man and states that he is a good walker, and that the foot never gives him trouble, it should cause rejection. When passed, its existence should always be noted, in case the man should be discharged for it later. The condition is more common but less important in negroes, but it may cause trouble in them also, particularly in the course of long marches.

Another type of weak foot is not really flattened but tends to rotate outward, thus bringing the inner margin of the foot nearer to the ground and more directly under the weight of the body, where a greater weight comes upon it, causing it to tire and eventually to break down and become flat. It may justify rejection.

The genitals should be well-formed and normal. Both testicles should be present and in the scrotum; it should be possible for the applicant to retract his foreskin, and the penis should **Genitals** not show gross deformities, such as the opening of the urethra being placed far back from the end. Scars on the penis should prompt investigation as to their nature, whether syphilitic, chancroidal, or otherwise. The man should be required to express the contents, if any,

16 THE RECRUIT AND HIS ENVIRONMENT

from the urethra. Pus should at once cause rejection, as should any ulcers or sores on the penis, redness, puffiness, or signs of marked irritation about the urethra, or the presence of vermin, commonly known as crab-lice. Inflammation of the glans beneath the foreskin, and venereal warts, while not in themselves disabling, will often justify rejection as capable of exciting suppurative buboes, and as resulting from venereal disease or uncleanness, or both.

Buboes are swollen and inflamed lymphatic glands, and are found in the groin. They may be single or multiple, on one or both sides, and may vary in size from that of a small cherry to that of an orange. While not all groin buboes are venereal, the great majority of them in applicants for enlistment are, and they justify rejection, even though the applicant insists, as he probably will if his venereal disease is not plainly evident, that they are due to "strain."

Varicocele is a condition of enlargement of the veins of the scrotum. It is most often found, and is apt to be most marked, on the left side. The condition frequently causes no symptoms, though it is often said to cause pain, and is apt to be pleaded as an excuse from arduous duty. If present, it need not cause rejection unless very marked or unless the applicant says that it produces symptoms, but when passed it should always be noted, and the applicant be made to agree to be operated upon if symptoms begin, or if the condition comes to interfere at all with his duties.

Hydrocele is an enlargement of the scrotum, usually on one side, caused by an outpouring of clear fluid into the membranous sac surrounding the testicle. It produces a tense, painless, and often translucent swelling,

which may at times be hard to distinguish from a tumor of the testicle or a hernia. Any one of the conditions, however, may suffice to exclude the applicant in time of peace.

While not much reliance can be placed on an applicant's denial of venereal disease and his assertions as to perfect genital health, he should be questioned on the subject, and should be asked as to the frequency of urination and his ability to pass a good stream of urine. Dribbling or inability to pass a good stream, or to hold or control his urine, should cause rejection.

The skin will necessarily be noticed and examined during the examination previously described. One of the first and most important points to be noted is its cleanliness. A dirty, foul, stink-

Skin

ing skin usually indicates the type of man not wanted in our army, and it justifies rejection without further examination. Certain men, however, appear with their skins soiled with sweat and dust simply because of lack of opportunities for cleanliness, and if their general appearance and their stories indicate that they are desirable men, they should be given an opportunity to bathe and then be examined. It must not be forgotten that a filthy man may bathe for examination, but reveal his usual habits by his linen. However revealed, filthy habits should cause rejection.

A skin showing the marks of severe and generalized scratching is usually indicative of disease or vermin infestation, and is therefore objectionable.

All forms of contagious skin diseases, such as ringworms and syphilitic eruptions, should cause rejection, or, in the person of an enlisted man, call for prompt treatment. As the line officer cannot be ex-

18 THE RECRUIT AND HIS ENVIRONMENT

pected to differentiate such affections, he will be on the safe side if he treats them all as suspicious, though he should if possible learn to recognize acne, the common "pimply" eruption seen on the chests, backs, and faces of many young men. It is usually not a severe affection, and as it may coexist with good health and good habits, it should not cause rejection.

While a few men who are not really physically fit get into the service, there are a good many more who are morally or mentally unfit, and they are the men who later spend much of their time in the guard-house or who swell the number of desertions. Against this class of men the recruiting officer works at a disadvantage. Unfortunately, almost any man, no matter what his character, can get letters of recommendation, so that such letters are really of little importance, and the officer is forced to rely largely on his own ability to judge men by appearances, — a notoriously unsafe thing, and an almost impossible thing for some individuals. Some aid is obtained, though, by attention to the applicant's general bearing, his facial expression, his quickness of apprehension, the condition of his person and his clothing, and such evidences of alcoholism, venereal disease, good habits, and truthfulness as the examination may bring out. When a man declares that he never drinks, although his breath is strong with alcohol at the time, or that he has never had venereal disease, though he shows a scar on his penis and another in his groin, or has a suppurative bubo that he says is due to a strain, the inference is fair that he is a liar and undesirable.

Under our system of voluntary enlistment, and as long as there are plenty of applicants, it is better to

reject all those about whom the officer cannot feel well satisfied; and this whole chapter is based on such an assumed condition of affairs. In time of war, or when the supply of applicants for enlistment is not up to the demands, many men can be passed who would be excluded under our assumed conditions; but in all such cases they should only be passed by medical officers exercising great care, and every defect should be noted, both as to its presence and its degree, as constituting a possible ground of application for a pension.

The man being in the service and any defect coming to his commanding officer's attention, he should be referred to a medical officer to have it corrected if practicable, but at any rate to have it noted, in the interests of justice to the man and to the government, and, if it be sufficiently serious, to have the man discharged before he breaks down in time of stress, with possibly serious results.

In this as in nearly all respects, the medical officer's duties can be performed more promptly and effectively if he has the coöperation of the line officer.

CHAPTER II

PERSONAL HYGIENE

THE recruit having been accepted for the service in good health and good physical condition, it is his duty to himself and to his government to maintain those desirable conditions unchanged except for the better, if it be possible for him to do so. It will be possible in large part, if he is taught how to do it, and it is therefore incumbent upon his officers to teach him at least the rudiments of personal hygiene, and those rudiments may be summarized as follows:—

First of all he must be taught the importance of an abundance of fresh air both day and night, as a health-giving, strength-producing, and life-prolonging measure. He must know that “colds,” sore-throat, pneumonia, and kindred troubles are not caused by cold air, but by micro-organisms; that those micro-organisms are much more numerous and much more apt to harm him in a warm, “close” room, with impure and rebreathed air, than in the open; and that such diseases are more common in cold weather, because in his efforts to exclude the cold man is apt to include such air. The habit of breathing deeply and thoroughly changing the air in the lungs is imparted to him by his gymnastic and military drills, and is helpful.

He should also learn early that his rest is important and should be taken regularly, in order to preserve his health and enable him, for that reason, to do without it more easily when the neces-

sity arises. He should get eight hours of sleep in each twenty-four, while four hours more should be allowed for meals and short rests, changing of clothing, etc. In the remaining twelve hours he should do his work and his play. Regularity in rest and work is of great importance, and that is probably one reason that so many men improve rapidly in physique after entering military service. On the other hand, it is debilitating and demoralizing to steal the hours from sleep for drinking and venery. It is desirable that the night's rest be taken in clothing different from that used in the daytime, and the latter should be allowed to air. Every man should possess and use night-clothing, and the not uncommon habit of sleeping in the underclothing should be discouraged. No man should have a "barrack odor" clinging to him, and, if ventilation and clothing are looked after, he will not.

The most beneficial exercise is that obtained incidentally to work or recreation. Exercise taken purely as such is usually repugnant, and takes on the aspect of work to men who feel that they **Exercise** already have an abundance of it. It should therefore be taken with some other incentive than is furnished by official orders. Baseball games, field sports, and other athletic competitions should be encouraged, and the effort should be made to interest and include all the men in them, and not merely a baseball nine and a few expert athletes in each post. The objects to be attained by exercise are the increase of the strength of the heart and the capacity of the lungs, on which two depend endurance, the increase of muscular strength, of nervous stability and control, of digestion and assimilation, and keeping the emunctories (the channels of waste excre-

22 THE RECRUIT AND HIS ENVIRONMENT

tion, such as the skin, kidneys, and bowels) active. No form of exercise at present encouraged in the army fails to promote some of these ends, so they are all good.

Cleanliness of person, clothing, and bedding should, and usually does, become a habit of life with the soldier ;

Cleanli- but some men require much watching and
ness admonition to make it become so, and frequent inspections should be made, in order

that the admonition may not be wanting. In general, it may be stated that dirty soldiers are either recruits or men who through drink or vicious habits have suffered in their self-respect. Body lice are rarely seen in our service except in guard-houses, where they may be introduced by captured deserters or other prisoners. Bed-bugs are occasionally found, and head-lice and itch less frequently. All of these pests should be carefully watched for and exterminated when found, as not only are they disgusting in themselves, and indicative of careless or filthy habits, but it seems quite possible that they may act at times as carriers of disease. Bed-bugs, when once introduced in barracks, are at times very hard to dislodge, particularly if the buildings are old and show many cracks in the walls.

Personal cleanliness should be understood as including the exclusive use of one's own linen and toilet articles. Disease may be transmitted readily by the indiscriminate or common use by several persons of the same towels, sponges, wash-cloths, combs, hair-brushes, shaving-brushes, etc. Among diseases which are most commonly transmitted by these means are itch, ring-worms and other skin diseases, granulated eyelids, gonorrheal infection of the eyes, body vermin, and,

occasionally, such infectious diseases as typhoid fever or smallpox.

It is not necessary that a man should bathe his body daily unless he is engaged in particularly arduous or dirty work, but he should bathe at least twice a week in summer and once a week in winter, and as much oftener as is necessary to keep his skin clean and free from all odor, especially from the feet, crotch, genitals, and armpits, and particular attention should be given to those parts. Once or twice a week, or oftener if necessary, the washable clothing,—shirts, drawers, and socks,—should be changed, and fresh put on. More frequent baths and changes of course do no harm, but they should not ordinarily be demanded.

The soldier cannot well give too much attention to his *feet*, as he thereby keeps them in good condition and directly increases his value as a soldier. Any trouble with them should at once be corrected by the man, by means of bathing, clean socks, properly fitting shoes, proper nail-trimming and cleaning; or, if of a kind not corrected by such means, it should be reported to the surgeon.

The *hands* are not subject to so many disabilities as the feet, but they should be kept clean and sound, the nails clean and trimmed, and it should become a matter of routine to wash them before taking food, as diseases so diverse as lead-poisoning and typhoid fever may result from the neglect of such a practice.

The *scalp* should not be neglected in the general cleaning, and its cleansing is facilitated by having the hair short.

Bedding should be aired and bed linen changed weekly.

26 THE RECRUIT AND HIS ENVIRONMENT

longer, at a time. Meanwhile, the man delays going to a dentist whom he will have to pay for his services, hoping that the army dentist will be at the post soon; and, knowing that the post surgeon is not a dentist, he does not report to him. Every company commander should therefore instruct his men somewhat as follows:—

1st. Every man should have a tooth-brush, and should so use it at least twice daily, and preferably oftener, as to remove from the teeth and gums all particles of food or other foreign matter, brushing away from the gums and in all cracks and crevices.

2d. That he should after each meal remove from between the teeth, with a wooden or quill tooth-pick, or with a thread, any pieces of meat or other food, and not allow it to remain and putrefy.

3d. That he should promptly consult a dentist or a medical officer about any trouble arising in the teeth or gums. The medical officer can at least give him proper advice, and in many instances proper treatment.

Much harm is done by careless spitting, urination, and defecation in improper places, by men ignorant of the harm they may thus do. Because apparently healthy themselves, it does not occur to them that they may, in answering a perfectly natural call, transmit disease to others. Yet, as will be shown in other places, they may be giving off the organisms causing typhoid or other infections in their urine or feces, or those causing diphtheria, tuberculosis, or other disease in their spit, and such organisms may then soak or wash into the water-supply, or may be carried as dust or

Spitting and al- lied Of- fenses

mud to the mouths, hands, food, or breath of healthy persons, and so infect them. Decency and patriotism should prompt a man to show as much regard for the health of his comrades as for his own.

The question of venereal disease will be dealt with in a subsequent chapter. Much of it could be avoided if the ignorance and misconception regarding sexual life could be dissipated. Every man should know that sexual continence

Sexual Life

is compatible with perfect physical health, while its maintenance is far greater evidence of character and "manliness" than is the association with prostitutes, the abasement of virtuous girls, or the lowering, self-respect-destroying and demoralizing practice of masturbation. The sexual organs and sexual desire are placed in man that he may procreate and replenish the race; but there is no penalty for his failure to do so, and the sperm necessary for the purpose will be discharged in sleep when it has accumulated in sufficient quantity. "Wet dreams" and involuntary seminal emissions are not, therefore, necessary evidences of "loss of manhood" or of any other disorder, but are seen to be the natural and healthy method of disposal of the sperm not used in the legitimate sexual intercourse of married life. Every normal man has periods of sexual excitement and desire, which constitute one of Nature's powerful influences in the perpetuation of the race; but self-respect should prompt and self-command enforce a determination that such desire shall not lead to acts that violate the laws of religion and society. These periods of excitement and desire can be made less frequent, and continence promoted, by the avoidance of lasciviousness in speech, conduct, reading, and thoughts,

28 THE RECRUIT AND HIS ENVIRONMENT

by thorough cleanliness and otherwise complete neglect of the genitals, and by a regular, sober, active life.

The habitual and excessive use of alcohol is everywhere conceded to be harmful, while the moderate or

Alcohol judicious use of it excites much discussion, but is, in this country at least, falling into

disfavor. Owing partly to the prohibition of the sale of wines and beer on government reservations, the conditions under which a soldier now indulges in alcohol are in most instances not conducive to "moderate" or judicious use of it; so that, laying aside all theory, and dealing only with the condition confronting us, it is our duty to advise and encourage total abstinence in the soldier. This we may be able to obtain in certain instances, but it does not seem probable that it will be a universal characteristic of soldiers during the active service of the readers of this paragraph. We should, however, particularly try to inculcate it in the young recruit, as the man who does not begin to drink before he is twenty-five years of age is much less apt to become a drunkard or a habitual user than he who begins before he is twenty, and there is a good chance that, if he abstains until that age, he will always do so. On the other hand, the impossibility of knowing who are, and who are not, capable of using alcohol judiciously should prevent our encouraging what we regard as judicious use in soldiers. Every military post shows in its guard reports, its sick reports, and in unrecorded ways, the injury and loss of service resulting from excessive use of alcohol; so that it is not necessary either to quote military statistics or to go to civil life for the lesson. However, the very nature of a soldier's calling so restricts his freedom of action, so excludes him from many

refining and restraining influences, and so throws him in the way of temptation to drink, that we should highly honor him in the mass for his good conduct, and encourage him in it as an individual.

If used at all, alcohol is preferably and least harmfully taken with or after meals.

It will do little good to warn men against the use of strong alcohol, of flavoring extracts, bay rum, and wood alcohol, as only abandoned drinkers use such preparations as beverages, and such men are not apt to heed the warning. However, it is important to know that wood alcohol and its preparations are more poisonous than ordinary alcohol, and that death, or complete and permanent blindness, may follow their use.

Probably the least harmful and the most general of the so-called "bad habits" is the use of tobacco in one form or another. In our army the methods of using it are smoking and chewing. Of **Tobacco** these chewing is probably the most objectionable as fouling the mouth and causing promiscuous spitting. Moderate smoking, indulged in after meals and in periods of relaxation, cannot be said to be very harmful, if at all so; but smoking in excess may do harm by causing digestive, respiratory, and nervous disturbances, among which may be heartburn, headache, palpitation or irregularity of the heart, insomnia, tremor, cough, and hoarseness. Just what constitutes excess may not be stated, as an amount harmless to one man may be enough to cause injury to another. In general terms we may say that an amount causing any of the above-named or other symptoms in any man constitutes an excess for that man. As to the various methods of smoking, the same general principles apply. Cigar, pipe, and cigar-

30 THE RECRUIT AND HIS ENVIRONMENT

ette are equally harmless if not used to excess. The main objection to cigar-smoking is the expense; to pipe-smoking, the irritation of the mouth, in rare instances eventuating in cancer; and to cigarette-smoking, the habit of inhaling the smoke and thus exposing a much greater surface to the fumes, causing irritation of the vocal organs and the bronchial tubes, and the habit of rolling and smoking a cigarette at each opportunity, so that the total number consumed becomes excessive. The habit of smoking and the proper indulgence of the habit do the soldier so little harm and so greatly increase his comfort and contentment that they should not be interfered with. Stale tobacco-smoke in quarters, however, gives a very unpleasant odor, and the room should be daily opened and aired sufficiently to prevent or dissipate this.

There should be no necessity for telling men to dress properly for the season and the climate, but the necessity nevertheless exists, owing to the carelessness, ignorance, or perversity of men.

**Temper-
ature
and
Climate** It is neither profitable nor wise for an officer to undertake to prescribe just what his men shall wear at all times, as men differ in their capacity to stand heat and cold, and one man may be able to maintain the highest degree of health and comfort by never wearing flannel underclothing, while his neighbor may promote both by wearing it the year round, and the rest of the men in the squad-room by wearing flannel in winter and cotton in summer. But men should be taught that exposure to either heat or cold lowers the resistance of the body and predisposes it to disease; that body-warmth is promoted by woolen clothing and that linen and cotton

make cool clothing; that alcohol is not a fit substitute for an overcoat or a rain-coat; that wet feet should be dried and dry coverings substituted for wet ones after marching or other exercise; that good intentions or thoughtlessness do not justify reckless exposure to extremes; and that it is their duty to the government, as well as to themselves, to use discretion in dress and to try at all times to avoid sickness.

Exposure to strong sunlight is by some persons alleged to be the cause of some of the evils heretofore credited to the heat of the sun, and it is further stated that these effects may be obviated by the use of hat-linings and underclothing of black, red, or orange, to exclude the actinic rays. These statements, however cannot yet be said to be established facts, but must await the results of further observations and experience.

In all climates and at all times the dress should be comfortable, and should hamper the movements, especially of the chest and abdomen, as little as possible. The equipment should be as light as circumstances will permit, should also be arranged with the idea of interfering with motion, particularly of the chest and abdomen, as little as possible, but should contain the necessary articles.

While the government supplies the necessities of the soldier and supervises his hygiene to a considerable extent, it is seen already that the man himself is responsible for much, and there are a few more golden rules that he should learn early for his own protection, and that will bear frequent emphasis and repetition, especially in campaign.

1. Be clean in person, clothing, and surroundings.

32 THE RECRUIT AND HIS ENVIRONMENT

2. Eat no food but good food, and only with clean hands.

3. Drink no water from unauthorized or doubtful sources, unless it is boiled,—plain, or in tea or coffee.

4. Abhor, avoid, and destroy vermin, whether lice, fleas, ticks, flies, mosquitoes, roaches, mice, rats, or other varieties.

CHAPTER III

FOODS AND THEIR PREPARATION

It is now trite and partly true that "an army travels on its belly," and plainly evident that the health and happiness of the individual soldier depend very greatly on what he puts into his belly. This is a matter largely in the control of the company commander, and it is therefore important that he should have some knowledge of foods and their preparation, and should pay great attention to mess-administration. Many company commanders do so, but others, knowing little of the subject, delegate the entire matter to their mess-sergeants, who may know less, and the men suffer. The government ration and the regulations concerning its use are very liberal, and if a company does not have an abundance of good, nutritious, and attractive food, the fault is nearly always in its handling after it reaches the company.

Foods are of different classes, and a judicious and proper mixture of these is necessary for the maintenance of health and strength. **Classes**

Proteids, or albuminous foods, are the class represented by the white of egg or lean beef. They occur also in vegetables, beans and peas being particularly rich in proteid, and this fact makes an exclusively vegetarian diet a health-sustaining possibility, which a proteid free diet is not. Animal proteid is more easily and completely digested than that from vegetables, and to most persons is also more agreeable. An exclusive meat diet

34 THE RECRUIT AND HIS ENVIRONMENT

can maintain health and strength for long periods of time, and Parkman relates in "The Oregon Trail" an experience extending over several weeks, during which he accompanied Indians on a buffalo hunt, and the whole party performed great amounts of work, during almost the entire time, on a diet of buffalo meat, relieved with a few berries on one or two occasions. Nevertheless, the tendency in those accustomed to a mixed diet is to suffer from albuminuria and diarrhœa when using an excess of proteids.

Fats are the class of food represented by the fatty tissues of animal bodies and the various vegetable oils. They are very valuable as foods, owing to their great fuel-value and the amount of nutriment contained in a small bulk; but if taken in excess they are difficult of digestion and cause dyspepsia and diarrhœa. They are especially valuable in cold climates, a fact which we recognize in our practice of eating pork and other fat foods more freely in winter.

Carbohydrates are the sugars and starches, represented by cane-sugar, fruit-sugar, and others, and by starchy vegetables such as potatoes and grains. For practical purposes we may say that the carbohydrates are of vegetable origin. Cellulose, as found in the husk and fibre of plants, is another form of carbohydrate, having a food-value for herbivorous animals but none for man. The carbohydrates have about the same fuel-value, weight for weight, as the proteids, and less than that of the fats. Starches may be taken in large amounts, but sugar, if taken in excess or in too concentrated a form, causes dyspepsia and other disorders.

Inorganic salts, as represented by common salt and

other salts contained in the foods, have no fuel-value, but are necessary in maintaining the nutrition of the body, and the same is true in even greater degree of *water*.

Organic acids, flavors, and condiments, as found in fruits, many vegetables, and in spices, are of value secondary to the preceding classes, and principally for the relish they impart.

For many years Voit's standard was taken as representing the approximate amounts of the three main classes of foods necessary each twenty-four hours for the maintenance of health and vigor in the average working adult.

Amount of Food

According to that standard the requirement was 118 grams of dried proteid, equaling about one ration of beef, 56 grams of fat, and 500 grams of carbohydrate, that is, of starch and sugar. The total value of this diet was over 3000 calories.

Recently, Professor Chittenden of Yale University has shown that health, strength, and vigor can be maintained on half the amount of proteid, without any increase in the fats or carbohydrates.

Meats are mixtures of proteids and fats, and contain water and salts also. They vary in palatability, cost, digestibility, and nutritive value; but all meats have nutritive value, though if diseased or decomposed they are not available for use.

Meats

In our service *beef* is the standard meat for issue, mutton, pork, fowl, and fish being substitutes issued only occasionally; and, considering the cost, nutritive value, palatability, digestibility, tolerance, etc., it is the most valuable meat.

Mutton is almost equally valuable for nutrition, but

36 THE RECRUIT AND HIS ENVIRONMENT

many persons dislike the taste of it, and almost all tire of it sooner than of beef.

Pork is fatter than either beef or mutton, and is therefore much more used in cold than in warm weather. It is more difficult of digestion than beef and not so generally liked, though bacon is much used in the army and is a popular and valuable food, especially in the field.

Veal is seldom issued to troops. Its nutritive value is somewhat less than that of beef, and it causes diarrhoea in some persons.

Fowl, as chicken or turkey, constitutes a special treat on national holidays and festal occasions. If freshly killed and healthy fowls are served, it is valuable as well as appetizing food; but when birds long kept undrawn in cold storage are issued, as is not infrequently the case, their quality should be suspected and each bird carefully scrutinized, and, if of doubtful appearance or odor, rejected.

Fish vary greatly in palatability and nutritive value, but all of them keep poorly, especially if allowed to die slowly and if not packed in ice or frozen. The meat should feel firm to the touch, and not crush on gentle pressure. Dried and canned fish, as issued, are practically always good, but they are not suitable for frequent or prolonged use, as men very soon tire of them.

Eggs consist of animal proteid and fat, and are classed with the meats. They are nutritious and usually easy of digestion. They may be cooked in a great variety of ways as a main dish, and are very frequently used in other dishes, such as puddings and cakes.

All meats should be eaten cooked, as proper cooking improves the flavor, increases digestibility, and destroys

parasites that might otherwise cause disease in the eater. Among the disease-producing organisms which may be conveyed by meats but which can be destroyed by heat are :—

1. *Tapeworms* of various kinds, the three most common ones being transmitted respectively by beef, pork, and fish, the meat in either case being “measly.”

2. *Trichina*, a species of larval worm which infests the muscles, causing great suffering, and, at times, death. It is conveyed to man by eating underdone pork.

3. *Tubercle bacilli*, the cause of tuberculosis or consumption.

4. *Bacteria causing some animal diseases*, which may reproduce the same diseases in man, or may merely cause intestinal trouble. Paratyphoid, a disease resembling typhoid, may be so produced.

5. *Ray fungus*, the cause of actinomycosis or lumpy jaw in cattle.

6. *Typhoid bacillus*, the cause of typhoid fever, may be conveyed in oysters or other shellfish coming from polluted waters.

7. The *bacillus of anthrax*, the cause of a very fatal disease.

8. The *bacillus* causing “*tremblcs*” in cattle and “*milk sickness*” in man.

Some diseases fatal to food animals do not affect man, and the flesh of animals dead of such diseases can safely be eaten ; but in general it is safe and wise to avoid all such meat, as, though the disease killing the animal may not affect man, chemical changes may have occurred in it to give rise to poisons, and ptomaine poisoning or “meat poisoning” may result. *Ptomaine poisoning* may

38 THE RECRUIT AND HIS ENVIRONMENT

also result from the action of bacteria contaminating the meat after death, or even after cooking, and causing changes in it. Such instances are seen when poisoning occurs from cold meat or hash cooked some days before use, or from canned meat taken from imperfect tins. When the ptomaine or other poisonous products of bacterial action are once formed in the meat, they often may not be destroyed by cooking, and all such meats are dangerous.

Some few kinds of meat, principally fish, are poisonous in themselves.

Meats are preserved for use in a variety of ways, only a small percentage of those furnished the army being used so soon after killing as not to require some form of artificial preservation. Of these methods of preserving, *cold* is the most satisfactory and best keeps the meat unchanged, and by this means it may be kept for months. It is applied so as to freeze or merely to "chill" the meat, and its only effects are to render the flesh more tender and to detract somewhat from its flavor. Frozen meat may be kept indefinitely, but that which is merely chilled does deteriorate in time, becoming soft, slippery, and flavorless. It should not then be used as food.

Cold cannot be depended upon to destroy the germs or parasites in meat, though keeping for a long time will destroy some of them. *Salting, pickling, and smoking* are much used in preserving meats, particularly beef and pork, and the results are very good. Ham, bacon, and corned or smoked beef are excellent and palatable meats for occasional use, but they are not adapted to long use to the exclusion of fresh meats, and when they are so used the health is apt to suffer. These processes

tend to free the meats from parasites, but cannot be depended upon to do so, and the eating of raw ham is a dangerous procedure. Canning, with its attendant sterilization of the meat, preserves all of its nutriment and destroys all parasites, but it melts the fat, gelatinizes the gristle, and softens the muscle, so that the meat comes from the can less attractive to the eye or the palate, but otherwise good and valuable.

Preservation by means of drugs and chemicals, other than those used in pickling or salting, such as boric acid or formalin, is forbidden, and meat so preserved may cause sickness; but for emergencies and occasional use it would not ordinarily do so.

It is not desirable here to consider in detail the various methods of cooking meats, but the company officer should exercise a certain amount of supervision in the matter, to see that his men's digestion is not injured by an excessive use of fried meats, that their appetites are not impaired by poorly prepared or poorly served roasts, stews, and hashes. A company mess is often good or bad as the cook knows, or does not know, how to handle, prepare, utilize, and serve meats. Rather common faults in company cooks are as follows:—

Roast or baked meat is overcooked and dry, is taken from the oven an hour or two hours before meal-time and sliced, and is served with hot, greasy gravy to make it warm.

Tough ends, scraps, and bones are not utilized in making soup stock, as they should be.

Excessive fat is not rendered and utilized.

Hashes and stews are served with unnecessary frequency, and become monotonous.

Lack of imagination or lack of knowledge on the

40 THE RECRUIT AND HIS ENVIRONMENT

part of the cook and mess-sergeant result in lack of variety and attractiveness of the mess. Intelligent supervision of the mess by the company commander, the preparation of bills of fare in advance, so as to insure variety, and the study of good cook-books will obviate this.

The principal carbohydrate food is bread, and its importance in the mess is scarcely, if at all, secondary to that of meat. The ordinary supply of

Bread bread is issued as such, and its quality depends largely on the capacity of the post baker, notwithstanding his proneness to credit any defects in it to the flour, the yeast, the hops, or the oven. The materials supplied by the government are, with rare exceptions, excellent, and the failure to produce good bread is usually to be attributed to either the ignorance, the carelessness, or the uncleanness of the baker. With the proper training of the bakers such failures will seldom be seen. Company cooks are permitted, and should be encouraged, to bake *biscuit* and *muffins* occasionally. They are somewhat more difficult of digestion than good issue bread, but are much liked and add to the attractiveness of the mess. Stale bread (that is, more than twenty-four hours old) is somewhat more digestible than fresh or hot bread; but during the process of staling it should be kept covered and protected from dust, and should not be unnecessarily handled, nor by any but clean hands; otherwise bacteria and moulds find lodgment and growth upon it and may produce disease. Toasting improves some breads and adds to their digestibility, but it is not often done in companies.

Pastries and *cake* are also bread, but are less easily

digestible than the ordinary forms. They add much to the attractiveness of a mess and should be served occasionally. The same remarks apply to the various kinds of batter-cakes.

Cornbread also makes a pleasant change from the monotony of baker's bread, and, as it is cheap, palatable, and nourishing, its use is to be encouraged.

Corn-meal mush, with butter or milk, is also relished, and when cold may be sliced and fried.

Hard bread is issued only for field use. Good teeth are necessary for its utilization as issued, but it may be softened in hot water or coffee without impairing its value.

A measure of economy, sometimes ignored, is the utilization of crusts and fragments of stale bread in puddings, as meat-stuffing, and in other ways. Such practices add to the variety and attractiveness of the mess, and allow savings to be made in other directions. To throw away such crusts because sugar and eggs are necessary in puddings and cost a little extra, is wasteful and wrong.

Vegetables supply all classes of foods, and are very important in the soldier's mess for this reason, and because they serve to impart variety and attractiveness to it. The most important vegetable issued (except wheat) is the *potato*, which consists of almost pure starch. It may be cooked in a much greater variety of ways than is usual in most companies, and the company commander should see that such is the case. In whatever way they are prepared, potatoes should be cooked until soft or mealy, and should be served hot.

**Vege-
tables**

Rice is another valuable starchy food, and in some

part of the cook and mess-sergeant result in lack of variety and attractiveness of the mess. Intelligent supervision of the mess by the company commander, the preparation of bills of fare in advance, so as to insure variety, and the study of good cook-books will obviate this.

The principal carbohydrate food is bread, and its importance in the mess is scarcely, if at all, secondary to that of meat. The ordinary supply of

Bread bread is issued as such, and its quality depends largely on the capacity of the post baker, notwithstanding his proneness to credit any defects in it to the flour, the yeast, the hops, or the oven. The materials supplied by the government are, with rare exceptions, excellent, and the failure to produce good bread is usually to be attributed to either the ignorance, the carelessness, or the uncleanness of the baker. With the proper training of the bakers such failures will seldom be seen. Company cooks are permitted, and should be encouraged, to bake *biscuit* and *muffins* occasionally. They are somewhat more difficult of digestion than good issue bread, but are much liked and add to the attractiveness of the mess. Stale bread (that is, more than twenty-four hours old) is somewhat more digestible than fresh or hot bread; but during the process of staling it should be kept covered and protected from dust, and should not be unnecessarily handled, nor by any but clean hands; otherwise bacteria and moulds find lodgment and growth upon it and may produce disease. Toasting improves some breads and adds to their digestibility, but it is not often done in companies.

Pastries and *cake* are also bread, but are less easily

digestible than the ordinary forms. They add much to the attractiveness of a mess and should be served occasionally. The same remarks apply to the various kinds of batter-cakes.

Cornbread also makes a pleasant change from the monotony of baker's bread, and, as it is cheap, palatable, and nourishing, its use is to be encouraged.

Corn-meal mush, with butter or milk, is also relished, and when cold may be sliced and fried.

Hard bread is issued only for field use. Good teeth are necessary for its utilization as issued, but it may be softened in hot water or coffee without impairing its value.

A measure of economy, sometimes ignored, is the utilization of crusts and fragments of stale bread in puddings, as meat-stuffing, and in other ways. Such practices add to the variety and attractiveness of the mess, and allow savings to be made in other directions. To throw away such crusts because sugar and eggs are necessary in puddings and cost a little extra, is wasteful and wrong.

Vegetables supply all classes of foods, and are very important in the soldier's mess for this reason, and because they serve to impart variety and attractiveness to it. The most important vegetable issued (except wheat) is the *potato*, which consists of almost pure starch. It may be cooked in a much greater variety of ways than is usual in most companies, and the company commander should see that such is the case. In whatever way they are prepared, potatoes should be cooked until soft or mealy, and should be served hot.

Vegetables

Rice is another valuable starchy food, and in some

42 THE RECRUIT AND HIS ENVIRONMENT

parts of the world it is the main article of diet; but in our country it is much less used and less desired than bread or potatoes. It is most popular with soldiers as a pudding, though it may be prepared in a variety of ways. It should be used to give variety and served in attractive forms, cooked with tomatoes or something else to take away its insipidity.

Corn is also to be classed as a starchy food. In addition to the uses discussed, it may be eaten in the whole grain, in its green, canned, or ripe, dry state. Green or canned corn is not so nutritious as the ripe, but it is much liked and should be used for the sake of variety. Ripe corn in the whole or the broken grain, as hominy or grits, is not so generally liked as the green, but it is very nutritious, and should be used occasionally as a principal dish.

Macaroni and other paste preparations are valuable foods, and admit of a variety of presentations, so that their use should be encouraged.

Onions are a staple article of issue, and are much used in the army. Their nutritive value is much less than that of the articles just discussed, but their flavor and the variety of ways in which they may be used make them very valuable. Monotony should be avoided in their use.

Tomatoes, likewise, have relatively little food-value, but stand high for their flavor and the variety they afford. When they are served cooked, it should often be with some otherwise insipid food, such as rice, macaroni, or stale bread.

Beans and *peas*, though possessing considerable starch, are the principal sources of vegetable proteid. They are both very attractive and palatable in the green state,

and, though less nutritive than when ripe, add much to the mess. When dried, they are particularly valuable in making soup (for use in which ham-bones and rinds should always be saved), and when served boiled or baked, with bacon. They are, however, somewhat difficult of digestion, and their cheapness should not lead to too frequent use.

Beets are another valuable source of food, as they contain much sugar. They are also cheap, and are relished by most men.

Most of the common vegetables, as cabbage, greens, carrots, spinach, radishes, and cucumbers, have little nutritive value, but they add greatly to the attractiveness of the mess and are also valuable in preserving health and preventing scurvy, and their use should be encouraged.

Salads are not used in company messes as much as they should be. They afford pleasing variety, and may be made to constitute an important dish, as when composed principally of potatoes or salmon.

Salads

As issued in company messes, desserts are nearly always very simple, and such is necessarily the case. When practicable, a dessert should be furnished once a day, and it may be made to use up otherwise unattractive articles, such as rice or stale bread made into puddings. Pies made of fresh or dried fruits are easy to make and serve, are inexpensive, and well liked.

Desserts

Milk and its products constitute important articles of food, though not entering so largely into the soldier's dietary as they might. Milk contains all classes of food, — proteid, fat, carbohydrate,

Milk

44 THE RECRUIT AND HIS ENVIRONMENT

water, and salts, — though not in the proportions desirable in maintaining adult life. Fresh milk is both more palatable and more wholesome than the canned or condensed article, if its purity and freedom from disease-producing germs can be assured, but unfortunately this is not always the case. It should be borne in mind that many serious diseases may be conveyed in milk, among them tuberculosis, typhoid fever, diphtheria, and scarlet fever, — some of the diseases most dreaded in military life. Milk sickness, a disease now rarely seen, was also conveyed in that way; while for many years the Mediterranean garrisons of the British army suffered very seriously from Malta fever, now known to have been transmitted by means of goats' milk.

In case of epidemic of any of the above-named diseases, the milk supply should be investigated, and this is particularly suggested by outbreaks of typhoid showing an explosive character, that is, epidemics in which many cases occur almost simultaneously. Because of this possibility of disease-production, milk whose origin and condition are not above reproach should be used in companies and post exchanges only in cooking or after boiling, and in camp or on the march men should be cautioned to the same effect, and warned against purchasing it by the glass from dealers or peddlers. Condensed or evaporated milk is more generally used in companies than the fresh article, and it is on the whole probably safer, as the process of condensation and the subsequent prolonged storage tend to destroy disease-producing organisms. A can of such milk, when once opened, should either be used promptly or kept cool and protected from dust until used. Boiling water should be used to dilute it, to kill germs

that may have settled and grown on it after the opening of the can.

Buttermilk is a valuable and pleasant drink, and where it can be obtained is much relished by the men.

Good *butter* is made from good cream and is everywhere highly appreciated; but unfortunately the good quality is not always obtainable, and much very poor butter is used in company messes. Oleomargarine is little inferior to butter in nutritive qualities and is to be preferred to poor butter, except that a prejudice exists against it in this country and men are apt to avoid its use. It has an advantage over butter in that it is not so likely to act as a carrier of disease germs, notably tuberculosis.

Both are valuable fat foods as well as relishes, and should not be regarded merely in the latter light.

Cheeses are very concentrated forms of food, containing about one third proteid and one third fat, though the different varieties differ in the percentages of these. They make agreeable and valuable additions to the mess, but are seldom used in large quantities because they are apt to be difficult to digest, and to cause constipation. Ordinarily they are used in companies rather as a condiment than for their food-values, though at times they may constitute an important part of a meal. In cooking, cheese is used principally with macaroni. Milk and cheese are liable to certain changes which cause the development of poisons, one of these, known as tyrotoxicon, being very powerful and producing severe symptoms or death. The ordinary souring of milk is not in itself dangerous, but may be beneficial in some respects. The justification for regarding sour milk with suspicion lies in the fact that the conditions

favoring the development of the bacteria causing the souring may also have favored the development of disease-producing organisms. Milk should be preserved either by refrigeration, to prevent the growth, or by heating, to kill, the contained bacteria.

For preserving other foods than meats and milk, refrigeration, canning, and drying are the three great commercial methods. Cold storage may preserve fruits and vegetables, as well as meats, eggs, and milk, practically unchanged for weeks, and immense quantities are so preserved and sold. Of even greater importance, however, is the process of canning, whereby the articles are cooked, sterilized, and hermetically sealed in cans or jars, to be preserved for months or years. This industry has increased enormously, and is now applied to almost all of the articles of food that have been discussed herein, and to a great number of others not so discussed.

The results obtained are excellent, and such foods are carried to all parts of the world and enjoyed.

Canned foods pall after a time, though, and should never, if it can be avoided, be used to the exclusion of fresh food. Otherwise nutrition may be impaired, and even scurvy may result. This may be because the heat necessary to preserve them destroys some content that is thus far intangible. The fear of metallic poisoning from canned foods is thought to be exaggerated, and the probability of such poisoning to be remote, except where lead is used, and that is infrequent and does not apply to the great bulk of canned foods used by the soldier.

Drying as a method of preservation is applied to fruits and vegetables, as well as to meats and fish. Desiccated vegetables find some use in the army, par-

ticularly in Alaska, where they constitute a real blessing, but their use is not general.

Dried fruits, however, are everywhere used and with very general satisfaction. Dried apples, peaches, prunes, and raisins are issued or sold by the Subsistence Department, and make valuable and pleasing additions to the mess, where they are served stewed or in puddings and other desserts.

Sweets are much enjoyed by the men, and are valuable for their fuel-value, and also, probably, as lessening a craving for alcohol. Syrup and jam are articles of the ration that are much enjoyed, while the sale of candy by the Subsistence Department shows that its worth is recognized. As stated before, an excess of sugar causes dyspepsia and should be guarded against.

Sweets

Coffee, tea, and cocoa are mild stimulants, but may almost as well be classed with condiments. Of the three, coffee is the only one extensively used in our service, and it is often poorly made and poorly served. The green issue coffee is of excellent grade, and if properly roasted and ground, and used fresh, is superior to most higher priced coffees obtained elsewhere, ready roasted. The preparation of good coffee is simple and easy, but most company cooks do not practice it. The company commanders should make sure that, in addition to proper steps in preparation, the following are observed :—

Bever- ages

The coffee must not be long boiled.

The coffee-pot must be emptied and cleaned after each meal, and no coffee grounds should be used twice.

The sugar and milk, if the latter be served, should be put in by the user and not served alike to all.

48 THE RECRUIT AND HIS ENVIRONMENT

Enough coffee should be used to give strength and flavor to the beverage, and then but a moderate amount of it should be served to each man.

In the field, where the water is of a doubtful character, the issue of large amounts of weak coffee is justified, as tending to reduce the amount of unboiled water used ; but in posts this is not necessary.

Cooks will insist that the men prefer their coffee weak, long-boiled, and ready sweetened, just as they will say that they prefer roast beef overcooked, dry, and swimming in greasy gravy ; but inasmuch as most men in civil life like to exercise a choice in such matters, it is thought that a like privilege might be appreciated in the service.

It is also thought that perhaps the use of tea in the service might become more general if the beverage were properly made, and not boiled or steeped until a strong, black tannin solution results.

The following is a bill of fare for one week, which, it has been demonstrated, can be supplied with little extra cost beyond the savings on the ration, provided that the cook is capable and careful and the mess-sergeant intelligent. Slight variations from week to week and to suit the seasons will permit such a bill of fare to be used for a long period and give satisfaction.

MONDAY

Breakfast : Stewed Fruit, Fried Eggs, Bacon, Bread, and Coffee.

Dinner : Roast Beef, Steak, or Meat Balls ; Baked Potatoes, Squash or Turnips, Stewed Tomatoes ; Pie or Cake.

Supper: Cold Sliced Beef, Fried Onions or Tomatoes, Bread and Jam, Tea or Coffee.

TUESDAY

Breakfast: Bananas or Fresh Fruit, Liver or Kidney with Bacon, Bread, Butter, and Coffee.

Dinner: Beef Steak or Hamburger Steak; Potatoes, Radishes, or Onions; Pudding, Bread, and Coffee.

Supper: Cold Beef, Fried Potatoes, Bread and Jam, Tea or Coffee.

WEDNESDAY

Breakfast: Stewed Fruit, Fried Mush and Molasses, Coffee.

Dinner: Boiled Ham; Boiled Potatoes, Boiled Cabbage; Pudding, Bread, and Coffee.

Supper: Cold Boiled Ham, Bread and Jam, Butter, Tea or Coffee.

THURSDAY

Breakfast: Stewed Fruit, Puffed Rice, Wet Hash on Toast.

Dinner: Roast Beef, Steak, or Meat Balls; Baked Potatoes, Corn, Squash, or Turnips; Pudding, Bread, and Coffee.

Supper: Cold Beef or Hash, Bread and Jam, Fried Onions or Tomatoes, Tea or Coffee.

FRIDAY

Breakfast: Stewed Fruit, Oatmeal with Milk, Biscuits with Syrup and Coffee.

Dinner: Baked Fish with Sauce; Boiled Potatoes and Onions; Bread, Butter, and Coffee.

50 THE RECRUIT AND HIS ENVIRONMENT

Supper: Salmon or other Fish Salad, Bread and Jam, Tea or Coffee.

SATURDAY

Breakfast: Stewed Fruit, Bacon, Rice or Corn Flakes, Toast, and Coffee.

Dinner: Baked Beans with Pork ; Pickles, Stewed Tomatoes ; Bread and Coffee.

Supper: Cold Beans or Bean Soup, Pickles, Tomato Catsup, Bread, Butter, and Coffee.

SUNDAY

Breakfast: Stewed Fruit, Oatmeal with Milk, Hot Biscuits, Coffee.

Dinner: Roast Pork, Veal, or Mutton ; Browned Potatoes, Baked Squash ; Apple Sauce, Pie or Cake.

Supper: Cold Meat, Bread and Jam, Tea or Coffee.

THE HYGIENE OF THE KITCHEN

From the hygienic and many other points of view the company kitchen is one of the most important places in the post, and the company cook a very important person.

These two bear a more direct relation to the guard reports, sick reports, and the general efficiency of the command than at first appears ; while the subjects of alcoholism and desertion are especially involved.

It is therefore an important part of the company commander's duties to see that both are as good as can be obtained. He should see, as stated before, that his cook is capable of preparing the food in attractive, satisfying, and digestible forms, that he is economical, a good manager, and can utilize the ration in a variety of ways and supplement it judiciously. But the cook may fulfill

these requirements and still do much harm by causing or spreading disease. Therefore he and his company officers should know something of the hygiene of the kitchen.

The primary and most important rule as to cook, kitchen, and the contents and surroundings of the latter, is cleanliness. All parts of the barracks and all persons therein should be clean, but the

Cook

kitchen and the cook should excel. The cook should bathe often, change his clothing frequently, always wash his hands after a visit to the toilet, after handling anything dirty, and before handling foods. He should always have a clean hand-towel in his kitchen, and not use for the purpose either a dirty rag or his dish-towels. He should wear white aprons, caps, and clothing, and change them when soiled. His finger-nails and his hair should be kept short and clean, and a nail-brush should be in the kitchen for use and should be scalded daily. A cook who has had typhoid fever, or, rarely, one who does not know that he has had it, may, though apparently in good health, be excreting and distributing typhoid bacilli; and a typhoid epidemic limited to a company should always cause this possibility to be investigated. Men who are dirty in their habits, who have syphilis, tuberculosis, or other infectious diseases, or who are persistent in unsanitary or careless methods of preparing foods, should not be allowed to cook.

The kitchen should be proof against the most rigid inspection for dirt. Some disorder and litter necessarily attend the preparation of food; but this can usually be cleared up almost immediately, and it should not remain any longer than necessary. The meat-block, bread-

52 THE RECRUIT AND HIS ENVIRONMENT

boards, carving-tables, and all utensils should all be inspected daily and all kept clean.

The ice-chest and milk-cans should receive special scrutiny, and should be clean inside and out, as evidenced to the eye, the nose, and the white glove.

Foods like cheese, codfish, and bacon, that possess a strong odor and do not readily spoil, should be kept on shelves or in boxes, protected by wire gauze, rather than in the ice-chest. Canned meats, fruit, and vegetables should be used as soon as possible after they are opened, or emptied into clean dishes, and not left about in the cans. All food should be used before it becomes mouldy, sour, or decomposed, or should be thrown away, though a slight and recent growth of mould on the surface of ham, cold-storage beef, bread, jam or other sweets does not necessarily injure the whole, and may be removed.

The kitchen sinks should always be kept clean, and should be well trapped. Care should be exercised that grease, crumbs, and fragments are kept out of them, that the pipes may not become clogged. Kitchen waste, the scrapings of food from plates, and unused fragments should be thrown into covered cans, which should be emptied and cleaned once daily, and should not serve either as a breeding or a feeding place for flies, roaches, rats, or other vermin. Particular care should be exercised to exclude all such from the kitchen at all times. For this purpose cleanliness and screens should suffice, but if fly-paper, roach-poison, and rat-traps are needed, they should be supplied.

The *kitchen police* is usually supplied by roster from the company, but at times the work is given as a punishment. This is undesirable. Instead of poor men,

good ones should be put on kitchen police, and the work should carry with it immunities and privileges to make good men desire it.¹

¹ For fuller and more comprehensive information on the preparation of foods and mess-management, the reader is referred to the book entitled *Handling the Straight Army Ration and Baking Bread*, by Captain T. R. Holbrook and Color-Sergeant Patrick Dunne, 5th U. S. Cavalry, and to the course of instruction given at the Army Training School for Bakers and Cooks at Fort Riley, Kansas, both of which are highly commended. No opportunity to have a cook trained in the school should be neglected. The Army Cook Book and the Subsistence Manual are also very important sources of information on this subject.

CHAPTER IV

THE HYGIENE OF THE BARRACKS

NEITHER the enlisted man nor his immediate commander has very much choice in the matter of barracks. They are assigned to duty at a post where certain barracks are provided, and must of necessity make use of them. In one post the barracks may be new, large, and built with a view of affording all possible advantages in the way of comfort, convenience, and sanitary arrangements. In another post they may be old, small, and apparently built without a thought of any of the advantages mentioned. In the one the capacity of the barracks may greatly exceed the size of the garrison, in the other, the reverse may obtain; but in nearly all instances the conditions are controlled by military necessity and not by the company commanders. It is therefore considered that the proper scope of this chapter is to indicate what is desirable, and to assist in making the nearest practicable approach to it under existing conditions.

The buildings are apt to be of almost any material, from brick, stone, or concrete, to nipa and thatch.

Build-ings Usually they are of substantial materials and well constructed, unless designed for merely temporary use in warm countries, when light board-structures or those of bamboo and palm are used. In the latter class, cheapness of cost is always a consideration; but it is not to be carried to the extent of making the structures unsanitary. Suf-

ficient room should be provided, adequate and convenient supplies of good water, proper kitchen arrangements, and protection from mosquitoes, flies, and other insect pests. The structures should be sufficiently raised above the ground to prevent dampness, if possible, and ample provision should be made for the care of excreta, garbage, and waste water.

In permanent barracks of the newer type, provision has usually been made for the sanitary requirements, and the conditions are satisfactory where overcrowding does not obtain. Barracks of concrete, stone, brick, or wood may all be satisfactory if other considerations are met. One of the first of these is location and exposure, and the desirable locations and facings vary with climate and local conditions.

In general, it is well to get much exposure to the sun in cold climates, and this is best accomplished by having the four corners of the main building point in the cardinal directions. In such places, however, the avoidance of prevailing winter winds may be of even greater importance, and may dictate a location. In any event, both sun and wind must be considered in their sanitary as well as their comfort-bearing capacities, the former as a great aid in cleanliness and disinfection, the latter as a promoter of ventilation. In any climate, the structures should be dry as to both site and walls. Damp cellars and surroundings may be avoided by selection or by grading and drainage. Damp walls are avoided by having a damp-proof course, as of slate or concrete, above the foundations, by double walls with air-spaces, by good ventilation, and, where excessive or hard-driven rains obtain, by waterproofing with paint, etc. If the walls are continuously of porous

brick or stone to the cellar, or to the surface of the ground, they will be continuously damp and chilly, in most climates, and only the insertion of a damp-proof course will correct the condition. If other considerations, such as size, ventilation, heating, or plumbing, are unequal, they should be allowed more weight, in governing the choice of barracks, than the structural material; if all other considerations are equal, the structural feature should govern choice, usually in the following order: concrete, brick, stone, wood.

The buildings should be sufficiently large to provide abundant squad room, recreation or day rooms, store-rooms, kitchen, pantry, and dining-rooms, **Rooms** small rooms for noncommissioned officers, tailor's, barber's, and cobbler's shops, a workshop for the mechanics, and ample water-closet and bathing facilities, unless a part of these are provided elsewhere.

The *squad-rooms* are the matter of greatest consideration, and they should be given first thought in selecting barracks. They should always be as large as it is possible to obtain, keep clean, and heat, and should provide a minimum of six hundred cubic feet of space and sixty square feet of floor-room for each man, exclusive of the room occupied by wardrobes and lockers. They should be well lighted both by windows and artificial light, preferably electric. The window-space necessary will vary with climate and location, but it should insure plenty of breeze in the tropics, and plenty of light in northern climates. If a good system of ventilation, other than by windows and doors, has been installed, so much the better, and it should be carefully studied until understood, and then kept in

maximum working order. In all barracks, ventilation and heating are closely related, and should receive the company commander's personal consideration. If it be practicable, it is a great help in ventilating most barracks to have a small open-grate fire going all winter, but there is usually no provision for it. The air in the squad-rooms should at all times be sweet and fresh, and free from the "closeness" and staleness that is particularly apt to obtain in the early morning hours of winter nights. The means of insuring this will vary in different places, but there is always one expedient that may be resorted to, — raising the lower window-sashes and fixing boards beneath them, with nails if necessary, so as to allow the air to enter in an upward direction between the two sashes. Inspections should be made occasionally, at the times indicated, to see that the air is good; while all arrangements to promote ventilation should be inspected frequently to see that they are in working order and have not been interfered with. In case interference is found, it is well to detail a man in each room to be responsible for their correct maintenance. All squad-rooms should be opened and thoroughly aired each day, in addition to the constantly working arrangements.

Always in the tropics, and during warm weather in other climates, all windows should be provided with screens to exclude flies and mosquitoes. If such is not the case, bed-nets must be used at night.

In cold climates double or storm doors and windows may be provided. They make the rooms more comfortable and easier to heat, but should not be allowed to impair the ventilation.

Water-closets, bath-rooms, and wash-basins should

58 THE RECRUIT AND HIS ENVIRONMENT

be entirely away from the squad-rooms, in which no plumbing should show. Articles of food, dirty and wet clothing, and useless trappings should be excluded as far as practicable. Floors should be sprinkled frequently with wet sawdust, and swept so as to create as little dust as possible; and at least once a week the entire room and its contents should be so thoroughly cleaned as to pass the most rigid inspection, no dirt, no dust, no disorder, no vermin being anywhere present. This inspection will be made on Saturday, and should be preceded by an outdoor airing of bedding and change of bed-linen on Friday. At the inspection all wardrobes, drawers, lockers, and boxes should be opened, and their contents exposed to view. The small spaces and corners about such articles, the space behind or under radiators, the tops of wardrobes and shelves, should all be examined, and if any man seem unclean, his linen and person. It is not thought wise to subject intelligent and self-respecting men to such personal inspections as a matter of routine.

The *day-rooms*, or recreation-rooms, should be suitable for their purpose, should have good light, good air, comfortable heating, and such provision in the way of chairs, tables, reading matter, billiard and pool tables, and other facilities for amusements, as will conduce to the interest, instruction, and amusement of the men, or such, at any rate, as can be obtained.

These rooms should be kept clean and neat at all times, free from dust, foul air, dirt, and vermin. The care should be such as to enable them to pass the same general inspection as squad-rooms. Smoking will be indulged in, in recreation-rooms, at most times of day, but the rooms should be opened and aired frequently.

Spitting will also be indulged in, and cuspidors containing water should be provided in sufficient numbers to prevent spitting on the floor, and any man then doing so should be punished. The cuspidors should be emptied and scrubbed daily.

Pictures and ornaments may be permitted to some extent, to detract from the bareness and add to the cheerfulness of the room; but, in general, unnecessary dust-collecting articles should be excluded.

The *dining or mess room* should be sufficiently large to seat comfortably at table at least ninety per cent of the command. A certain number of men will necessarily be absent from each meal, — on guard, in the kitchen, on pass or furlough, and for other reasons, — but provision should be made for the maximum number that may be present. Lighting, heating, ventilation, and screening are all important here as in other rooms. Screening, particularly, should be carefully provided, and every means employed to keep the place free from flies. The cleanliness of this room should be as nearly perfect as possible, and while it is permissible to have the room attractive, dust-gathering trappings should be excluded. The mess-tables are used uncovered, or covered with white oil-cloth. The former is the preferable method, as it insures thorough cleaning and scrubbing, if the tables look clean. An oil-cloth can be made to look almost or quite its best by a small amount of indifferent rubbing, though it really may not be free from grease and dirt.

As said before, the men should be required to observe the decencies at table, and they are more apt to do so if the decencies are observed toward them. The table should therefore be orderly and attractive, though

60 THE RECRUIT AND HIS ENVIRONMENT

plain, the dishes and utensils should be perfectly clean and arranged in a proper manner, the food should be served in an orderly, attractive, and appetizing way, and with a sense of fitness. While parlor manners may not be expected in the barracks, gross indecencies of manner, speech, or action should be prevented. The room and table should be well cleaned after each meal, and the dishes and utensils should be carefully inspected daily and their perfect cleanliness assured. Once a week, or oftener, the tables and floors should be thoroughly scrubbed with lye and water, and vinegar-cruets, salt-cellars, mustard-pots, etc., well washed and filled. In warm weather, fly-paper and, if necessary, occasional fumigation with pyrethrum fumes, should be used to supplement the screens.

No man should be allowed to eat with unclean hands, and it should soon be a matter of habit for all men to come to the table clean as to hands and face, and neat in appearance. Leisurely eating and thorough mastication should be encouraged.

After the squad-rooms the *kitchen* is probably the most important room in the barracks. Its importance has been discussed elsewhere, but may be emphasized here. It should if practicable be a large room, well lighted and well ventilated, opening on a large porch, and connecting with store-rooms in the cellar or elsewhere convenient, and with a roomy pantry, with abundance of shelving and drawers, also well lighted and ventilated. Good tables, sinks, running water, ample drainage, sound and smooth meat-block, an amply large range, and abundance of good kitchen furniture and utensils should be provided. A large ice-chest and some screened shelving should be near at hand. It is

desirable, if practicable, to have a large hood over the range, to catch and conduct away, through a ventilating shaft, the odors, vapors, and smoke arising from the cooking. All of these articles should be frequently inspected to make sure that they are always cleaned as soon as possible after use, and never put away in any other condition. The cook, above all men in the company, should live, think, and dream cleanliness, and he should be held most strictly to account for shortcomings in that line. The kitchen police should be rewarded for efficiency in such a way as to make the detail desirable, and in this way the kitchen should be made an object-lesson and a school of cleanliness. The inspection of the kitchen should include the pantry and ice-chest, should take note of, and remedy, waste or poor management, and should always embrace inquiries concerning, and search for, flies, roaches, mice, and other vermin. Cats, dogs, and other pets, as well as men not there on duty, should be excluded at all times. Garbage-cans should be outside of the kitchen, should be emptied at least once a day, and then cleaned inside and out; otherwise they speedily become a foul nuisance.

The general remarks as to the lighting, heating, ventilation, and cleanliness apply to the offices and the shop-rooms for the baker, tailor, and cobbler, and the inspection of these latter should not be omitted, otherwise they are especially apt to form accumulations of scraps, dirt, and dust; and they are especially favorable distributing points for vermin.

The company barber should be required to keep his brushes, combs, razors, and utensils thoroughly clean, and to sterilize them frequently. He should be obliged to keep an abundance of clean towels and aprons, and

62 THE RECRUIT AND HIS ENVIRONMENT

to be liberal in their use. He is not apt to be so of his own volition. He should be instructed not to shave or cut hair for men with skin diseases without first getting the surgeon's assent. Should his own skin or hands show disease, he should consult the surgeon.

Store-rooms do not need light, air, and heat in such liberal amounts as do living-rooms, but they should be dry, free from vermin, and should answer the purposes for which they are required.

Bath-rooms should be well drained, well ventilated, and well lighted, and when such is not the case every effort should be made to correct the condition. Lighting is facilitated and cleanliness promoted by having walls, doors, and partitions painted white. Where tubs are installed, they should be thoroughly washed and cleaned after each use, and the floor should be kept clean at all times. Each man should have his own soap and towels, and should take them to and from the tub with him. When tubs are not sufficiently numerous to accommodate the men as they wish to go, they should be detailed by roster to bathe, at least once or twice a week, and as much should always be required of them. Shower-baths are much more economical of both time and water, and are installed at most posts. They, as well as tubs, should have both hot and cold water connections, be frequently inspected as to condition, and be kept scrupulously clean. The drain openings should not be allowed to clog with soap and hairs, and the entire bathroom should be scrubbed at least once a week, and as much oftener as is necessary to keep it clean and sweet-smelling, and free from mustiness. Gratings should be lifted daily if necessary and the space beneath flushed and scrubbed. The same general remarks apply to the

bowls for face and hand washing. The stoppers for these, and the chains holding them, should always be clean, the outlet openings likewise, and watch should be kept to prevent the formation of a film or coat of soap and dirt inside the bowls. The throwing of tobacco, paper, cotton, and dressings into bowls, tubs, or bath compartments should be forbidden and prevented. The man detailed in charge of the bath-room should be cleanly and trustworthy, and should have authority to enforce the rules.

The water-closets are frequently in close proximity to the bath-room, and if both are kept clean there is no objection to it. Flush-closets and urinals are installed in practically all posts now, and where such is not the case one of the various expedients for disposing of excreta in the field will have to be used, and reference is made to their description. Flush-closets should be kept clean at all times, the bowls scoured, and the seats wiped off occasionally with a damp cloth. The floors about them should always be clean, and the room should be free from odor. Standing on the seats, defacing or defiling them, throwing stiff paper, matches, and other things likely to cause obstruction, into the bowl, and writing obscenity on the compartment walls should be punishable offenses. When a closet gets out of order in any way, it should be put out of use until repaired.

It is to be borne in mind that the use of carbolic acid and chloride of lime is but rarely necessary about flush-closets, and then not for the purpose of destroying odors.

Constant bad odors indicate lack of care or poor plumbing, or both, and should be corrected. When concealed by other odors, such as those of chlorine or carbolic acid, they may not be noticed, but the evil

64 THE RECRUIT AND HIS ENVIRONMENT

condition of which they gave warning is unremedied. Tissue paper is issued for toilet purposes, and no other should be permitted to be used, as being apt to scratch or irritate the anus, and to clog the drain. By signs in the room, and by verbal instruction, the men should be taught to wash the hands after every visit to the closet. It should also become a habit for them to inspect their feces before flushing the closet, as disorders may thus be brought to their attention, and remedies sought. The passage of blood, pus, large amounts of mucus, or worms should prompt early consultation of the surgeon. The closet-rooms should be screened, and should be floored with cement to permit flushing, and to do away with cracks and crevices. Roaches and flies should be excluded. Each is indicative of unsatisfactory conditions. *Urinals* are ordinarily flush-bowls, for use by one person at a time, and should be sufficiently numerous to meet the demands of early morning and of the evening. The floor and walls immediately about them should be of smooth, hard, waterproof material, such as slate, and should be washed thoroughly each day and then lightly wiped with a moist rag sprinkled with kerosene. Otherwise a crust of urinary salts is apt to form, and odors of decomposing urine to arise. When proper care and cleanliness are exercised, neither of these appear, and the use of deodorants is not necessary. The flushing of the bowls is usually automatic, and should be of sufficient frequency to prevent deposition of salts, though its action should be supplemented by washing and scrubbing of the bowl. If the outlet of the bowl is not already screened, a piece of wire gauze should be placed over it and held in place by a small stone or other weight, otherwise it may get clogged with bits of

cotton, cigarette-stumps, and other waste thrown into the bowl. The throwing of such articles into urine bowls should be punished, if detected, but detection is not usual.

It is neither desirable nor practical to have the men do their own laundry-work in barracks, but a couple of tubs, running water, and scrub-brushes should be provided, if possible, to enable them to do a certain amount of emergency work and cleaning.

The company commander should familiarize himself with the system of plumbing in his barracks, with a view both to detecting leaks and other faults, and to taking action for the institution of improvements where indicated, and likewise to fit himself to distinguish between poor plumbing and inefficient care of good plumbing.

The water-supply of the barracks should, if possible, be of water fit for drinking without boiling or other special preparation. If this is not the case, the men should be warned and the proper drinking supply carefully defined and located. The general supply should be abundant for all purposes, and should amount to one hundred gallons per man per day, though so much need not ordinarily be used, and the amount really used will depend on a number of varying factors, such as temperature and climate, the character of the work done by the men, the care used, economy exercised, and the condition of the plumbing.

Water-Supply

Where tub-baths are used, the consumption will be greater than if showers are used. While care should be exercised to prevent waste, it is usually false economy to hinder any reasonable use of water,

as in bathing, scrubbing, watering plants or grass, or laying dust.

The water should be piped to the kitchen, the bath-rooms, water-closets, urinals, wash-basins, furnace-room, and to such places in and about the building as may be necessary for police purposes or fire protection. All faucets, exposed pipes and fire-plugs, and other parts of the water-supply apparatus, should be frequently inspected for leakage and as to their working condition.

Except in the ways indicated above, the water-supply in barracks is not usually controlled by the company commander.

Good plumbing presupposes good materials, and such are usually furnished. Iron pipe should conduct the water-supply, and all joints should be tight. **Plumb-** Faucets are usually of brass, and with proper **ing** usage rarely need repairs, except new washers. Lead pipe is objectionable as being liable to cause lead-poisoning. Some water after a time will cause the pipes to fill partially or entirely with a deposit of mineral salts, or a growth of low vegetable forms, with a consequent lessening of the size of the stream. Such conditions, as well as advanced erosion, may necessitate renewal of the pipe. Kitchen sinks and wash-bowls should have their outlet-pipes and traps exposed and open to inspection, not closed in to become musty hiding-places for roaches. Kitchen sinks are usually made of galvanized or enameled iron or zinc, wash-bowls of enameled iron or porcelain. In any event, the surface should be smooth and capable of complete cleaning.

All should be effectively trapped to prevent the re-gurgitation of air or gas from the drains. Sinks should

preferably have grease-traps, which should be easy to open and clean.

All bath-tubs should be of enameled or porcelain-lined iron or of zinc, and closet and urinal bowls of enameled iron or porcelain. All should be thoroughly trapped, and ventilated on the ground side of the trap, by communicating with the ventilating pipe. No odor should come from them, and they should flush freely and effectively on all occasions. Whenever one of them is found at all out of order, flushing poorly or imperfectly, or emptying slowly, it should be put out of use until the fault is corrected. Such trouble, more often than not, comes from the use of newspaper as toilet-paper, or the throwing of matches and other improper objects into the bowls. Any leaking from a closet, urinal, or drain should be immediately corrected, as especially apt to cause disease. The maintenance of thorough cleanliness in the closet is incidentally the most ready way of learning of such defects, while the faults arising from its neglect are at times unjustly attributed to the plumbing.

Of the various systems of lighting in use in our barracks, electricity is by far the best. Both oil and gas consume and contaminate the air, produce dirt and much heat, may be blown out, and give relatively poor light. **Lighting** Electric lighting produces no dirt, neither consumes the oxygen, nor adds to the combustion products in the rooms, produces very little heat, cannot be blown out, and gives a brilliant, steady light that is very much better than either of the others. The danger of explosion is absent, and that from fires is reduced to a minimum by proper wiring. The lights should be sufficiently numerous to illuminate well all

parts of the building, while in the reading and recreation-rooms they should be numerous enough and placed low enough to make reading and writing practically as easy as in the daytime. Dull and flickering light, such as is often produced by gas or oil, is very trying to the eyes, and causes strain and fatigue; while the vitiation of the air by the combustion of these substances necessitates much more liberal ventilation, or causes suffering. Economy in lighting should be practiced by extinguishing lights when the necessity for their use has passed, as in the dining-room and kitchen after they are cleaned, in the squad-room late in the evening, thus having more for use in recreation-rooms.

As stated earlier, the subjects of ventilation and heating are so intimately connected that no consideration of the one is complete unless it includes the other. This is so for the reason that gravity is the most important factor in ventilation, and the weight of a given volume of air is less or greater as it is warm or cold. In other words, we influence or control gravity by means of artificial heat. The same force is the principal factor concerned in the diffusion of heat and heated air throughout barracks.

In summer, doors and windows can be kept open, and ventilation is seldom a very serious problem; but in winter or very cold weather, when doors and windows are closed and all the outside air is so cold that its impact causes discomfort, the case is different.

Open fires are suitable for heating single, small rooms, but not large ones, though the presence of an open-air fire aids in the ventilation of the latter as well as adds its cheer.

Stoves give more heat and less ventilation, for the

same amount of fuel used, than do open fires. They may be made to promote ventilation greatly by being surrounded by a sheet-iron cylinder or jacket, into which a fresh-air shaft empties at the bottom. The air is warmed as it enters, and rises into the room through the top of the cylinder, and later escapes by way of stove-door, window-cracks, and sundry other outlets, including outlet-shafts, if such are provided.

The inlet should be screened to keep out dust and dirt, and the top of the stove or of the cylinder should always be provided with an open vessel of water, to moisten the air by its evaporation.

Cast-iron stoves give off carbon monoxide, a poisonous gas, when red-hot, and may so do harm. The air used in the combustion of fuel is taken from the room and passes up the chimney, fresh air replacing it from the outside, and quite satisfactory ventilation may at times be so maintained, even with unjacketed stoves, but it is not wise to rely on such means alone, especially in crowded barracks. Slow-burning stoves, such as anthracite base-burners, exercise much less influence in this way than rapid-burning wood or soft-coal stoves.

Hot-air furnaces operate by heating fresh air, which is brought to them through a ventilating shaft from the outside, and which, when heated, rises through the tubes and passages to the registers, whence it passes into the rooms. One advantage of this method of heating is that, with increased heat, in very cold weather it sends up an increased amount of fresh air. This air, though, is usually very dry if the water-pan of the furnace is not carefully watched and kept filled. This is due to the fact that air doubles its capacity for absorbing moisture with each increase of 15° C. in its

70 THE RECRUIT AND HIS ENVIRONMENT

temperature ; so that outside air which at zero C. does not seem dry is excessively so when raised to 25° C. For this reason all systems of artificial heating should make some provision for moistening the heated air ; which otherwise causes discomfort and harm.

The method of heating just described is not very much used in our barracks, being regarded less favorably than the steam and hot-water systems.

Steam-heating is very satisfactory and is much used. In this system the heat from the fire is transmitted to water, which is made to boil, under pressure if necessary, and the heat is carried throughout the building by the steam, which travels through tight iron pipes that expand into radiators in the various rooms. It is a rapid method of heating, and one easily controlled by an intelligent man, and such a man should be detailed in charge of it. He should maintain an even, steady fire, enough steam to make the rooms comfortable, and keep the water in the boilers at a constant and proper level. Such plants usually show their maximum efficiency when a steam pressure of one or two pounds is maintained.

Water should be kept evaporating from the radiators, or, with a good man in charge of the system, steam may be allowed to escape from them to moisten the air.

Hot-water heating is somewhat similar, but the pipes are filled with water instead of steam. It is a very steady method of heating, but slower in either heating or cooling than steam. High and low pressure systems are used, only the latter in our service.

The air heated by the radiators is dried thereby, and provision should be made for moistening it.

When steam and hot-water heating are used, they are made to promote ventilation by having the fresh air enter the rooms through or under the radiators. It is thereby warmed and made lighter, and at once rises, to be later distributed through the room by gravity, convection, and diffusion. In order that it may not again be drawn through the radiators and redistributed, a line of less resistance should be created for it by ventilating shafts opening near the floor and leading to the roof, or by an open fire-place to answer the same purpose.

As there is also danger of fire in and about barracks, and as that danger is increased in cold weather, all chimneys, flues, shafts, stoves, furnaces, etc., should be thoroughly inspected and put into first-class condition in the late summer or early fall, before fires are started. Competent men only should be detailed in charge of fires, fire-drill should be held sufficiently often to familiarize the men with their duties in case of conflagration, and fire-buckets filled with water should be distributed about the buildings. As a matter of cleanliness, and still more as a matter of hygiene, and especially in the tropics, this water should be renewed once or twice weekly, to prevent its serving as a breeding-place for mosquitoes. Frequent inspections should verify the thoroughness and effectiveness of this measure.

The avoidance of mosquitoes, flies, roaches, lice, and other vermin has already been shown to be principally a matter of screening, cleanliness, and police; but if an old barracks is infested with bed-bugs or fleas, it is often a most difficult matter to get rid of them.

The most effective way to do so is to give the building and furniture a thorough overhauling, scalding out

Vermin

72 THE RECRUIT AND HIS ENVIRONMENT

cracks and crannies of beds, floors, and walls, with boiling water, washing them with *five per cent* carbolic solution, then closing all cracks with putty or strips of board, and painting over all. Infested clothing or bedding should be washed or steamed.

Frequent inspections should be made for the early detection of any recurrence or reintroduction of the pests, when their eradication from one room, or even from one bed, may suffice to end the trouble.

CHAPTER V

CAMPS

THE soldier's time is partly spent in camps, and occasionally the periods so spent may extend to many months. The conditions of camp-life are in many respects radically different from those in barracks; the old sanitary problems may have to be met in very different ways, while new ones arise, and sickness once started has opportunities to spread such as are not offered in the barracks. Our own unfortunate experiences in the camps of concentration in 1898 clearly show that the assemblage of large bodies of raw troops into great camps is in itself a dangerous matter, and that, if wise care is not exercised to prevent it, diseases in epidemic form may create greater havoc than the enemy in arms. It is the duty of the medical department to take the initiative, and to be responsible for much of that care; but its work can only be made effective and really preventive of disease when backed by the intelligent and sympathetic coöperation of the officers and men of the line. However great may be the powers and responsibilities of the medical department in sanitary matters, they can never be so great as to interfere with some urgent military necessities, or to free it from the need of the coöperation mentioned. It is therefore necessary that the company officer familiarize himself with many of the sanitary problems presented by camp-life, with a view to teaching his men to take proper care of themselves and so to assist him and the sanitary authorities in maintaining their efficiency.

74 THE RECRUIT AND HIS ENVIRONMENT

At times the choice of the camp-site will devolve upon the company commander, at other times a site is assigned to him by higher authority, and he has no choice in the matter. It is, however, usually within his power to improve a poor site, or to impair or destroy the value of a good one ; and whether he is more likely to do the one thing or the other depends largely upon his knowledge of military hygiene. The general principles regarding the selection, arrangement, and care of camps, as laid down in the Field Service Regulations, are to be observed, though they can at times be improved upon, and when such is the case it should be done. Let it be remembered that from a sanitary standpoint the important desiderata are: first, that the site should be clean and healthful; second, that it should be kept clean and free from infection during its occupancy; third, that it should be left so after the departure of the troops.

The consideration of the site of a camp must include an investigation of the healthfulness of the surrounding country, and in certain instances a very desirable site is wisely abandoned for one apparently less so, because the health conditions may be known to be better in the latter. For example, in the Bitter Root valley in western Montana there occurs during the spring months a fatal fever, which is conveyed by the bites of infected ticks. These are found principally on the west bank of the stream, the eastern side being almost free from the disease. It would be wise, in that territory and at such a season, to forego considerable advantages that might be found on the western side, and to take the trouble of crossing the stream, rather than to run the greater risk of infection with the "tick fever." A somewhat similar

disease occurs in certain river-bottoms in Japan, and in that case the camp could be safely placed on the infected *side* of the stream, *above all levels subject to flooding*, as the infected insects are met with only at such levels. In Africa sleeping sickness is transmitted by a fly that is found only in or about thick underbrush close to water, and relapsing fever by infected ticks that appear principally in regular rest-places or camps. It is possible that other diseases of equally strict localization may occur, and it is important to avoid them if practicable.

The qualities desirable in a camp-site vary somewhat with the season, breeze and shade, for instance, being desirable in summer and undesirable in winter. The site should have a natural slope, to insure the speedy disappearance of rain-water from tents and streets. The surface should be as regular and smooth as can be obtained, free from numerous rocks and hollows. Stumps and trees are at times objectionable, particularly if numerous and close set, but at other times they can be put to a surprising number of uses, and those that are most in the way will disappear in a short time after the camp is made. The inequalities of surface due to numerous large roots or lightly covered rocks are, however, harder to overcome.

The soil of the camp-site should be dry and firm, so as not to give rise to a great amount of dust in dry, or mud in wet, weather. Clayey soil, if the surface slopes gently and drains speedily, is good. Sandy or very soft loam is undesirable, and, in very windy seasons or neighborhoods, the dust it gives rise to may be a plague. A surface well set with grass is good, if not damp; but the grass will soon die in tents and be

tramped to pieces in the streets, and should be well cleaned up. The neighborhood of marshes or standing water should, if possible, be avoided, principally because they are apt to be infested with mosquitoes, but also because the ground may be wet. Ordinarily, high-ground water (shown by its nearness to the surface in wells or pits) makes for insalubrious conditions, much as do marshes.

Winds may be desirable or the reverse, according to circumstances. In regions where mosquitoes are abundant, a prevailing wind blowing them away from the camp is a great comfort, one in the opposite direction a great nuisance. In wintry weather a strong wind always produces discomfort, and in summer it may do so because of the dust and dirt it blows, and in such circumstances shelter from it should be sought.

As sites are unhealthful or infected, in most instances, either because of the presence of disease-carrying and infected insects, such as mosquitoes or ticks, or because of their previous infection from human occupancy, old camps, marshy grounds, and places apt to present the conditions mentioned, should be avoided. While other things than germs may and do cause disease, they can, to a relatively large degree, be safely ignored if infection be prevented; and the original choice of the camp-site may be of little importance as compared with the measures taken for such prevention. While the avoidance of malarious swamps and of such disease-centres as have been mentioned above is of the greatest importance in regard to certain diseases, especially those that are insect-borne, it is usually camp diarrhœa, dysentery, typhoid, and other infections disseminated largely by man himself that create the

greatest havoc in camps. The danger from these usually increases with the length of time the camps are occupied, and with the increased opportunity for their defilement. This being so, bivouacs, temporary and permanent camps, are apt to be defiled in an increasingly dangerous degree in the order named, and care in policing is of increased importance in the same order, though it cannot be safely neglected in any of them. Even a dirty bivouac-ground may constitute a menace to the inhabitants of the region, to other troops later following the same route, or to the command leaving it, on its return that way.

The danger of infectious diseases is lessened by attention to the following points:—

The ordinary shelter of the camp is furnished by tents, which may be of any of the varieties issued for the service, though the shelter-tent is only exceptionally used, in one camp, for more than a few days at a time. All tents are apt to be hot in summer and cold in winter, objections partly met by the proper use of tent-flies. All are liable to be crowded, and the air to contain many micro-organisms when they are kept closed. This can be met by thorough sunning and ventilation, with walls raised, for a part of each day when the weather permits, and by the frequent removal, cleaning, and airing of all blankets, clothing, and accoutrements to which the moulds and other organisms adhere, and the exclusion of articles in or on which they may grow, such as food, dirty straw, trampled grass, empty but unclean bottles, cans, and boxes. All tents are apt to be damp in rainy or wet weather, and to lessen this they should be well ditched, so that all water from the tent-walls and the

surrounding ground is carried away and the tent-floor kept dry. If the camp is to be occupied for any considerable length of time, board floors should be furnished, or the interior be filled with fine gravel and rammed, while company streets should be raised and made hard and smooth by pounding or rolling.¹ If floors are used, they should be raised occasionally and the space beneath thoroughly cleaned and aired. The general tent ventilation is also important in promoting dryness. Because of the dampness, men should always try to sleep off of the ground. In permanent camps they may have cots. Straw sacks are helpful, but they also should be raised if practicable. Cots or beds may be improvised from timber, or easily from bamboo, in a country where that abounds. Small branches of pine or other evergreen boughs serve to make a well-ventilated and comfortable bed where they are found. Boughs, cornstalks, loose straw, or dry leaves may be available in bivouacs. In other camps than bivouacs these materials should be stirred up and aired daily and renewed before they become mouldy or much broken up.

Owing to the crowding in camps and the frequently poor bathing facilities; the opportunities for the spread of body-vermin are unusually good, and all precautions as to cleanliness, clothes-washing, and sterilization, if necessary, should be exercised to prevent their introduction and dissemination. Dogs, cats, and other pets should be excluded from tents and usually from camps. Mosquitoes are to be avoided by means to be considered later, but if they are present, nets must

¹ Sprinkling the streets with crude petroleum and rolling them, as is now done in parkways and some other public roads, might prove of value in permanent or semi-permanent camps, in lessening both dust and mud.

be used. Flies, roaches, rats, and mice are best kept down by the maintenance of such good police that they find no breeding-places, and no access to food in or about the tents; hence the exclusion of food, boxes, cans, etc. In some places, as the neighborhood of San Francisco, the camp is very apt to be full of fleas, and, as these insects do not stay long on their hosts and breed in the sand, floors, or bedding, cleanliness of tents, floors, and surroundings is most important, as the young live on decomposing animal and vegetable matters found there. The adults, however, can live only on liquid food, practically only on blood.

Because of ignorance, laziness, or viciousness, men at times urinate in or near tents instead of seeking urinals. The offense may not be readily detected, but it is none the less filthy and dangerous, and should be severely punished.

Huts, cabins, and dugouts may be used instead of tents in long-established camps, particularly in cold climates or seasons. Because they are not so accessible to light, ventilation, or cleaning as are tents, even closer and more frequent inspections and greater care in cleaning are necessary, and particularly so in the prevention or elimination of vermin. At least once a week all clothing, bedding, accoutrements, and other accumulations of personal belongings should be moved outside and the places thoroughly inspected and cleaned. Such places as well as tents will require heating in cold weather, but as long as proper precautions are taken against fire and to provide for the escape of combustion products, this is desirable, as promoting dryness and ventilation, as well as comfort. Dryness of shelters and abundance of fresh air should be assured by all possible

means, and to every man. No one should be allowed to spend all of his time inside hovering over a stove, but all should work out-of-doors a part of each day. Wet clothes should be dried and dirty ones washed outside of the shelter, if it be at all practicable. Men with clothing or shoes wet from rain, snow, or slush should change them for dry at the first opportunity, and before lying down to rest or sleep. So far as military necessities permit, men, even those on guard, should be allowed to shelter themselves from inclemencies of the weather, and to overcome the results of over-exposure as soon as possible, and by other means than alcohol, which many seem to regard as the only remedy.

The water-supply to permanent or semi-permanent camps may be piped throughout them and delivered from stand-pipes conveniently placed. In **Water** other instances, it will have to be hauled or carried from wells, cisterns, streams, and other sources of supply, to the place of consumption. If it is known to be good and free from infection, it may be drunk as delivered; but when doubt exists as to its purity, and such will most often be the case, it must be boiled or filtered before use. Boiling destroys all disease-producing organisms that are found in water, and renders it safe. Prolonged boiling (the Field Service Regulations prescribe it for 20 minutes) makes assurance doubly sure, but if fuel or time is precious, making certain that the water comes to the boiling-point once may suffice, as that will destroy the organisms of typhoid, cholera, and dysentery. But it is at best a time-consuming process, and if the water is not well cooled it is not so refreshing, and men seek it from other sources. Other methods of purification are therefore resorted

to, and of these, precipitation and filtering by means of the Darnall filter promises to be simple, rapid, and effective. This filter, when obtainable, will be accompanied with printed directions for its operation, and these should be carefully followed. Whether the water be boiled or filtered, reliable men should be detailed in charge of the drinking supply, preferably under the direction and supervision of a medical officer, and should be well instructed in their duties. Forbes-Waterhouse sterilizers, Berkfeld filters, and other means of purification, are at times resorted to. Many of these are valuable, but are not used because of the difficulty of keeping them in order or the slowness of their operation. They should not be resorted to unless their working and results are understood, and the line officer is advised to put the responsibility for the purity of the water-supply on the medical officer, and then to lend him every assistance in assuring it. However purified, the drinking water should be kept pure and protected from all sorts of contamination until used. This is provided for in the directions accompanying the Darnall filter, but special arrangement has to be made in other cases. A simple and effective way of doing it is to have one drinking-place for each company, and to have at that place two or three barrels or large cans fitted with covers and with faucets near the bottoms. Water should be boiled, or otherwise purified, and run into one of these, covered, and allowed to cool. When cool it is marked as ready for use, and the water is drawn off as needed through the faucet. Meanwhile the next canful is cooling. When the first can is emptied, it is rinsed out with boiling water and refilled, to cool while the second is in use, and so on. The barrels or cans

82 THE RECRUIT AND HIS ENVIRONMENT

should be elevated on a platform so that the faucet may be convenient, and a soakage-pit should be dug beneath the drip and filled with small stones or gravel to prevent the formation of a mud-puddle. Waste of the water should be discouraged, but if it amounts to more than can be cared for by the pit, which should be about one foot deep by two to three feet square, outlets a foot wide and six inches deep, and of such length as necessary, may be made and filled also with stones or gravel, to lead away the excess and permit of its absorption.

Outside contamination is prevented by the use of the tight-fitting cover. If faucets are not to be had, siphons may be improvised, if rubber-tubing is available. Otherwise, the water will have to be dipped. If so, a clean dipper should be provided for the purpose, and the water emptied from it into the drinking vessels. No cup should be dipped in the water after it has been contaminated by hands or mouth. The dipper should be scalded from time to time, and should be kept clean in every way. If reliably clean ice is obtainable, it may be added to the water. If its quality is doubtful, it may be packed around vessels of water, but should not be put into them. The general care of the water-supply as prescribed in Field Service Regulations should be observed.

There are no tests that can be quickly applied that will enable a medical officer to pronounce a given water-supply safe. He may form an opinion to that effect from a consideration of the source and surroundings of the supply, but any idea that he can, by a simple and quick chemical test, or a microscopic examination of it, gain positive knowledge that it is pure

and safe is fallacious, and should not be entertained. If the supply probably receives seepage from privies, stables, or homes, or washings from the same, or drainage from near-by cities or towns, it must be regarded as unsafe; and in other cases, where sources of contamination are not so evident, it is wisely precautionary to take the same view. In fixed camps there may be opportunities for the full and careful examination of water-supplies; but even when such is the case, and the water is found satisfactory, the condition may change in a day because of some accident or the carelessness of one or a few men. Boiled water should be used for many kitchen purposes, such as washing food-receptacles and such foods as are eaten raw. Boiled or actually boiling water should be furnished the men for washing their mess-kits. A good arrangement is to have for each company a box, a row of large kettles of hot water, at least one of them actually boiling, and a small mop or scrub-brush. As a man finishes a meal, he takes his mess-pan and implements to the place where these are, scrapes any food-remnants into the box, rinses the articles clean in the first kettle, passes to the second and there scrubs them with the brush, and then scalds them in the boiling water of the third kettle. Proper provision should be made, by means of a trough-container or otherwise, to prevent soiling of the ground with bits of food or greasy water.

The water-supply should be sufficient to provide baths for the men. If the camp be situated on a stream or other body of water, it may afford proper facilities in warm weather. Where such is **Bathing** not the case, and in cold weather, special provision will need to be made. Such provision will include a proper

supply of water, sheltered and warm bathing-places, and a proper disposal of the dirty water.

The supply of water may be piped to the bathing-place, or it may have to be carried there. The former condition need not, ordinarily, concern the company officer, except that, if the water as delivered is very cold, most of his men will not bathe in it in cold weather. In case it has to be carried, each man should carry for his own use, and economy will then be exercised. The most effective use is then made of the water if the bather first scrubs himself from a basin, using soap and a rough cloth, and later rinses off the soap with a shower, which may be improvised by means of a large tin can and a piece of rope. Warm water should always be available, and may be kept so if a large can of water be kept over a fire near the bathing-place, and any man desiring to bathe be allowed to take from it, replacing an equal amount of cold water. A sheltered and warm bathing-place may consist of a room in a convenient building, of a special house, shed, or tent, heated, if necessary, by a stove, an extemporized brick oven, or other means. The floor should be hard and dry, or should be covered with wooden gratings to lift one above the waste water. If for lack of time or other reason a suitable bathing-place has not been prepared, and if it be cold weather, hot water may be furnished and men allowed to take baths in their tents. In such cases they should be careful not to spill water.

If bathing is possible only at the cost of considerable labor or inconvenience, a proportion of the men will neglect it, and it is therefore desirable that a proper place be provided and the men required to bathe at

least once a week ; while the feet, face, and hands should be bathed oftener, and facilities should be provided for that. These may be just outside or inside the tents. The disposal of waste water is at times as great a problem as the procuring of the fresh supply, and it will have to be solved in different ways under different circumstances. In large or permanent camps underground drains may lead it away ; at other times trenches may conduct it to a near-by stream or to dry wells or pits, and occasionally it may be necessary to collect it in barrels and haul it away. Always, care should be exercised that it does not flow or soak to the source of supply and contaminate that, and that it does not form breeding-places for mosquitoes. Where other provision is not made, fairly satisfactory results may usually be obtained as follows: A suitable space inside of a tent, say five or six feet square, is dug to a depth of four inches, and filled with gravel or fine stone, which is rammed. Over this a wooden grating is placed, and on the grating the men bathe. From the lower side of the space, trenches four inches deep and a foot wide are dug and filled with gravel or stones, and they conduct the water downhill to an open trench that leads it away, or to a soakage-pit about a yard square and deep, or larger if necessary, that is filled with large rocks, whence it soaks into the ground. Such pits, if in porous soil or gravel, will dispose of a great amount of water in a day. If in clay or close-grained soil, their working is not so satisfactory, and it may do better to run the water out in numerous, shallow surface-trenches from which it can evaporate. Similar precautions may be taken to provide for the waste water from drinking-troughs for animals.

Facilities for washing clothing should be provided in each company, though they need not be more elaborate than a supply of good, and preferably warm, water, a board, and a scrubbing-brush and soap. The waste water may be disposed of as indicated for bath-water. If a boiler can be provided for clothes, it is desirable to have it. In rainy weather a heated tent should be furnished in which to dry them, but at most times they can dry in the open.

As in barracks, so in camp, the kitchen is one of the most important parts of the command, and on its proper administration depends much of the cheerfulness, health, and efficiency of the men. **Kitchens** The food should be abundant and good, and its preparation such that the men will be nourished and satisfied by it, and not under the temptation to gorge themselves with pies, milk, and soft drinks of doubtful character from outside sources.

The mess in fixed camps may be made quite as good as in barracks, and, except in rare emergencies, the articles of the ration are both good and abundant. As the problem of cooking is presented somewhat differently in the field and in barracks, the company commander, the mess-sergeant, and the cooks should all study the best methods for field service. Some excellent garrison cooks are not nearly so good in the field, while others excel there. Economy, good management, and cleanliness are as important in the field as elsewhere, while, owing to the outdoor life and active exercise, the appetite is usually increased and more food required. Owing to the liability of chilling, exposure to wet, and the greater opportunities for intestinal infection, irritating, indigestible, and slightly nourishing foods,

such as green corn and boiled cabbage, should be used sparingly, while green or overripe fruits should be forbidden. In the presence of epidemics of intestinal diseases, it makes for safety if only cooked food is eaten. Underdone cereals or vegetables are apt to cause indigestion, and as they are more difficult to cook thoroughly in the field because of the trouble with fires, wind, etc., particular attention should be paid to them. Rice, beans, hominy, oatmeal, and potatoes are probably most apt to be underdone. The methods of cooking in camp are various, the facilities ranging from the most simple and crude to the elaborate. Soup-carts, baking-ovens, and so-called fireless cookers give promise of solving some of the problems; but the knowledge, industry, and management of the company commander, mess-sergeant, and cooks must be relied on to solve most of them. They must utilize to the utmost the articles of the ration, and such other good food-supplies as they can obtain, and the equipment furnished for cooking and such additions to it as they have or can prepare; and must put to use or minimize the harm of natural factors, such as wind, rain, snow, or poor fuel. It is the ability to do this last that enables some men who are not very good cooks in garrison to excel in the field. While a man knowing more of cookery may let a meal be spoiled or uncooked because of poor fuel, rain, and adverse winds, another man may have an excellent fire and a well-cooked meal in spite of all three. Camp craft results from experience and ingenuity, and the one should be furnished by manoeuvre-camps and practice-marches, and the other encouraged by observant company commanders when it manifests itself. The man possessing these qualities will get the most from

88 THE RECRUIT AND HIS ENVIRONMENT

his utensils and his fire, whether he has a fully equipped camp-stove, a spider and pots, or merely a trench of his own digging.

As far as possible, men coming in from a march or a guard should be supplied with hot food at once, and when camp is changed an effort should be made to supply a hot meal as soon as the regular meal-time comes. This means the use of kitchen carts or fireless cookers, or the early establishment of the kitchen and the use of food that can be quickly cooked.

As in garrison so in camp, cleanliness of cooks, kitchen, and food are secondary only to the actual supply of food, and the lack of it will do more harm than partial starvation. Flies have greater opportunities to breed in camps than in garrison, dust is more abundant and blows about more freely, fecal contamination of hands, shoes, and clothing occurs more readily, and the facilities for storing food so as to protect it from all these are poorer. Consequently, the amount of care to be exercised is greater, and the necessity for minute and thorough cleanliness more urgent. The hands, persons, and clothing of the cooks and assistants should receive the greatest care. Tables, benches, blocks, cooking utensils, and everything about the kitchen should be cleaned as soon as used, and scalded, if it be possible, with boiling water. Boxes and bags containing food should be frequently inspected inside and out, and their positions shifted, in order that insects may not gather in or about them. They should always be kept clean. Kitchen floors and surroundings should be raked, swept, or scrubbed after each meal, so that all particles of food and everything that might attract flies will be removed. Slop-buckets and garbage-cans should be washed after

each emptying, and not allowed to become crusted with a greasy coat of filth. At least once a day a rigid inspection of the kitchen should be made and all points of kitchen police investigated. Reliable men should be put on this duty and rewarded for its proper performance. Mosquito netting or wire-gauze covers should be provided for the protection of food while it is awaiting preparation or serving. Sheeting or boxes should protect it from dust.

It is questionable whether screening the entire kitchen is always advisable. Cleanliness is necessary in keeping down flies, and if a kitchen is screened all around it is darker and has more angles and corners in which dirt and scraps may collect to attract them. As the door is of necessity frequently opened, they enter and are then kept in by the screening. Screening is an aid in preserving the food uncontaminated, but it does not justify even the partial neglect of the more important matter of scrupulous cleanliness.

The kitchen fire should constitute a small crematory for kitchen-waste, and the cook should endeavor to be as independent as he can of civilian scavengers and fatigue parties. Nearly all scraps and solid waste that are to be thrown away can be burned by him, and at times he can take care of most or all of his dirty water by devices to be described later. Every tin can that is emptied should be thrown into the fire and all organic matter in or on it destroyed. Later it may be hauled away or buried.

Major H. A. Arnold has described the following arrangement as working satisfactorily in the Pennsylvania militia. A pit is dug about 60 inches long, 30 inches wide, 48 inches deep at one end, and 36 inches deep at

the other. It is filled loosely with stones to a height a few inches above the general ground-level, and is banked on all sides to protect it from surface-water. The kitchen fire is built on these stones, and is inclosed on the sides, but the ends of the trough thus made are left open. All watery material, such as waste coffee and dish-water, is poured on the stones at the shallow end of the pit. The heat of the stones evaporates it. All solid waste, including tin cans, is burned in the fire. The ashes and *débris* are removed as necessary. It is thought that this might be somewhat improved by having the liquid waste first strained by passing it through a layer of straw or excelsior six inches deep, which would catch much of the fine solids and grease in it. The straw should then be burned and fresh supplied once or twice daily. It might be placed in the bottom of a large tin or other box, or in a shallow pit that drained into the larger one. At other times, when the particular conditions would warrant it, the water could be led away to a stream by such a filled trench as was described for the bath-water, provided it were first strained through straw. Otherwise the trench would soon become foul with a deposit of soap, grease, and food-particles. When garbage and waste water are hauled away for disposal elsewhere, the matter of emptying them from cans and loading them into wagons should be carefully done under the supervision of a responsible man, with a view to the prevention of spilling and the immediate cleaning up of any that may occur. Receptacles should be washed inside and out immediately after emptying, and should at all other times be kept tightly covered. Lime-washing the cans and the stands on which they rest renders the detection of dirt easier and is therefore recommended.

If abundance of good food is provided in the messes, the men have no proper excuse for indulging in improper food obtained from peddlers, and such persons should be excluded from the camp and its environs.

Papers, sweepings, and other ordinary camp litter, as well as horse-manure and stable-dirt, should be burned, and such disposal of it is easy and convenient. If not so disposed of, it is apt to become scattered, create a nuisance, and, particularly, to do harm as affording breeding-places for flies. A crematory may readily be improvised, and if a good draught is furnished, it may be kept burning continually with waste as fuel. Three forms of pit or crematory are as follows, and any one of them may be easily and quickly constructed:—

Disposal of Waste

1. A cylinder of about the height and capacity of a barrel, built of blocks of sod about a foot square. It may be steadied by having posts planted on two or three sides of it, and being wrapped with wire. An air-inlet about ten inches square should be left at the bottom, and if iron piping or rods are available a grate may be made of them.

2. A circular pit with a cone of stones, around which the fire is built, piled up in the middle.

3. A horseshoe-shaped mound of earth, or a niche dug in a bank, or a trench open at one end.

Other forms may be improvised to suit conditions, and bits of stove-pipe or other means of improving the draft or protecting the fire from rain can be utilized. These pits must be cleaned as necessary, and the ashes and unconsumed matters, such as tin cans, placed where they cannot constitute a nuisance.

The proper disposal of the products of human waste,

feces and urine, so that they may not create a nuisance or constitute a source of danger, is one of the most important problems of camp sanitation. The nearest approach to its solution that has ever been made is their incineration, and the difficulties in the way of that, except in camps of some permanency and at considerable cost, are so great as to be in most instances insurmountable. In our manœuvre camps and permanent camps the *McCall incinerator* has proven a great boon, but whether it will prove as valuable and as workable when used by troops that move frequently is doubtful. Even as used now, though, it is of great value and justifies the rather large expenditure it necessitates. Care in its management is necessary to keep down unpleasant odors, and repairs must be made promptly, as the need for them arises, or the whole apparatus may get out of order and cease to operate satisfactorily. A competent and intelligent man should have supervision of each one, should be familiar with its construction and the manner of operating it, and should see that it is operated properly and in accordance with directions. In some of our manœuvre camps it has been possible to install inexpensive but satisfactory systems of *water-carriage latrines* that empty into the sea or other safe place. Such a system consists of a long iron or zinc trough, much like the Reed trough, sloping from its upper end, where water enters it, to its lower, where is an outlet which is closed by a cylinder that screws into threads in it and fits tightly. The water-supply is from a faucet, and may also be from a water-flushed urine trough that empties into the upper end of the main or large trough, above which is a long board with seat openings, which

may or may not have hinged covers. The whole plant is housed in a cheap but clean shed. The trough is filled with water, into which the feces and paper drop, and is flushed by gravity when the care-taker removes the cylinder that is screwed into the outlet, the entire contents then flowing into the drain of glazed earthen pipes that lead it away. Such a system is open to the objection that it applies only to fixed camps.

The *Reed trough, pump, and wagon* as described in G. O. 170, A. G. O., 1899, offer another fair method of disposal of excreta, inferior to those discussed above where they are applicable, and so cumbrous as not to be available except in camps of some permanency.

The methods of disposal actually used in short camps, or by troops that change positions at all frequently, are the use of deep trench latrines, of shallow trenches, and of no trenches, but rather a disregard of the subject. All of these are filthy, dangerous, and objectionable. They are apt to constitute disgusting nuisances to the eyes and nose, dangerous sources of fecal contamination of shoes and clothing, sources of camp-pollution by seepage and leakage, and, worst of all, vile and dangerous breeding-places for flies that carry contamination from them to the food or persons of the men. These objectionable features may be lessened by proper care, and it is most reprehensible to neglect it.

The *deep trench* is probably most generally used by our troops in camp, and is prescribed in Field Service Regulations. A trench two and a half feet wide, six feet deep, and of varying length, is dug, a seat is provided by means of a small log or piece of timber supported on forked or crossed pieces, and the whole more or less protected from sight by a brush screen. The care

exercised varies from the occasional perfunctory covering of feces with earth to the abundant use of lime and the daily burning out of the whole pit with crude petroleum. Whatever measures are used, they are usually insufficient to keep down odor and flies in hot weather. The best results are probably to be attained by having a man stationed at the pit, whose business it is to keep all fecal matter and soiled paper covered with loose, dry earth at all times. The ground about the pits should be kept thoroughly policed and protected from contamination with feces and urine. If crude oil is available, it may be advantageously sprinkled into the pit daily, as an aid in keeping away flies or destroying their larvæ, and for the latter purpose its use should be rather lavish. The whole pit should be protected from surface-washings by ditching and embankments, and from rainfall by a tent-fly. No more disgusting sight can be imagined than that afforded by such a pit partly full of feces, alive with maggots, and flooded by tropical rains.

When flies begin to breed in a pit, and the larvæ cannot be destroyed by oil or otherwise, it should be filled and another dug. Burning the pits out daily with straw and crude petroleum may aid in keeping down flies, but as often practiced it will not entirely do away with them or their larvæ. Immediate and rather deep covering of dejecta, and thorough policing of the pits and their surroundings, are the most efficient means of caring for them. They should always be placed as far from the kitchens as is practicable, and where they will do a minimum of damage to water-supplies. Toilet-paper should be provided and kept in covered boxes, so that it may not be blown about camp. The pits should be

inspected at frequent intervals by an officer, who will see that the man in charge is doing his work properly, that flies are being kept down as thoroughly as possible, and that men who willfully or carelessly pollute the surroundings with urine or feces are punished. It is always essential that a separate urine pit be provided, as a proper state is more readily maintained in the feces pit if it be dry.

Major William Jepson, M. D., I. N. G., devised the arrangement described below, and it was put in successful use in the encampment of the 56th Iowa in July, 1908. The quotation is from a report by Major Daniel H. Boughton, U. S. A.

"Near each kitchen fire a hole eight or nine inches in diameter and six or seven feet deep was dug with a post-hole auger. Into the mouth of this hole was fitted a bucket with a wire bottom to catch solid matter as slops were poured in, the top being flush with the surface of the ground and provided with a lid. The liquid matter ran through into these post-holes, and the solid matter was thrown into a garbage-can and then removed in the usual manner by police parties. Instead of a bucket specially prepared to catch solid matter, a piece of wire netting or gunny sack could be used.

"One of these holes sufficed for each company kitchen during the seven days' encampment. The ground where this method was used consisted of about three feet of black loam with a subsoil of clay, the most difficult ground in which to dispose of liquid waste; nevertheless the method was satisfactory. Sand or gravelly subsoil is better. If pits are dug with the spade they should be kept covered with pieces of thin muslin or other material.

“Latrines were prepared by digging with the auger a series of holes at the proper distance apart, and also by digging trenches in the usual manner. Over these holes or trenches boards with the necessary openings were placed at the proper height, and thin muslin was tacked to the edges, and stretched to the ground, where it was pinned with small wooden pegs; covers for the openings were also provided, making the pits absolutely fly-proof. These covers in some cases were made of tin, and in others of thin muslin to which a piece of wood was tacked to hold it over the opening when not in use. Urinals were made by taking pieces of the same cloth coated with gas tar and suspending them in the form of troughs in rear of the latrine seats. Each trough communicated with the trench by a pipe made of the same material as the trough.

“The latrines were screened from view but left open at the top, and though the deposits were not covered nor disinfectants used, the odor was scarcely distinguishable a few feet away. It was constantly escaping into the air. Latrines of this nature can be readily constructed by troops and at very little cost. The secret is in keeping out the flies. As the deposits were not covered with earth or ashes, a single trench accommodating six seats was sufficient for a battalion of over 200 men during the entire week.”

The *shallow trench* system is prescribed by Field Service Regulations for camps of one night only, but it has found more general use in the British Service, and the following advantages are claimed: 1st, That the walls do not have time to become foul, and the odor arising is very much less than from deep trenches. 2d, That flies are much less numerous about them. 3d, That

the feces become speedily transformed into simpler matters, so that no signs of fecal matter as such remain after three or four weeks. 4th, That disease-producing organisms are more quickly destroyed. As used during the manœuvres in Hampshire in 1907, the system was as follows, and was said to give excellent results. The trenches were dug one foot deep, one foot wide, and three feet long, and were arranged in lines two and a half feet apart. If ground was scarce they were made two feet deep. Enough trenches were dug to accommodate *five per cent* of the command. The turf was carefully removed and laid aside, the dirt taken out was pulverized. A man was at all times detailed in charge of the pits, to see that each soldier thoroughly covered his own deposits with pulverized earth, for which purpose cans or shovels were provided. The trenches were usually found to last two days, when they were carefully and solidly filled and the turf replaced, new trenches being dug in the interspaces. The same precautions as to dryness, blowing paper, complete covering of excreta, separate urinals, and general police have to be observed as when deep pits are used.

The *dry earth* and *pail systems* do not need discussion, as they are not satisfactory for use in camps, and where conditions are such as to permit of their installation they will also permit of the use of McCall incinerators or Reed troughs, both superior to them.

As stated above, urinals should be provided separately for the use of men not desiring to defecate at the same time. The McCall incinerator and the Reed
trough system both make such provision, **Urinals**
but when pits or trenches are used, it is frequently neglected. The provision should be for both night and

day. *Night urinals* consist of large cans or half-barrels, to be placed in the company streets at night, and to be carried away and emptied and washed in the morning. If one emptying does not suffice to prevent overflow or spilling, they should be emptied also at ten o'clock at night. They should be marked at night with lanterns, so that they may be easily located. During the day, when not in use, they should be kept clean and dry and exposed to the sun. The *day urinal* may be similar, or it may be a trench, pit, or box. A circular pit may be dug, two feet in diameter and three feet deep, and filled with stones, clinkers, or sawdust, and the men allowed to urinate into it; or a barrel or box with both ends knocked out may be half-buried and similarly filled. If pine-needles or sawdust are available, they should be used for the filling, as they are credited with great virtue in preventing the development of bad odor. Shallow trenches, leading to pits filled with coarse rocks, may be used, and, as they become foul, filled, and new ones dug, until the pit itself is foul or the entire surface about it has been utilized for trenches. The Jepson urinal has been described.

While urine pits or trenches do not afford the fly-breeding facilities of fecal accumulations, they do present opportunities for the spread of typhoid infection, and care should be exercised to prevent dust or mud from them being carried to camp by the wind or on men's shoes, and particularly to prevent the pollution of water-supplies from them. Urine is more manageable than feces because of its fluidity, and, if necessary, it can be chemically disinfected in the barrels or cans, by the use of bichloride, carbolic, or formalin.

There is a tendency to look upon the matter of camp

**Camp
Police**

policing as an affair of military discipline rather than of great sanitary importance, and for that reason it is strongly recommended that whenever it is possible a medical officer should be made police officer. He understands better than the line officer the sanitary dangers of the camp, while his interest in having it present a neat and military appearance is quite as great. He is more interested than any one else in having a small sick report, and he alone, in most instances, finds interest and profit in the study and prevention of vermin and pests, in the disposal of excreta, and in studying the tastes and history of the fly and the mosquito. He should have at his disposal a sanitary squad under a sanitary sergeant, such a squad consisting of at least one man from each company and one noncommissioned officer of the hospital corps. In addition to that, the provost sergeant and fatigue parties should report to him, rather than to the quartermaster, for instructions regarding all work having a sanitary bearing. He should be responsible for sanitary conditions, but he should have, with responsibility, a proper authority and proper freedom of action. The camp police will, at times, necessarily extend beyond the camp proper. It may embrace the improvement or protection of the water-supply, the draining of a swamp, the petrolizing of pools, the clearing of brush quite outside of the camp limits. Under certain conditions it may include a quarantine system, a war on rats, a house-to-house inspection of civilians, the suppression or regulation of traffic, as that in milk, green vegetables, or other foods. In all these matters the medical officer should have as free a hand as possible, and should be able to carry out his own recommendations under no

other control than that of the commanding officer. It is confidently believed that such an arrangement as is here advocated would make for greater efficiency and quicker and better accomplishment of sanitary measures, without in any way causing discord or impairing the value or dignity of the services of other persons.

CHAPTER VI

THE HYGIENE OF MOVING TROOPS

THE march, the battlefield, changes by ship and rail, all present their special problems of sanitation, neglect of which may result in disaster to the careless command or to its friends and allies. Many men and some officers are apt to ignore the latter phase of the question and to think that when they have avoided harm to themselves and their companies they have performed their full duty in sanitary matters. Nowhere is a man his brother's keeper to a greater degree than in an army, and it is hoped that the fact may become more fully appreciated.

Article V of the Field Service Regulations deals well with the subject of marches; but the rules there given should be applied with discretion. It must be remembered that the various possible **Marches** states of weather, roads, and personnel may result in an almost infinite variety of circumstances, all of which could not possibly be considered in any set of printed rules. Much must depend on the judgment of the commanding officer, in sanitary as in tactical matters, and his knowledge or lack of knowledge of sanitation may determine the success or failure of a movement. If good judgment is used in adapting the regulations to the circumstances, the rules laid down therein will constitute a reliable general guide. The length and speed of the march, for instance, must be materially influenced by climate, weather, roads, water,

transportation, weight carried, amount and character of sickness in the command, the rations, character and state of the clothing, the spirits of the command, the objective, temperance in the use of intoxicants, and other factors.

The length and frequency of rests must be influenced by much the same factors, and the commander who sets a pace and determines the frequency and duration of rests without regard to these influences is not doing justice to his men. The time of day during which the march is to be made will also influence the length, speed, and rests, and the choice of it must in turn be influenced by diverse considerations. Thus the temperature may make daytime marching very trying, yet the state of the roads, the possibility of malarial or other infections, the presence of sick, and other considerations, may make it appear the lesser of evils. It is always well, therefore, to consider these various influences and to bear in mind that the command brought into camp in good spirits and good condition after what may be a relatively short march, is in better condition for large accomplishments on the morrow, than another command that has marched five miles farther, but has reached camp with the men footsore, weary, discouraged, and with their vital resistance so lowered as to be ready to yield to the inroads of any chance infection. The influence of *climate* and *weather* is so quickly manifested on the men, and in such obvious and well-known ways, as not to need much discussion, and a moderate amount of common sense and consideration for his men should cause an officer to make allowances for them. Heat, wind, snow, rain, fog, slush, glare, mugginess, all tire, discourage, and tend to

sicken the men, shorten the march, and increase the rests.

Closely related to these is the *state of the roads*. Mud, excessive dust, loose sand or gravel, very rough and uneven roads, all increase delay, vexation, and fatigue, and, even when weather conditions are good, may cut the march to a fraction of what is desired. Not to be separated from the question of roads is that of transportation. If the former are good, abundant wagon *transportation* may be used, if available, lessening the loads of the men, supplying them with abundant food, tentage, and other necessities and comforts. If they are bad, transportation facilities may be much limited in consequence, or, if furnished, may prove a source of vexation and weariness, the men having to spend hours of toil extricating mired animals or wagons, in unloading, loading, and carrying freight. Such trials of body and spirit react on the health, and when, in addition, because of failure of the transport wagons, men have to lie down supperless and without proper shelter, the question of transportation has assumed a sanitary importance of magnitude.

Intimately related to the question of transportation is that of the *weight carried* by the men, and the attempt is wisely made to reduce this to a minimum; yet under some conditions, and generally they are those in which men should have to carry the least, as in bad weather, over bad roads, on forced marches, after very trying campaigns that sicken man and beast, transportation may fail and the men have to carry extra ammunition, food, and clothing. If this is a necessity the best should be made of it and the load carried, but the march should, as far as possible, be regulated

to suit the altered conditions. We are not able to breed or buy men as we do horses, with a view to their strength, speed, endurance, or spirit, though we do attempt to select them with a view to these qualities. Nevertheless, each organization will have its strong and its weak men, its fast and its slow, its cheerful and its despondent, and it is the less desirable of these that must be most considered under trying conditions. The weak man must not be overloaded, the lame man overmarched, because the more able men can march farther or carry more. Otherwise they will become sick, and the problem is worse complicated than before. Disregard of this simple fact has been known to result in most serious consequences. Whatever the weight carried, it should be so disposed on the person as to carry as comfortably as possible, and *not constrict or compress the chest*. As stated before, it is on the free play and efficient working of the heart and lungs that strength and endurance depend. If the chest is compressed and its mobility lessened by tight clothing, crossed straps, or other cause, that free and efficient working are impossible, and the man becomes exhausted under a load that might not greatly inconvenience him if more wisely disposed. Recent experiments have demonstrated that inhalations of oxygen enable athletes to run farther and faster, and with less inconvenience, than is the case without such inhalations. When the action of either the lungs or heart is interfered with, the oxygenation of the blood is hindered and early tiring, weakness, and exhaustion follow. These conditions constitute not only disturbing and delaying factors of a temporary nature, but also weaken the resistance to disease and prepare the way for infections.

Sickness in the command that is moving always occasions delay, often very much extra work, and, quite as often, danger of epidemics. All sick should therefore be left behind, if possible, when the march begins; all hygienic measures should be employed to prevent other men becoming sick; all who do become so should report to the medical officer as soon as practicable so as to get the benefit of early observation, treatment, and care, and all who do not favorably and quickly respond to such early care should, if possible, be provided for on the way, in houses, camps, or hospitals, or sent back, and not allowed to hamper and constitute a danger to the entire command. The Field Service Regulations prescribe that the sick shall be eliminated before the start is made, but in practice certain classes of sick are not eliminated and later give much trouble. Among these are venereal cases. Many such do not appear on sick report unless frequent inspections are made and they are ordered to report. These men, though doing full garrison duty and feeling well, are likely to develop buboes on the march and be unfitted for either walking or horseback riding.

Men recently discharged from treatment in hospital for malaria, dysentery, and other debilitating diseases are apt to become exhausted or to have recurrences of their sickness when subjected to hard marches.

Weak hearts and weak feet are liable to break down under similar conditions, and men known to suffer from either should be excluded. Not only should sick men be excluded from the march, but well men should be taught and, if necessary, compelled, to conduct themselves in such a way as to maintain their efficiency and strength. Alcoholism, particularly, should be discour-

aged, and all straggling to get alcohol and all drinking in ranks should be punished; for not only does the indulgence lead to inefficiency from drunkenness, but it predisposes to exhaustion, infections, injuries, to heat-stroke in hot weather and freezing in cold. While not useful nor to be encouraged in camp, drinking, if not carried to excess and not prolonged into the hours belonging to sleep, is then less directly harmful than on the march.

Cheerfulness and *bright hopes* are powerful stimulants to accomplishment, and it should be the aim of the company commander to keep his men in such a state of mind. Difficulties may often be laughed away, while if given too much thought or gloomily dwelt upon they readily become insurmountable. One of the great items in Napoleon's success was his faculty of having his men accomplish the impossible.

Hardships may be magnified in retrospect; on the march they should be minimized and belittled, if unavoidable.

Another occasional cause of exhaustion is said to be the reckless and excessive indulgence in *water*. The

Water Field Service Regulations state that under "ordinary conditions a canteen of water should last a man a day." Ordinary conditions in the field are, however, hard to define, and the amount of water needed under varying conditions and by different men varies greatly. There is no doubt that many men do use water injudiciously, and that, by training or habit, they may teach themselves such moderation and restraint as to be enabled to make the contents of one canteen last them through any ordinary day's march. As the canteen holds two and one-half pints

and the march rarely covers more than eight or ten hours, it may be seen that this should not involve any hardship for a well and fit man. It is, however, a well-recognized fact that men who begin a march with the consumption of large amounts of water, and who early exhaust their canteens, are very apt to become exhausted or to fall out before the march is completed. The consumption of the water is not always the sole cause of such exhaustion, but both may be the effect of a common cause, such as sickness, diarrhœa, alcoholic indulgence, excessive smoking, or injudicious eating the night before. By the avoidance of such indiscretions the moderate use of water is made easy; but if a man has spent a part of the night in alcoholic indulgence, smoking, and eating cheese and salt herring, his system is so loaded with poisonous waste-products that he needs a large amount of water to dilute and remove them, and depriving him of it will not improve his condition in any way. Aside from such indiscretions, more water is required by men whose food is very salty or very dry; by those marching in a hot sun or losing much water in the form of perspiration, or those marching in dust, as at the rear of a column. Thus we see that while "excessive" use of water is harmful, its liberal use is not. On the other hand, insufficiency of water is harmful, even when not causing suffering through thirst, in hot weather as predisposing to sunstroke, and as causing such concentration of the urine as to cause very severe bladder irritation, with burning and pain that occasion great suffering and alarm. It is nevertheless highly desirable that men should so conduct and so train themselves that they can be able to get through the marching hours without using more than one canteen-

ful of water; and if for any reason they cannot do so, it is usually better that they should go thirsty than that they should drink water from unknown sources or of doubtful character, as a few hours of discomfort from thirst, or an attack of pain in the bladder, are to be preferred to an attack of typhoid, cholera, dysentery, or parasitic infestation. Sunstroke, however, may be as fatal as any of these, and if symptoms of it appear, water must be used, under the direction of the surgeon, if one be present.

The water problem should be handled as follows:—

Before the march is begun, an abundant quantity of good water, boiled or filtered if necessary, should be supplied. If the water is boiled, that should have been done the night before, in order to have it cool in the morning. If other vessels are not available, it may be put in the canteens the night before and allowed to cool in them. Each man should drink what he wants before starting, using his cup for the purpose, and have his canteen filled from the good supply. Except rarely, he cannot know that the water he sees later in the day will be suitable for drinking, and he should therefore drink then only from his canteen, and, in order that it may suffice, exercise great care and moderation in his use of that supply. As soon as camp is reached in the evening, the preparation of a fresh supply should be begun, and, if boiling is the form of purification used, the first drink furnished may be tea or coffee. Reckless or careless indulgence in water from supplies of unknown character may have the most disastrous results; but the subject will be dealt with at greater length in a subsequent chapter. The Field Service Regulations say that “a medical officer near the head of the column may be called

upon to decide whether water is fit to drink." Again, let it be remembered that he can only base his opinion on the source and surroundings of the supply as he can see them or learn of them from others. The appearance, odor, taste, and such other tests as he is able to apply under the circumstances can in no wise assure him that the "water is fit to drink."

In the paragraph following that from which quotations have been made, it is stated that "good opportunities for watering (horses) on the road should not be neglected, as this is of great benefit in hot weather and on severe marches," the regulations thus recognizing in the horse a need for abundant water that is denied for the equally hard-worked animal, man. It is true, though, that the horse is not so apt to suffer from dissipation and to need the relatively large amount of water required by the dissipated man. It is sometimes urged that cold weak *tea* or *coffee* be carried in the canteen and drunk on the march, in order to assure the use of boiled water. Aside from the possibility of their being diluted with unboiled water, and of tea being made with cold and unboiled water, these drinks are not so refreshing and thirst-satisfying between meals, to most Americans, as plain water, and it is not thought that they are as good for use on the march as the latter, provided that it is properly purified or sterilized.

The food-supply on the march, as elsewhere, is very important. Because of the greater amount of work performed, and the increased tissue-changes due to the outdoor life, the amount of food **Food** required on the march exceeds that required in garrison; while because of the few and imperfect kitchen facilities, the new and strange environment, the absence of water-

closets, and for other reasons, food infection is more apt to occur, and every effort should be made to obtain cleanliness and to serve all food sterile, rendered so by heat. Because of the exposure to which the men are subjected, diarrhœal troubles are readily induced, and irritating foods should be avoided. Except in rare instances it should be possible to furnish two hot and substantial meals a day to the marching column, and that should be done. A hot breakfast should be served before the march is begun, and it should be a substantial meal, such as one of meat, potatoes, bread, and coffee. Eggs and similar articles, if obtainable, may be used. The noon meal, also, should be prepared and issued to the men before starting, and should embrace much nutriment in relatively small bulk. Fried bacon, cheese, sliced meat, and bread meet the requirements well, while a pickle, a bit of fruit, or jam adds to the enjoyment of it. In the evening, after the hard work of the march is ended, the men should have another hot meal, and as they will have leisure to digest it, it should be more bulky than the other meals. Stews, if well made, answer admirably, and may be supplemented with such additions as are obtainable. Because of lack of fuel, delay in the arrival of company wagons, the necessity of starting fires and doing the cooking after the halt is made, the evening meal is often late ; and if they can obtain other food many men eat it, often with little regard to its suitability. It is hoped that soup-carts, ambulance kitchens, and fireless cookers will correct this by having a ready cooked and hot meal prepared when the halt is made. Because of their portability, canned and preserved foods are much used on marches, and the usual precautions are necessary to see that no spoiled cans are

used. If this be done, they constitute fairly pleasant and very wholesome food-supplies, but they should be supplemented with fresh articles when these are obtainable. The general care in the preparation of foods that has been indicated as desirable in camps should be exercised in the short camps made by marching troops.

The government provides suitable clothing for all varieties of climate and weather, and the company commander should see that his men are provided with the proper kinds and amounts to **Clothing** meet the probable needs of the march. Flannel shirts should be worn even in tropical climates, as they prevent too rapid cooling of the body and afford protection from the night chill. Blouses or coats are rather an impediment than a help in the tropics, if flannel shirts are worn. The poncho should always be carried and the blanket nearly always, even in hot seasons, as the night chill is often penetrating and is accentuated if the clothing or the ground be wet. *Mosquito nets*, while not articles of clothing, may be mentioned here as most important in tropical countries, or hot weather. Nothing that the soldier carries is of greater value in proportion to its weight. It is his most important protection against yellow fever, malaria, and dengue; while it may be quite as efficient in protecting him from tick-bites, from the flies that infest with screw-worms, from poisonous insects and reptiles. Its use, though the importance of it has long been recognized, is apt to be neglected on marches.

It is not considered necessary to discuss in detail the various articles and kinds of clothing to be worn, but some few articles need discussion. The drill regulations allow the marching soldier three pairs of *socks*, one pair

on his feet, one in his blanket-roll, and one in his surplus kit on the company wagon. In order to keep his feet in good condition he should begin each day's march with clean feet in clean socks, and must always have a dry and clean pair of socks for emergencies, such as the accidental falling into mud-puddles or streams. It is therefore necessary that he should wash his feet and one pair of socks each night, and the company commander should make sure that he does so, and that, if possible, he then has facilities, such as a fire, for drying the socks. The socks should not only be clean and dry, but they should fit properly. If too long, they fold or wrinkle and make pressure or rub. If too short, they cramp the toes and may be as important in causing or aggravating corns, bunions, foot-cramp, and ingrowing toe-nails as poorly fitting shoes. They should be of such weight as is suited to the individual man, in general being rather light for foot-troops so as not to cause excessive sweating or burning of the feet. Any disorders arising from the use of improper socks or from other cause should at once be reported to the surgeon, who, by timely advice or treatment, may be able to avert lameness or disability.

Shoes likewise are frequent causes of lameness, and in many instances the company officer is to blame for not seeing that his men get proper sizes and fits. At other times the quartermaster's department is at fault in not having proper sizes for issue. In such event the man chooses a shoe of improper size, or buys them of improper shape and material from civilian dealers. The footgear of the army has improved in quality of late years, however, and most of the shoes now issued are excellent, both in shape and material. A proper shoe

should be sufficiently long and broad, so that the toes are not crowded even after a long march. It should not make painful pressure on any place, but should fit snugly and evenly over the instep and about the heel, to prevent rubbing or forward slipping of the foot. The heel should be broad and low. Two pairs of shoes are allowed, one pair on the feet and one pair in the surplus kit. If practicable, the day's march should always be begun in a clean and dry pair, and it is therefore important that a part of each evening's work should be the cleaning, drying, and oiling of one pair. For the last-named purpose, a piece of fat meat will answer if other oil is not obtainable. Shoes left wet and dirty soon become hard, wrinkled, and uncomfortable, and "run over" and break more readily than those kept clean and supple.

Two suits of *underclothing* are allowed, one on the person and one in the surplus kit. As underclothing is even more important as a protection from dirt than from cold, these will also require frequent washing. Neglect of this may result in attacks of boils, ringworms, and other skin infections. When a day in camp permits it, and coffee tins or other receptacles are available for the purpose, underclothing should be boiled.

Another article of the field kit that will require frequent washing and occasional boiling is the *towel*, but one being authorized.

For general purposes, and use in almost all climates, it is doubted if a more satisfactory headgear than the campaign hat has been devised. In arctic weather the fur cap, and in the tropics a cork or pith helmet will excel it at times, but for general use it is excellent. It should preferably, however, be freely ventilated and

worn "peaked" or uncreased, so as to have a large air space between the top of the hat and the top of the head. Much has been written as to the advantage of red or orange hat-linings and underclothing as a protection from the actinic rays of the sun. The value of such provision is not yet determined.

The general rules for camp police should be applied to marches as far as practicable, and they can be applied to a far greater extent than is ordinarily the case. In order to do this, proper instructions should be given in advance, and each company should have, as in camp, at least one man on sanitary police duty, whose business it would be to see to the proper disposal of waste matters, to prevent the careless and indiscriminate soiling of halting-places, to assist in preventing the use of improper water, and to do such other work of a similar character as the circumstances necessitate or the medical officer recommends. He should carry a spade. Men should not, except in urgent cases, be allowed to fall out except at regular halts. If the halt is a short one, the company commander indicates to the sanitary soldier a suitable spot for urine and feces. The man goes to that spot and remains there. All men needing to fall out go there and use the place indicated, afterward covering their dejecta with earth. If the halt is longer, half an hour or more, the sanitary soldier repairs at once to the designated spot and digs a shallow trench for feces and a shallow pit for urine. All men should be instructed to use these places only, and the trench and pit should be filled in again before the march is resumed. In case a man, from urgent necessity, does fall out on the march, he should be instructed to cover his feces with earth, using his

bayonet, if necessary, for the purpose. Similar rules should apply to men on outpost or picket duty, and in this way much may be done to prevent the surroundings of a camp from becoming an ill-smelling, fly-breeding nuisance. Scraps of food, greasy papers, and other articles thrown away by the men at the halting-place, should be thrown into one pit and burned or buried.

On arrival at camp, the sanitary soldier should immediately begin shallow trenches in a proper place selected by the medical officer, and men should be punished for urinating or defecating elsewhere. After that, the other business of making camp may proceed. Without it, the camp should not be made, as its site will almost certainly be defiled.

In bivouac, or while actually marching, the men should take all possible precautions to protect themselves from, or to rid themselves of, vermin and insects, and in warm weather the use of the mosquito net should not be neglected.

The hygiene of the battlefield has been largely set forth in the consideration of the camp and the march. All the general principles there set down hold good and should be observed as far as possible. Often, however, the military necessities allow no time to be devoted to sanitary matters; but even so, the soldier trained to a realization of the importance of such matters will be able to take better care of himself and to do less harm to his comrades than the man not so trained.

The man going into battle should, if possible, observe all of the following rules in addition to the general care outlined above: —

1. Make sure that you have a first-aid packet, that

**Battle-
field
Hygiene**

it is in good condition, and that you know how to use it.

2. Have your canteen full of boiled water.
3. Be clean in person and clothing, be bathed, and have on fresh underclothing.
4. Have your bowels and bladder empty.
5. Eat only good, nourishing, and digestible food.

All of these rules have an important hygienic bearing in case he should be wounded. The introduction of the first-aid packet in its present form was probably the greatest improvement in military hygiene ever consummated, and it has prevented an immense amount of suffering, mutilation, and death. For a large proportion of wounds, a properly applied first-aid dressing constitutes the most important part of the treatment, and, in many cases, the whole of it. With such a dressing at hand and a knowledge as to how to apply it, the use of water on wounds is very rarely necessary. But if water is used, it is most important that it should have been boiled. Otherwise it may infect the wound most seriously. Cleanliness of person and clothing are very important for the same reason, to prevent infections. Most infections of wounds are caused by bacteria derived from the skin or its coverings, and by simple cleanliness of these, surgeons are daily able to, and do, make extensive wounds that heal without a trace of infection. The importance of having the bowels and bladder empty, or relatively so, lies in the fact that wounds of these viscera are exceedingly dangerous, but that the danger is directly lessened with the chances of excremental soiling of the tissues. For this reason, eating, drinking, and moving, all of which increase or facilitate the escape of intestinal or bladder contents,

are to be avoided in the case of belly wounds, and for the man so wounded we may formulate another rule :

6. If wounded in the belly, lie down in the most comfortable position, eat nothing, drink little or nothing, do not move, do not get excited or restless. If overlooked or forgotten, you may be better off than if found and transported.

The sanitary disadvantages under which men travel on our army transports are numerous, but they are mostly beyond the company officer's power to remedy. The transport regulations provide for such sanitary measures and daily inspections as would seem to assure good conditions, and the conditions are such that with the aid derived from sea air and changes of scene the health of the men is usually well preserved or improved. *Over-crowding* is an evil frequently encountered on transports, and one from which others flow. The company commander is quite powerless to prevent this, but he should lessen the evils of it by having his quarters as freely ventilated as possible, having them policed each day, and vacated by both men and their belongings when opportunity offers. He should see that his men bathe frequently and have frequent changes of underclothing, and that they get suitable opportunities for clothes-washing.

Transports are not infrequently infested with *bed-bugs*. The company officer can only partially correct this evil by the police and cleanliness of his quarters, and, at times, by bed-bug hunts and the use of insecticides.

The *food* on transports is usually of good quality and very nourishing, but the methods of *cooking* and

**Transport
Ships**

serving it are almost always monotonous and unattractive, and result in unnecessary waste, improper methods of eating, and some disgust with the food. The company officer should try to influence these conditions for the better, so far as he can do so. Owing to the long confinement in exceedingly narrow limits during the trans-Pacific voyage, unremitting attention to all sanitary details is urgently demanded. All cases of sickness or disease of any kind should be reported to the surgeon early, in order that he may isolate contagious diseases, as well as give treatment. The two ills from which a majority of men suffer are sea-sickness and constipation, the former at the beginning of the voyage, the latter throughout it. The tendency to *sea-sickness* is lessened if the man goes aboard the boat in good general condition, with his bowels cleaned out and his system free from the poisons produced by over-eating, alcoholism, and other excesses. Constipation should be prevented, if possible, by the use of rather bulky foods, such as oatmeal, fruits, and vegetables, by daily exercise and a well-maintained habit of visiting the closet regularly and making the effort to defecate.]

Under varying conditions, troop trains may be composed of any kinds of cars that are in use. The gen-

**Troop
Trains**

eral principles of troop hygiene should be applied as far as practicable, whatever the accommodations. The most common faults in troop trains in time of peace are poor policing and dirty cars, and an insufficiency of water for drinking and cleaning purposes. Both of these faults are usually attributable to the railway officials, but the troops are partly to blame in that they are sometimes careless and wasteful of the water, and make very little

effort to keep the cars clean. The company commander can therefore partially control the evils. He should also make it his particular care to see that the car closets are well supplied with water for flushing purposes, and that the supply is replenished as opportunity offers.

CHAPTER VII

THE HYGIENE OF HOT AND COLD CLIMATES

THE general principles of hygiene are the same the world over and may be applied in any climate. They aim to keep the body strong, sound, free from infection, and at its maximum efficiency. Inasmuch, though, as the dangers of infection as well as of injury from extremes of temperature are somewhat different in the tropics and the arctics, it is well to give these conditions separate consideration.

Hot Countries

The danger to life and health is greater in nearly all tropical countries than in those that are temperate. This is due to a number of causes, among which we may note the almost total neglect or ignorance of sanitation among tropical peoples, the high, even, and often moist, temperature in which disease-producing organisms flourish and multiply, the great abundance and variety of insect-life, the common pollution of water-supplies, the habits of dress, and the heat itself. It is sometimes said that the safest procedure in matters of hygiene in strange countries or climates is to conform to the habits of the natives, but this is often a very serious error, and it has undoubtedly cost many lives. It is faulty habits in the natives that help to keep alive, in the tropics, many diseases that have long since almost or entirely disappeared from temperate climes, while the people sicken and die of preventable infec-

tious diseases in far greater numbers than should be the case.

The native of the tropics is very apt to suffer from water-borne diseases, and cholera and dysentery, although not limited to tropical climates, are so much less prevalent elsewhere as to be **Water** relatively rare, even if not unknown. The native is influenced in his choice of a water-supply by convenience, superstition, habit, and the fact that his ancestors did certain things, but not often by sanitary considerations. For this reason he suffers from the diseases mentioned, and partly for this reason he practically always harbors one or more varieties of intestinal worms. The only safe rule for general application to drinking-water in the tropics is that it should be sterilized by boiling, distillation, or passage through germ-proof filters. But this alone is not sufficient. The water must be kept sterile until consumed, by being put in sterile containers and closed against all possible contaminations. Both Filipino and Chinese servants in the Philippines are so entirely ignorant of the very meaning of sanitary precautions, that they are continually guilty of such action as stoppering bottles of distilled water with corks they have held in their mouths, wiping out a drinking-glass with a dirty rag or paper picked from the floor, drinking direct from bottles, and bottling unboiled water rather than take the trouble of boiling it, and it is feared that not all soldiers are free from some of these faults.

It is generally asserted, and is theoretically true, that less food is required in the tropics than in colder climates, and that the fats and proteids in particular should be considerably reduced. **Food**

This may be admitted, but we must also admit that

lifelong habits, exercise, and temperament exert an influence that cannot be ignored, and any arbitrary attempt to put American soldiers, especially if they be doing field-work, on a ration approaching the rice-and-fish diet on which many natives of the tropics live, will result not only in discontent, but also in impaired nutrition and in clandestine indulgence in food or drink of improper character. The present army ration is thought to be excellent for the tropics, as for home, provided that the men *work and exercise as at home*. If such is not the case, the ration is too heavy and should be reduced as indicated. The sugars and starches are well borne in the tropics, are not such "heating" food as fat and proteid, and do not tend, in the same way and same degree, to cast an excessive amount of work on the liver and kidneys. Neither do they offer opportunities for such harmful forms of intestinal putrefaction. Organic acids, such as fruit-juices, are especially valuable and important, as well as agreeable, in the tropics.

Care must be exercised in the use of fresh fruits and green vegetables, not because they are in themselves harmful, but because they are apt to be contaminated, from unclean water or from fecal manuring, with the parasites causing intestinal diseases. The only safe plan in using them in countries where dysentery, cholera, and intestinal worms abound is to have them sterilized: the green vegetables and some fruits by cooking; bananas, mangoes, oranges, and similar thick-skinned fruits, by thorough washing in boiled water. A large part of the food consumed by white people in the tropics is preserved by refrigeration or by canning. It is as good there as elsewhere, but is apt to

decompose rapidly after exposure to the ordinary temperature and atmospheric conditions, and should therefore be used promptly after such exposure.

Owing to the facility with which potatoes and similar vegetables spoil, the tropical resident should early accustom himself to the use of rice, yams, and similar products. The use of condiments and spices is generally more liberal in the tropics than elsewhere. This is, partly at least, a matter of custom, but it may have a physiological basis in that these articles are stimulant to the alimentary tract. They are not known to do harm.

Alcohol is quite as much abused by northern peoples in the tropics as in their own homes, or more so. In real moderation, that is, in the amounts that can be completely oxidized in the body **Alcohol** and used as food, it is not known that alcohol is more harmful in the tropics than elsewhere. In greater amount it is so, for the reason that alcohol in excess is in all climates an irritant to the liver, and in the tropics the liver is already hard-worked, irritated, "sluggish," and subject to congestions,—conditions aggravated by the irritation from the alcohol. Like the proteids and fats it does less harm if plenty of exercise is taken to promote its oxidation and the elimination of the combustion-products. As in other climates, its use is only infrequently indicated, and abstinence from it is advisable. Its very general, and often excessive, use among soldiers proceeds not so much from any needs of the system as from idleness, vacancy of mind, evil examples or careless associations, laxity of public opinion on such matters, and absence of such restraining influences as home-life and respectable female society. It

would therefore seem that the company officer can best combat the evil by attention to these points. Heat-stroke and heat-exhaustion are much more apt to occur in a man who is drinking, and to be more severe when they do occur. All drinking to excess is debilitating and probably lessens the resistance to infections. Recurrences of malaria and dysentery are not infrequently brought on by drinking-bouts or sprees. Alcohol also predisposes in more indirect ways to disease-infections, as by rendering a man neglectful of the quality of his food or the use of his mosquito net, and because he may use impure water with his drink, erroneously trusting to the alcohol to sterilize it.

In general terms, the dress should be adapted to the climate and to insure comfort and protection. The clothing issued to the army fulfills these requirements, allowing variations in the underclothing to suit individual tastes and requirements. The cotton khaki for outer dress is cheap, strong, durable, fairly cool, and easily washed. The flannel shirt for the field and that of blue cotton for barracks meet the requirements. British writers lay some emphasis on the desirability of always wearing woolen clothing in order to protect from chill, and much has been said as to the value of the woolen belly-band or abdominal binder. The writer's personal experience, and observations as to the general American experience, in the tropics do not indicate that this is a rule of general application. As stated before, most tropical diseases are infectious in nature, and if proper precautions be taken to avoid the infections, it is not thought that the discomfort produced by too great warmth of body, excessive perspiration, and prickly

heat is apt to offer additional protection. In the field, where the chill of night air is more keenly felt and where wetting may occur from rain or from fording streams, the flannel shirt should be worn, and it and the blanket meet the requirements. There is no objection to the use of light flannels by men who find them comfortable, while in the cold season and in high lands, as at Baguio in Luzon, they may be much needed; but for most of the service to which our men have been subjected in the tropics they are a source of harm rather than of good. Light-weight cotton undershirts, with short sleeves, and muslin drawers not coming much below the knees are preferred. The clothing worn in the tropics requires frequent washing and frequent changing, and a good-sized laundry bill is a necessity. The clothing is usually sun-dried, and that is an important aid in destroying germs on it. Most native wash-men do not boil it, but, if feasible, they should be required to do so.

The campaign hat, as previously stated, is an excellent article for field use, but it should be well ventilated by openings in the sides, and should be worn in such a manner as to preserve a large air-space above the head. The khaki cap offers very little protection from either heat or light, and should be superseded for garrison use by a straw hat or a helmet, the latter preferably of pith or cork. Much has been said and written of late about the value of protection from the actinic or invisible rays of the sun, and helmets with an interlining of metal foil, and hat-linings, underclothing, and spine-protectors of black, red, or orange material, have been advocated as means of protection from them. The value of such protection and of these means are yet in doubt and they

must remain so for some time. The evil effects ascribed to the actinic rays are principally those manifesting themselves as functional nervous disorders. Such disorders are notoriously hard to trace to a definite cause, and, on the other hand, the sufferers from them are the most favorable subjects for treatment by mental suggestion. Therefore, the fact that a number of such sufferers are benefited by the use of red underclothing is not conclusive evidence that their ills were caused by actinic rays.

Shoes should always be worn in the tropics as a protection against infections. Comfort and fit should be considered as elsewhere, but because it is comfortable to go barefooted is not sufficient reason for doing it. One of the greatest causes of sickness and debility in the tropics is the hookworm, and as it usually gains entrance to the body through the skin of the feet, the general use of shoes is looked upon as an important method of protection against it. Plague infection usually occurs through flea-bites on the feet and legs. Other infections occur in the same way; and the subject will be discussed at greater length in another chapter. Shoes and leggings also protect against leeches, ticks, mosquitoes, and other insects.

Because of the heat, the relative cheapness of native servants, and his comparative wealth, the American is apt to neglect outdoor exercise in the tropics. **Exercise** is. It is very important that he should not do this, as abundant experience has shown that it lessens the liability to the nervous breakdowns that constitute one of the chief causes of invaliding from the tropics. It is desirable that the glare and heat of midday be avoided as times for such exercise, the early morning or

the evening being preferred. The exercise should be incidental to interesting work or play. Tennis, polo, football, horseback-riding, and hunting are excellent forms. Marching is also good if the men are interested and cheerful. After the exercise, which should be sufficiently violent to produce a good sweat, the body should be bathed, well rubbed down, and dry clothing put on. Swimming is also a good form of exercise and one that should be encouraged, as every soldier should be a good swimmer.

With proper precautions as to exposure to the sun, the prevention of chill from wet clothing, and the avoidance of great fatigue, it is thought that exercise will do only good. Many white men in the tropics suffer from lack of it, very few, and they are usually diseased, from excess. Its value lies in the promotion of the active functioning of the skin, liver, and other organs of waste elimination, and the resulting improved nervous control, rather than in increase of muscular strength. Without exercise, alcohol, tobacco, and excesses in food are all doubly harmful.

Because of the freer perspiration and the greater liability to skin diseases in the tropics, more frequent baths and changes of clothing are necessary than in temperate regions. The bath water **Bathing** should not be sufficiently cold to produce shock or chill, but as such water cannot usually be obtained in the tropics, that feature need not require much consideration. Persons who have suffered from malaria or dysentery or other intestinal troubles should use tepid or warm water. It should be remembered that typhoid, dysentery, cholera, and certain infestations with animal parasites may be contracted by bathing in polluted

streams or ponds, and such places should therefore be avoided. Whatever the source of the water, care should be taken that it does not get into the mouth and so carry infection. After the bath the body should be thoroughly dried, particular attention being paid to the arm-pits, crutch, and groins, and, if these parts sweat profusely, it is well to dabble them with some antiseptic solution and then dry them again, after which they may be dusted with talcum powder. Fresh and dry underclothing must be then donned. If prickly-heat eruptions are present or appear easily, a weak formalin solution, about one teaspoonful of formalin to a pint or a quart of water, may be used to sponge the body after it is washed and before drying. It is thought that the measures here advised are as good as are known for the prevention of the prickly heat, small boils, chafing, and "dhobie itch" or ringworms that cause so much annoyance and irritation in the tropics. In addition, it should be remembered that some of these affections are contagious, and contact with persons having them should be avoided. Soldiers occasionally borrow articles of clothing from one another, and that is another method of spreading skin diseases. In Japan it is still a rather common custom for many people to bathe in the same tub or tankful of water, and that probably accounts for much of the abundant skin disease seen in that country. Such a practice, though economical of fuel and water, is to be avoided.

The heat, the bright light, the new and strange environment, the native habits of both dress and conduct, unite to stimulate and excite the sexual desires of the new-comer in the tropics, while his money and the native habits, poverty,

**Sexual
Hygiene**

and views of morality, constitute a set of circumstances enabling him to gratify them. In consequence he is apt to indulge in such sexual excesses as impair his nervous control, unless, as is frequently the case, he is soon or often checked by venereal disease. Later, when he becomes debilitated by his excesses or by other causes, his sexual vigor fails and he begins to worry about that, thus aggravating and accentuating his trouble. Nothing that is equally harmless so profoundly affects the mind of the average young man as the loss, even though it be temporary, of his sexual vigor, and nothing so prolongs and aggravates that loss as continual brooding over it; so in this combination we have the making of neurasthenia and melancholia. Each man should know that his sexual needs are not greater in the tropics than elsewhere, that sexual or other excesses may bring about nervous exhaustion and temporary loss of sexual power, and that the best rules in the tropics, as elsewhere, are: 1. Sexual continence; 2. Avoidance of obscenity and indecency; 3. Cleanliness of the genitals; 4. Complete avoidance of thought or handling of them, except for legitimate and proper purposes.

The necessity for thorough policing and cleanliness of houses and grounds is even more important in the tropics than elsewhere, because of the greater prevalence of vermin and of vermin-borne diseases. Malaria, yellow fever, dengue plague, and other insect-borne diseases, occur principally in the tropics, and are best combated by such methods of policing as do away with the feeding or breeding places of the insects carrying them. In addition, roaches and ants

**Police
and
Cleanli-
ness**

abound in numbers and with a persistency not seen elsewhere, and, as they run impartially in water-closets and food-chests, it is easily conceivable that they may transmit disease. Rats and mice constitute not only an annoyance but also a great and positive source of danger, because it is among them that epidemics of plague begin, and it is from such rat epidemics that those among people take their origin. Parasitic skin affections prevail very widely among persons, domestic animals, and fowls, and it is partly by cleanliness of habitation that we escape them.

All the usual sanitary precautions as to cleanliness, plumbing, and good policing are to be observed in the tropics as elsewhere ; but additional care must be given to all standing water both inside the house and out, to prevent mosquitoes breeding in it. Such collections should not be allowed if they can be avoided. If allowed to exist, the water should be completely emptied and renewed once in four or five days at most, or should be kept covered with a film of mineral oil, which can be renewed at like intervals. Ponds or other large collections should be stocked with fish, and the general precautions against mosquitoes, to be outlined later, should be observed. Mosquito-proof houses, while in quite general use in our own country, are still not in use by us in our tropical stations, and the fact is inexcusable. Our army practice has not nearly kept pace with our knowledge, and it is hoped that officers will spare no pains to have their barracks and quarters screened as a measure of comfort and of sanitation.

Very Cold Climates

The principal sanitary dangers menacing the dwellers in very cold climates, aside from lack of food and other things outside of our present consideration, arise from exposure to cold, resulting in frost-bite or freezing, from poor ventilation, from too intimate association with diseased persons or animals, from improper food or lack of fresh food. Infectious diseases are less numerous than elsewhere, as their germs do not find such favorable surroundings in which to grow, and there are fewer persons to harbor and distribute them.

The outdoor air of arctic regions is purer and freer from disease-producing organisms than that of other regions. Man-polluted air is as dangerous there, however, as elsewhere. As ventilation **Air** and heating are, from a sanitary standpoint, inseparable, and, as free ventilation makes heating more difficult because of the greater volume of air that must be warmed, it may be understood at once that the natural tendency in an arctic climate is to have poor ventilation. Every increase in ventilation lets in cold and allows heat to escape, and the discomfort of this is more acutely and keenly felt than that due to foul air; so the effort to keep warm may cause the air to become vile. As animal heat helps materially to warm a crowded space, and as crowding lessens the consumption of fuel, man and dogs, sick and well, are thrown into close contact. If one man in a crowded, ill-ventilated room has tuberculosis, diphtheria, tonsillitis, measles, smallpox, or other contagious malady, the contagium or poison is so concentrated as to make all present unusually liable to contract the disease.

Ventilation, therefore, should be as free as possible, care being taken to warm the incoming air. Crowding should be avoided, and as much time as possible be spent in the open. Particularly should intimate association in crowded rooms with sick persons be avoided. The intimate contact with sick and with animals is also productive of disease in other ways than that indicated above. A man having gonorrhœa or syphilis is in such circumstances much more apt to infect innocently and unknowingly those about him than he would be under other conditions. Echinococcus disease, due to infestation with a certain dog tapeworm, is almost limited to persons brought into such intimate contact with dogs as is here mentioned. Cleanliness is almost impossible under such conditions, and wounds are thereby very apt to become infected.

A pure water-supply is not such a rarity in the arctics as in the tropics, both because pathogenic germs do not flourish in arctic temperatures and because men are less numerous as sources of pollution. The danger of pollution of a supply is greatest in the spring and summer, when melting snow may wash into it the excrement that was scattered in the neighborhood during the very cold weather. Excrement is apt to be so scattered, for the reasons that other disposal may be more troublesome and that snow and freezing prevent its constituting a nuisance to the nose and eyes. Especial care should be exercised to prevent anything of the kind, and drinking water should always be taken from points above any possibility of such contamination. Attempts should not be made, particularly on marches, to quench thirst with snow or ice, as they detract from the local and general heat of the body, and are liable

to irritate the mouth and throat. Snow and ice may furnish the water-supply, but they must be melted and the water heated, when it may be profitably used in making tea, which is then refreshing, comforting, and stimulating, and adds to the warmth of the body.

The food-supply of the arctics consists largely of canned goods, and may be quite generous. Every effort should be made to supplement that diet

with fresh articles, whether of animal or vegetable origin, as scurvy is otherwise apt to make its appearance. It is generally thought that scurvy is due principally to deficiency of fresh vegetables, but a few years ago an antarctic expedition was saved from disaster from this disease by the addition to the diet of fresh penguin meat, and numerous other instances indicate that fresh meat may be as efficient as fresh vegetables

Food

• in preventing or curing it.

Because of the necessity of greater bodily heat-production in the arctics, more food is required, and, as meat and fat give rise to more heat during their oxidation, and as they are also more readily obtained, it is in them that the increase should be made. This need is recognized by the government, and is met by an increase in the ration for troops serving in Alaska. Where game or fish are obtainable they should be used to supplement the ration. The methods of preparing the food are not essentially different from those used elsewhere. Foods should be served hot, and the heat of the body so conserved.

Alcohol should not be used in the arctics except as a drug, as it increases heat-radiation and loss. Particular care should be taken that it is not used during the period of exposure to cold, as

Alcohol

that is the time when loss of heat is to be especially avoided. If used at all, it should be after the period of exposure, when the body is chilled and the surroundings warm and comfortable. In such circumstances it may be of considerable value, but its routine use should not be practiced.

Fur, feathers, and wool as conservators of heat, and canvas and leather as protection from wind, constitute the main clothing-supply and bodily protection in the arctics. Fur, leather, and canvas, with the exception of shoes, are for use out-of-doors only, wool for general wear both indoors and out, while feathers or down are used in bedding. Extreme cold is nearly always dry, and may therefore cause less suffering than higher temperatures when the air is moist. Wind of course greatly increases the suffering and the danger from either dry or wet cold. The reason that fur, feathers, and wool make such warm clothing is twofold; they are themselves poor conductors of heat, and they contain in their interstices a large amount of air, another poor conductor. For the latter reason a multiplicity of light garments is warmer than the same amount of material woven into one heavy garment, and cotton or silk wadding may be used also with good results. It is because they prevent the displacement of the warmed non-conducting layer of air within the clothing that leather or close-woven canvas affords such excellent protection against wind.

Fur caps and gloves, blanket-lined canvas boots, canvas overcoats lined with blanket or sheepskin, woolen outer and under clothing, woolen socks, oiled shoes, felt overshoes, and canvas leggings are issued for very cold service, and in far northern posts buffalo coats are still

furnished for use by men on guard. By proper use, these articles can be made to furnish adequate protection against any cold that is likely to be encountered. The parts most apt to suffer from cold are those most exposed and having the weakest circulation, such as the nose and cheeks, the ears, hands, and feet. The fur cap protects the head and ears, but not the neck and face. It should therefore be used with the overcoat collar turned up, or with the hood of the coat, in very cold weather. The blanket-lined canvas hood covers the neck and part of the face and therefore affords better protection. A veil or a strip of cloth may also be worn across the lower part of the face if necessary. The fur gloves adequately protect the hands except in the most extreme cold or in cases of long exposure. If necessary, woolen gloves may be worn beneath them. When, in spite of them, the hands become numb and cold, they should be beaten or exercised to warm them. For work not requiring the use of individual fingers, woolen mittens under canvas afford good protection for the hands. Care should be exercised about removing the warm hand from its glove to do delicate work in the cold. A soldier has been known to have both hands frost-bitten because he removed his gloves in the wind and in a temperature of 40° F. long enough to adjust his saddle. The feet are easily kept warm during walking if the government issues of footwear are properly used ; but when men are riding and not using the feet, or when snowshoeing and the feet are bound so tightly that the circulation is interfered with, the problem is more difficult. Except in wet weather, the foot-covering should not be impermeable, as the feet when comfortably warm are apt to perspire, and such a covering as a

rubber shoe causes the perspiration to condense against it and to be frozen there, so that the foot is almost or actually in contact with ice. An oiled shoe is less objectionable, as it does permit the vapor to escape, in part at least. In very dry cold, however, a felt shoe, or two or more pairs of woolen socks with canvas or soft leather moccasins, may be better still. Fur-lined boots or stockings are very warm. As free circulation of the blood is a necessity in the prevention of frost-bite, the feet should not be bound any tighter than the absolute requirements of the work demand, and the circulation should be stimulated by rubbing and kneading as opportunity permits. Wet feet should of course be dried and dry coverings put on them as soon as possible. Should frost-bite occur, it is important to remember that serious results are less likely to follow if the part is thawed and the circulation restored gradually, the beginning being made by friction with snow or cold water until the pallor and hardness have disappeared, and cooling applications being then applied, or the part left exposed to cool air which is very gradually warmed. A healthy and well-fed state of body are important in preventing freezing, and of course over-exposure on guard or otherwise is to be prevented if possible. Fatal freezing is preceded by languor, dullness, sleepiness, or stupidity, and men should not yield to sleep when very cold.

This is a form of blindness due to exposure to the glare of bright sunshine on snow. It is usually due to spasm of the lids and such sensitiveness to the light that the eyes cannot be kept open. The results may be very serious, and care should be used to prevent the trouble. Prob-

**Snow
Blind-
ness**

ably the most effective measure when exposure is not avoidable is the use of "snow eyes," or wooden spectacles with narrow slit-like openings through which vision is possible. Dark goggles are less protective, and have the additional disadvantage that they usually bring metal into contact with the skin and may thus cause freezing. Metal parts should be wrapped or covered with wool or fur. The practice of marching in close column and each man keeping his eyes fixed on the back of the man in front of him, the leading men falling to the rear of the column as their eyes tire, is also helpful in preventing it.



PART II

THE CAUSES OF DISEASE

"He discovereth deep things out of darkness, and bringeth out to light the shadow of death."

JOB 12: 22.



CHAPTER VIII

THE REMOTE OR PREDISPOSING CAUSES OF DISEASE

FOR our purposes it may be assumed that most of the diseases with which we are concerned are due to animal or vegetable parasites; but not every man exposed to infection by such parasites contracts the diseases caused by them. A special predisposition or liability is necessary, otherwise everybody would have tuberculosis, the entire army at Chickamauga in 1898 would have had typhoid fever, and all physicians would die of infectious diseases within a short time after entering upon their profession. This matter of liability or immunity to diseases is one of the most interesting, complicated, and important in the whole realm of medicine, and many of the world's greatest thinkers and investigators find it worthy of their highest efforts and constant attention. It is the purpose of this chapter to deal with these questions in some elementary aspects that should be known by the company officer, in a practical and simple way, without technicalities.

One of the most important of the factors concerned in predisposition to disease is heredity. Every man inherits from his ancestry something that renders him liable to tuberculosis, but im- **Heredity** mune to rinderpest and chicken-cholera. He possesses in a degree not possessed by any other known creature a liability to malaria, syphilis, gonorrhœa, yellow fever, cholera, smallpox, and other diseases, while he shares with many other animals his susceptibility to tuberculosis, suppuration, plague, and anthrax, and at the same

time he is not at all subject to some diseases that are very fatal to his domestic animals.

In addition to its influence in this general sense, heredity has long been credited with great influence in predisposing certain families or individuals to certain diseases. Consumption was long looked upon as a hereditary disease, but, with our more complete knowledge as to its nature, the tendency is now to regard its appearance and persistence in families as an evidence of familial contagion rather than of heredity. In the case of cancer, another disease prone to "run in families," the same explanation is now frequently advanced, though no germ-cause is known for the disease. The influence of heredity, though still considered weighty, is not now so much emphasized as formerly as a cause of insanity. In all these diseases, however, it is still contended that a predisposition, a type of tissues endowed with resisting powers below the normal, is inherited, even if the disease has an extraneous cause. Disease can at times be born with the child, and not manifest itself until later, the actual disease germ being present in the body all the time. Syphilis is the ailment most often showing this. Other diseases, such as typhoid and smallpox, may be contracted in the uterus, and the child may be born sick. It is probable that sometimes disease may be suffered and recovered from in the mother's womb, and the attack confer immunity through later life.

Certain persons appear to possess all their lives an immunity to certain diseases, never contracting them in spite of frequent and direct exposure.¹

¹ Captain C. F. Craig and the writer were quite unable to infect certain individuals, who volunteered for the purpose, with dengue, using methods usually successful and injecting doses of infected blood many times as great as would be received in natural infections.

Certain nervous affections have no known cause but heredity, the disease appearing in the affected family with great regularity. "Bleeders," persons who bleed excessively or fatally from very trifling injuries, in most instances inherit the tendency, and in a curious way, as it usually descends only *through* females to manifest itself *in* males. The drink habit is often spoken of as inherited, but this is probably not the case. An unstable nervous system, which allows its possessor to fall more readily a victim to drink or other excesses, is inherited. It is common observation that types of body are inherited, that children look and are built like their parents, and it is but natural that they should be predisposed to the same diseases, so far as bodily conformation exercises any influence.

That this does exercise an influence is indicated by the table quoted on page 5, as well as by popular belief. The common remark that such a man looks as though he might have apoplexy any day, or such another man will probably die of consumption, are based on facts long noted by physicians and the public. It is partly for the purpose of eliminating persons so predisposed to disease that the government demands not only that each man shall be in good health at the time of his enlistment, but that he shall also conform to certain standards not having much bearing on his present efficiency.

Certain diseases are known generally as those of childhood, as they are so rarely seen in persons beyond that age. Scarlet fever, measles, whooping-cough, and diphtheria may all occur at other stages of life, but it is with ever-decreasing frequency that they do so, and the phrase is, in general,

**Types of
Body**

Age

correct. On the other hand, cancer, apoplexy, cataract, and several other troubles are so nearly confined to aged persons that they may be called diseases of old age. Certain infections are seen principally in early adult life, so that it is the young soldier that usually shows them in the military service. Among them we may mention typhoid, gonorrhœa, syphilis. The question of age is related to that of recklessness and folly, and it is partly for this reason that the young are in general more disposed than the old to infectious diseases. The prospects of recovery from an illness are also influenced by age. Pneumonia and injuries are much more serious in an old than in a young man. On the other hand, diabetes and epilepsy are not usually so serious in elderly persons as in children.

Certain persons are born with peculiarities of person that render them particularly liable to certain diseases, and no adequate explanation can be offered for them. Some persons cannot eat fish, others strawberries, others eggs; and if any of these overcomes his dislike and forces himself to eat of the forbidden thing, he is made sick, possibly dangerously so. A large number of persons are born with such a susceptibility to the effects of certain pollens that they have to leave their homes when these pollens are ripe, or suffer from hay fever. Some men suffer from asthma if they go about a stable or ride behind a horse, others are made sick or faint by the presence of a cat or the odor of its urine. Of two healthy men, one may suffer great discomfort after taking a tenth of a grain of iodide of potash, while the other can take three hundred times as much and not suffer at all. It is not impossible that similar peculiarities

**Personal
Pecul-
iarity**

influence the development and course of infectious diseases.

Worry, fear, and homesickness strongly predispose to, if they do not actually cause, mental and nervous diseases. They also appear to predispose to infectious diseases, possibly by lowering the general resisting powers and influencing the circulation.

**Mental
States**

The aphorisms of all languages recognize the value of training in almost any occupation or pursuit in life.

Military authorities recognize it by the fact that they require the soldier to spend years

Training

in training for an emergency that may never arise, or that, having arisen, may last but a day or an hour. It is not less valuable in sanitary matters than elsewhere. The child or the man who is trained to reason, to obey, and to conduct himself properly, is in much less danger from infectious diseases in camp or elsewhere than is the reckless, disobedient, or headstrong individual who knows no law but his own will and appetite. It is the latter who drinks bad water, eats poor food, contracts venereal disease at each opportunity, urinates and defecates in forbidden places, avoids vaccination, seeks alcoholic indulgence, sleeps without his mosquito-net, spits on the floor, and is otherwise a source of much worry and mischief. It is because of this lack of training in hygiene that recruits are so apt to become sick when brought into camp, and, what is worse, to scatter their sickness in all directions. It is this same lack of training that constitutes the greatest weakness and the greatest evil of our military system to-day, and gives origin to the fear that when we again mobilize an army of a quarter or a half million of men, — more than the

regular medical staff can oversee,—the conditions of 1898 may be repeated.

Lack of training on the part of troops is the most important predisposing cause of disease in armies. It is the duty of all officers, of whatever rank or branch of the service, to endeavor to overcome this defect, for which they are mainly responsible. The fact that a man is ignorant or headstrong at the time of his enlistment does not justify his being so a year or three years later. Medical officers should be required to give lectures or other instruction, but the company officer should see that example and wise precept are so constantly before the minds of his men that hygienic living becomes a matter of habit.

Aside from habit in the large sense just used, many small and unimportant personal customs have an influence in predisposing to disease. Eating raw meat predisposes to infestation with trichinæ or with tapeworms, carelessness in hand-washing increases the liability to many infections and poisonings. The habit of going barefooted may be of very great importance in predisposing to plague and hookworm disease. Careless habits in the disposal of waste are responsible for so many sanitary ills as to make it evident that the main purpose of sanitary training is the establishment of proper habits.

The influence of race as a predisposing factor of disease is closely related to that of heredity, habits, and environment, but race in itself exercises some influence. The reason for this is not always known, but in some instances it is probably due to long racial exposure to certain diseases, whereby only the more resistant strains or families survive. At

other times it is probably the case that the disease in question may be uniformly incurred in childhood, and so confer protection through later life. Thus, the question has been raised whether the present repression of yellow fever will not permit the growth of a generation of Cubans having no natural immunity to the disease. Whatever the reason, Cubans and West African natives are less susceptible to yellow fever than Americans and Europeans. Jews are especially subject to diabetes, and negroes are much more apt than other peoples to suffer from keloid, a form of tumor resembling overgrown scars.

Exposure to inclemencies of weather acts as a predisposing cause to numerous diseases. Tonsillitis, pneumonia, rheumatism, and influenza are all germ-diseases, yet the frequency with which an attack of any of these is ascribed to wetting or cold shows that these influences are not without effect. The germs causing pneumonia, diphtheria, spinal meningitis, and tonsillitis may be present in the mouths or noses of healthy people and produce no symptoms, yet after exposure the person may sicken. In a similar manner a person may carry the organisms of malaria or dysentery, and only become conscious that he is ill after a wetting or chilling.

Injury predisposes to very many diseases. Pneumonia, meningitis, and typhoid, of general diseases, may follow injuries in such a manner as to appear related to them. Abscesses, kidney and liver troubles, and tumors may follow injuries that do not break the skin, while anthrax, erysipelas, most suppurations, "blood-poisoning," hospital gangrene, lockjaw, hydrophobia, and some other affections

Exposure

Injury

are preceded by breaks in the skin in nearly all instances, though these may be so small as not to be noticed at the time. It is through the wounds that the disease-producing organisms gain entrance to the body. In the strict sense it is also through wounds — insect bites — that malaria, yellow fever, plague, and sleeping sickness are transmitted. The great class of suppurations, however, is the principal diseases following injury, and the proper use of the first-aid packet for the purpose of preventing them has almost done away with one of the greatest horrors of war.

Injury may also act as a means of diffusing and generalizing an affection previously isolated and relatively harmless. It may rupture an abscess due to appendicitis, and so give rise to general and fatal peritonitis, or it may loosen the infected clot in an inflamed vein, and cause acute general "blood-poisoning."

As is shown in Part I of this book, the soldier's environment includes so much that it must exercise an

**Environ-
ment**

influence in predisposing him to, or protecting him against, nearly all sorts of diseases. It is so large a matter that it cannot be adequately mentioned in a paragraph, and the whole subject of hygiene touches on it. The consideration of some phases of it occupies the remainder of this chapter.

There are quite a number of "occupation diseases," so called because their development depends on injury due to the employment of the individuals.

Occupation

Among such diseases may be mentioned writer's cramp, painter's colic, chimney-sweep's scrotum, housemaid's knee, and many others. During and after the Civil War a certain type of dilated and irritable heart was known as soldier's heart,

and in the stress of a hard campaign many more cases may develop. In the early months of 1899, when the State troops were doing much hard marching in the Philippines, many such cases were seen. Occupation has a great effect, however, in other and less obvious ways. Thus it may expose a soldier to mosquito bites and so predispose him to malaria, yellow fever, or dengue; it may put him on guard in a prison, where he will contract typhus or relapsing fever. Men working about stables are more apt than others to suffer from lockjaw after wounds, as the germ occurs in horse-manure. Occupations involving the breathing of much dust, such as marble-cutting, scissors-grinding, and some weaving, predispose to tuberculosis. Sedentary occupations may impair digestion, allow muscles to atrophy, and cause hemorrhoids. Callings involving great strains or exertion predispose to diseases of the heart and blood-vessels. It is stated that the majority of Japanese jinricksha men are dead or invalids from circulatory diseases before they attain the age of forty. Occupations causing much worry or nervous strain predispose to nervous exhaustion. Overwork in almost any calling may predispose to disease, either by lowering or exhausting the general resisting powers of the body, or by exhausting the nervous supply.

The quality of the air-supply is very important as predisposing to disease, by its temperature, dryness, or purity. Too great heat, especially if moist, causes much discomfort and may result in **Air** heat-exhaustion, heat-stroke, and death. Rooms that are too hot cause the men to sweat and to expose themselves unduly to cold or draughts, which may result in congestions of the lungs or kidneys. Rooms that are

too cold allow the body to become chilled on the surface and congested internally, making easy the development of coughs, sore throat, and other troubles. In barracks, however, such troubles are usually more closely related to foulness and contamination of the air than to its temperature. One case of sore throat in a squad-room may suffice to contaminate the air and expose every man in the room to disease if the ventilation is poor. This condition is aggravated with every increase in crowding. Carbon dioxide and other waste-products of the body may also make the air poisonous without considering infection.

Foggy or damp air predisposes somewhat to rheumatism. That which is too dry causes dryness and irritation of the respiratory surfaces. Dusty air introduces many infections, among them tuberculosis, and it is for this reason that spitting on floors or streets is very dangerous. The spit dries and is pulverized, and eventually becomes dust, in which condition it may be blown or inhaled into sound noses or throats, carrying with it the disease-producing germs. Dust-laden air is responsible for many more "colds" than is cold itself.

The use of contaminated water predisposes, of course, to water-borne infections. In addition, the use of very

Water hard water or of water containing chemicals may so disturb the digestive organs as to make infection easier. Excessive water-drinking disturbs the digestion and causes unnecessarily profuse sweating and urination; deficiency of it causes irritation of the kidneys and bladder, constipation, and general insufficiency of waste elimination. Greater insufficiency of course causes direct suffering from thirst.

Aside from the very numerous infections conveyed

by uncooked or improperly cooked foods, which are mentioned at greater length in other places, the kind, amount, and preparation of the **Food** articles used has so great an influence on health as to make the subjects of catering and cookery worthy of the company officer's best thought. He should learn what foods are good, their relative digestibility, food-values, cost, and the amounts in which they should be furnished. The method of preparation should be such as to make them attractive and digestible. The ration forms an excellent basis.

Excess of food predisposes to stomach and intestinal disorders, to gout, obesity, vascular and kidney diseases, and to various forms of auto-intoxication, or self-poisoning by waste products. Insufficiency of proper food leads to weakness, loss of flesh and a lessened resistance, that renders infections more easy. Great epidemics of typhus and relapsing fever have followed famine and deprivation so many times as to earn for them the title of famine fevers.

Alcohol is the direct cause of such diseases as delirium tremens, alcoholic neuritis, and gastric catarrh; but it is also a predisposing cause of many other **Alcohol** diseases, — of the blood-vessels, heart, kidneys, and brain. It also predisposes to infections, both by making its user careless in regard to them and by lowering the resisting powers. Drunkards are especially subject to pneumonia, and are also especially unfavorable subjects for that disease. The chronic gastric catarrh of the old alcoholic may make him an easy victim for cholera or typhoid. Alcohol predisposes to heat-exhaustion in hot and to freezing in cold climates.

Insufficient or excessive clothing may predispose to disease as do heat and cold. Insufficient covering in the tropics may lead to severe sunburn, or, if **Clothing** of the head or spine, predispose to heat-stroke. Lack of foot-covering not only leads to injury of the feet, but predisposes to diseases that usually enter through the skin of the feet, including hookworm disease, chiggers, Guinea worm, and plague. Insufficient clothing may also predispose to mosquito-borne diseases. Excessive clothing may predispose to heat-stroke, exhaustion, and diarrhœa, and this is especially true when men are turned out for inspection or drill in too heavy uniforms.

Warm baths, if followed by exposure or chilling, may predispose to disease, as may cold ones in persons who do not react promptly. Insufficient bathing **Bathing** predisposes to skin diseases, especially in hot countries; bathing in polluted waters, to water-borne infections. In addition to typhoid, dysentery, cholera, and common intestinal worms, there is reason to believe that *schistosomum* disease — infestation with a worm that lives in the veins of the liver, and causes bleeding and other serious bowel and bladder symptoms — may occur during bathing, the young parasites entering through the skin. This disease is widespread in Africa, and varieties of it are seen in the West Indies and the Philippines, so that it may yet assume importance with us.

The use of dirty water also predisposes to skin diseases, as noted in regard to the oft-bathing Japanese. The exposure incident to bathing in streams may increase the liability to diseases conveyed by mosquitoes and other insects.

Natural sleep in proper amounts is an important factor in the prevention of disease ; the lack of it causes weakness, exhaustion, and depression that render infection and injury easy matters. **Sleeping**

Insufficient sleep for a long period of time causes nervous disorders and predisposes to insanity. Sleeping in improper surroundings may greatly increase the liability to disease through chilling, wetting, exposure to mosquito bites, and in other ways. Sleeping in close contact with persons suffering from certain contagious diseases would strongly predispose them, as would sleeping in the beds or clothing of such persons.

Poor sanitary police is a feature of poor military training, and predisposes to all of the epidemic forms of infectious diseases. It does this by allowing infected dust and papers to blow about **Police**

and pollute the air, by allowing soil and water contamination, by leaving or providing breeding-places and food for flies, fleas, mosquitoes, rats, and other vermin, and by failure to remove sources of infection. In these ways it encourages the development of typhoid, cholera, dysentery, yellow fever, plague, pneumonia, influenza, tuberculosis, and other diseases.

By attention to police and general cleanliness of houses and neighborhoods some diseases that formerly occurred in great epidemics have been almost banished from civilized countries, and even famine does not bring them forth as in earlier times. Typhus and relapsing fevers are striking examples. Tuberculosis and other diseases have been reduced in part by similar measures.

As the best police is possible only when all unite in striving for it, it is essential that line officers and enlisted men should become interested in the sanitary

importance of the question, and should look on good policing as a source of pride, comfort, safety, and efficiency.

A poor camp-site may constitute a predisposing cause to many kinds of disease. It may expose the men to fog,

Camp- unnecessary dampness, wind and cold, or to
Sites dust and heat, to mosquitoes, flies, fleas, bad water, or contaminated soil, and through

these factors promote the development of the worst camp-diseases. While it is highly desirable, therefore, to select as good camp-sites as can be obtained, it must ever be borne in mind that proper sanitary precautions may nullify the evil effects of a bad site, and that neglect of them is almost certain to destroy the value of a good one. Polluted sites, those that have been much used by troops, and those that expose the men to bad water and disease-bearing insects are particularly to be avoided, as these factors predispose so strongly to various diseases that only the best trained and disciplined troops may come off unharmed.

Practically every case of infectious disease is derived from some other one, and it is therefore obvious that

Presence the existence of one case constitutes a fac-
of Disease tor predisposing to more. One man with typhoid, cholera, smallpox, or plague, in a camp, constitutes a danger to the entire command, and this fact is so well recognized in these and other diseases that it forms the basis of the entire system of quarantine and isolation. The origin of individual cases of infectious diseases may be difficult to trace, the typhoid-carrier, the unsuspected syphilitic, or the man with a slight diphtheria, not even realizing that he is sick, and unknowingly spreading disease broadcast. One

disease may predispose to the development of another or to new manifestations that are spoken of as new diseases. Thus blood-poisoning, gonorrhœa, typhoid, and, more particularly, rheumatism, predispose to valvular heart-disease, syphilis to certain affections of the nervous system, diphtheria to varying forms of paralysis, dysentery to liver abscess. Other instances might be detailed in considerable numbers, but enough has been said to indicate that disease is a factor that may predispose to other diseases.

CHAPTER IX

THE IMMEDIATE OR EXCITING CAUSES OF DISEASE

THE immediate or exciting causes of some diseases, among which may be mentioned such important ones as scarlet fever, measles, yellow fever, and typhus, are unknown, but those that are known are so numerous and of such varied character that they could not possibly be discussed in this chapter, even if it were desirable that they should be. We will therefore omit all discussion of the causes of large and important groups of diseases, and consider only briefly the various classes of disease-producing factors which it seems it would be interesting or profitable for the company officer to know. Numerous as they are, such factors may be gathered into a few groups.

Of these groups, the first to be considered is that of mechanical causes, and, of such, *traumatism* or violence is probably the most important. Like the other components of this group, it may predispose to disease, but it is also the direct or exciting cause of many ills, among which we may class practically all varieties and instances of wounds, from the slightest abrasion to the most extensive shell-wound, fractures, dislocations, and bruises. Concussion, laceration, or rupture of internal organs, such as the brain, spinal cord, liver, kidneys, stomach, intestines, or bladder, may result from blows or other injuries that cause no break in the skin, or even, in rare instances, leave no external mark. At times the violence may be

Mechanical

slight in degree, and apparently out of all proportion to the resulting damage. Thus, a relatively slight blow on the chin may be so directly transmitted to the brain, and so jar and shake it, as to cause immediate and deep unconsciousness; or a blow over a distended stomach or bladder may rupture it and cause death, to the great surprise of all witnesses. Or the evil effects of the injury may not be manifested at once. A knee or hip, for instance, may be injured and give rise to no symptoms at the time, but later cause prolonged suffering or lameness; or a man may receive a severe blow on the head or the abdomen, and continue at his work all day, yet he may have received an injury from which he will die in a few days. Officers should, therefore, exercise much care and forbearance before they pronounce a man a malingerer because some accident from which he suffered appeared trivial or harmless. Violence may be so transmitted as to manifest its effects on a distant part. Thus a man may fall from a height and land on his feet, and yet fracture his skull; or he may fall on his hands and fracture his collar-bone; or he may receive a blow on the right side of the head and suffer a laceration of the left side of the brain. As the manifestations of violence may be almost infinite in variety, so may its effects. *Pressure* may also cause disease in a variety of ways. Applied to the feet it may cause deformity, bunions, or corns; to the waist, displacement of internal organs; to a nerve, as by a bullet or a bone-fragment, paralysis or great pain; applied to a blood-vessel, it may cause ulceration and rupture, with fatal bleeding, or it may cut off the blood-supply of the part to which the vessel runs, and cause gangrene or death.

Heat and cold, besides predisposing to disease, as shown in the preceding chapter, may directly cause injuries varying from the slight reddening of the mildest burn or chilblain to the loss of entire limbs, or of life itself. They, as well as violence and pressure, in addition to the immediate damage done, may cause lifelong suffering or deformity, from the formation or contraction of large scars, or by means of secondary infections.

Increase or diminution of atmospheric pressure may cause disease or death, the former in divers or caisson-workers, the latter in mountain-climbers or balloonists. In the case of increased pressure, the trouble most often comes from too sudden return to normal pressure, which allows the liberation of nitrogen gas in the blood.

Electricity may cause disorders varying from slight pain or nervous disturbance to deep burning or sudden death. Mechanical injuries that are not usually thought of as such are exemplified by the rashes or irritations due to handling or contact with some *caterpillars* or *moths*, and are caused by the introduction and retention in the skin of the minute barbed hairs that partly cover the creatures.

The group of chemicals that cause disease or death is very large and embraces practically all poisons. They may occur in any form, gaseous, liquid, or **Chemical** solid, and may enter the body in any way, as by inhalation, by mouth, by the rectum, through the skin, or through wounds.

Gaseous poisons usually enter the system by inhalation. Those most commonly causing trouble are carbon dioxide, water gas, coal or charcoal gas, or carbon monoxide. These and other gases may produce death

without occasioning great suffering. Another group, including formalin, ammonia, chlorine, and sulphurous fumes, cause marked irritation of the respiratory tract and excite efforts at escape, so that they are, in that respect, less dangerous. Certain others, such as some compounds of arsenic or phosphorus, may cause poisoning by inhalation in small amounts, and be very difficult to trace. Chloroform and ether usually, and wood-alcohol and some other liquids occasionally, enter the body in a vaporous or gaseous form. Gases may produce death in any one of several ways: by acting mechanically to exclude oxygen, and so cause suffocation, as in the case of nitrogen; by forming combinations with the blood to prevent its taking up oxygen, even if it has the opportunity, as in the case of charcoal gas; by causing paralysis, or change of nerve-tissue, as in the case of chloroform or wood alcohol; by causing such irritation of the glottis as to close it by swelling, and so induce suffocation, as does ammonia.

Liquid poisons may also cause injury or death in a great variety of ways. The group includes a large number of substances. Some of them, such as *sulphuric* and *nitric acid*, do injury by their strong corrosive properties; others, as alcohol, act on the nerves. They usually enter the body by way of the mouth, but, as in the case of the acids, they may act on the surface, or enter through wounds.

The poisons occurring in *solid* form are even more numerous than the others, and they enter the body in the greatest diversity of ways, and produce manifold symptoms. *General poisons* include a great variety of substances, such as arsenic, phosphorus, antimony, mercury, lead, and a great many more. They may be taken

knowingly with good or evil intent, or in ways not known at the time, and often difficult to trace. The amount taken at one time may be almost infinitely small, yet its frequent repetition may result in poisoning. Thus, a painter may contract lead-poisoning from the minute amounts of lead getting on his food from unwashed hands, a beer-drinker from the lead taken into solution from lead pipes through which the beer is drawn.

Alkalies, such as *caustic potash*, *caustic soda*, *lye*, and *quicklime* are powerful irritants or caustics.

Vegetable poisons include substances producing disease or death in a great variety of ways. Some of the most common and familiar of these are *opium*, *strychnine*, *cocaine*, and *jimpson weed*. Most of the enslaving drugs — alcohol, opium, cocaine, and others — are of vegetable origin. Habitual users of them may get to using enormous doses, some of them several times the amount that would be fatal to persons not so habituated. Some vegetable poisons are very irritating to the skin and cause marked eruptions. Croton oil causes pustules that resemble those of smallpox. Poison ivy owes its evil qualities to an oil that occurs in and on the leaves and that causes the well-known inflammation of the skin.

Many of the chemical poisons are the products of germ action. Alcohol, the best known, results from the action of yeast cells on sugar. *Ptomaines* result from the action of germs on meats, fish, milk, and other food-stuffs. There are a great many of them, and their effects vary greatly. Some of them resemble drugs, such as morphine, strychnine, or atropine, in their action, and, as their presence may not be suspected at the time they

are taken into the system, the symptoms produced by them may prove very puzzling. It should be borne in mind that ptomaines may not be destroyed by boiling, and cooking should therefore not be depended upon to render partially decomposed meat safe. Many of them are harmless, but neither can that fact be depended upon to show that a given article of food is wholesome. Their poisonous character depends on the type of organism producing them, rather than on the stage of decomposition. Thus one piece of meat may be quite rotten, and yet not poisonous, while another that is free from odor and presents a fairly good appearance may be very dangerous.

The foods that most often give rise to *ptomaine poisoning* are those that have been preserved imperfectly, such as beef or fish in cans that have become perforated or blown, cold-storage beef that is not well frozen or chilled, poultry imperfectly chilled, especially if it be undrawn, and imperfectly cured hams and sausages. The symptoms vary considerably, depending on the source, character, and amount of the poison ingested, but in most instances they include evidence of marked gastro-intestinal and nervous disturbances, such as vomiting, diarrhœa, headache, dizziness, and cramps. *Self-poisoning* or *auto-intoxication* may be due to ptomaines or other decomposition products resulting from the putrefaction of food, occurring in the stomach or bowels, or to simple retention and absorption of poisonous products which would ordinarily be eliminated, but which, because of constipation, kidney disease, or for some similar reason, remain in the body. Such poisonings are promoted by overeating, lack of exercise, and by excessive drinking or smoking, as well as by diseases.

Animal poisons, in addition to the products of animal waste to which reference has just been made, include such substances as *snake-venoms* and those of stinging insects and spiders. These vary greatly in their composition, and their effects range from the slight itching and burning of a mosquito bite to the agonizing deaths following some snake bites. The venoms are introduced through wounds, many of them being harmless if applied to the unbroken skin or even if taken by mouth.

The most important group of the larger animal parasites that infest man is that of the intestinal worms.

**Large
Animal
Parasites** Their distribution is world-wide, and in some places they are so prevalent and their effects so serious, that they exercise very important influences on the morbidity and mortality rates. They may be grouped in three classes: the flukes or sucking-worms, the tapeworms, and the roundworms.

Flukes or *sucking-worms* are not common parasites of man in our own country, but in some parts of the world they are very prevalent and cause serious illness. Several varieties occur in the Philippine Islands, and some of them cause death. One variety occurs in the lungs, and causes blood-spitting and cough that may lead to the belief that tuberculosis is present; or it may lodge in the brain and cause paralysis or other cerebral symptoms. Other varieties live in the veins of the liver and may produce disease of that organ, while their sharp-pointed eggs lodge in or perforate blood-vessels of the bowels or bladder and cause bleeding, irritation, and other symptoms. This is a type of the *schistosomum* disease mentioned in the preceding chapter. Although it is claimed that this particular class of flukes may enter

the body through the skin, this is not yet a generally acknowledged fact; and the usual history of the flukes is that they pass a part of their lives in the bodies of snails or other small water-animals,¹ and enter the human body only by way of the mouth. Even though both methods of infection be possible, the obvious means of prevention are, first, to prevent infected or egg-bearing feces or urine from reaching water; second, to boil or otherwise sterilize the drinking water; third, to abstain from bathing in infected water unless it also be boiled or filtered; fourth, to avoid the use of uncooked foods that may have been contaminated by bad water.

The *tapeworms* are better and more widely known as human parasites. There are many varieties of them, but the best known are those large species ingested with beef and pork, though others are important, and some of these, such as the broad tapeworm from fish, may be very important in places. The life history of tapeworms, while not so complicated as that of flukes, involves, in most instances, residence in two animals, and knowledge of it enables us to understand the manner in which infestation occurs, and the method of avoiding it. Life begins in the eggs, which are passed from the worm into the bowel-contents of its host, and thence to the outer world. Here they may fall into water or on vegetation, or the excrement itself may be ingested by animals. At any rate, the eggs must, in one way or another, get admission to the alimentary canal of the second host, for which purpose a particular species of animal is necessary. In the case of the beef tapeworm,

¹ Flukes in general, so far as their life-history is understood, have at different times four forms or stages and two or three methods of multiplication.

this host is the ox, in that of the pork tapeworm, the pig. Rarely this development may occur in other animals, or in man himself, but ordinarily development will not occur in any but the usual host. Having reached the stomach of this host, the embryo is set free when its enveloping shell is dissolved, and penetrates the wall of the stomach, whence it passes by the blood vessels to the liver, muscles, brain, or other part of the body, where, after undergoing some further development, it becomes the head and neck of a future worm, inclosed in a small bladder, like cyst or measles, from one fourth to three fourths of an inch in length and about one third of an inch in diameter in the case of the pork measles, and somewhat smaller in the case of the beef measles. If now the flesh of the infected pig or ox be eaten raw, or not cooked sufficiently to kill the worm, it may pass unharmed to the bowel of man and there attach itself and grow. The head and neck as ingested are very small, bearing comparison in size to a small pinhead and a piece of fine twine. From this head and neck the rest of the worm develops in the bowel. The part so developing consists of a great number of segments, at times hundreds, each flat and white, so joined together as to form a long flat ribbon that may attain a length of thirty or more feet, each segment having a nervous and alimentary system and male and female generative organs, and each developing, fertilizing, and discharging eggs, so that each tapeworm, though an individual, may be said to be also a community. It may be readily understood from the above that allowing pigs to act as scavengers of human ordure, as is often the case in the tropics, is not a wise sanitary measure. It was stated above that man might rarely be the measly host. In such cases he might

become infected with eggs by means of contaminated water, or of lettuce or other green vegetables manured with human feces, by getting the minute eggs on his hands in water-closets or from his own stool, or by having them pass from the bowel to the stomach during vomiting. Several other varieties of tapeworms may infect man, — one of the most dangerous, and one for which man may shelter the cystic stage with fatal results, being derived from the dog, and another dangerous one possibly being derived from the rat, which may deposit the eggs on food articles with its feces.

Many varieties of *roundworms* infest man, and their life-histories vary greatly. A few varieties will be briefly discussed in order to indicate more clearly the necessity of certain sanitary precautions. Those most commonly inhabiting the human intestine — *the large, white roundworm*, the *whipworm*, and the *pinworm* or *seat-worm* — require no other host, and infection occurs, directly or indirectly, from man to man. The eggs are laid in the feces and pass from the anus, whence they may get on the hands, into drinking water, or on green vegetables, and so obtain entrance to the mouth and stomach of the same or another person. The embryos are set free in the stomach or intestine and develop into adults. Probably the most common method of infection is by means of polluted water. Children and adults who are infested with seat-worms may reinfect themselves many times, by reason of the fact that these worms often crawl from the anus and give rise to most intense itching, in the efforts to relieve which the hands and nails pick up many of the microscopic eggs, which may later be transferred to the nose or mouth.

The *hookworms* are probably the most important of

all the worms infesting man, both because they may and do infect the great majority of the inhabitants in some tropical countries, and because they so seriously reduce health and strength. In Porto Rico, the Philippines, and our own Southern States, infestation with these worms is common, and thousands and thousands of persons are debilitated or incapacitated, and have their resistance to other disease reduced by it, with the result that industrial progress is delayed and physical degeneracy and high sickness and death-rates promoted. Formerly it was believed that infestation with this parasite occurred in the same manner as with those just described, but it is now generally admitted that it usually, if not always, occurs in another way. The usual history is about as follows: The adult female produces a great number of eggs that pass out from the host with his feces. If passed on moist, warm earth, the eggs hatch and liberate microscopic embryonic worms which eat feces or other organic matter, live and grow in the moist earth, and eventually, if fortunate, find an opportunity to get on the bare feet or other skin-surface of a person, and at once penetrate the hair follicles and the true skin, producing itching and irritation, gain access to the lymph or blood-vessels of their host, and are carried, by way of these channels, to the lungs. Here they leave the blood and pass out on the mucous membrane, crawl up the windpipe to the throat, from there down to the stomach, and then to the small intestine, where they take up their abode, reach adult life and a length of one third or one half an inch, and in turn produce eggs to infect more polluted soil. In addition to this, they produce a very serious anæmia or thinning of the blood, which

gives rise to weakness, shortness of breath, dropsy, mental dullness, and, in children, checked development and stunted physique and mentality. Such conditions impair the body's resisting powers, and the victims more readily succumb to malaria, dysentery, and other diseases.

The short outline of the life-history of the worm just given indicates the wisdom of the following measures to prevent the spread of the infestation: first, treatment of all infested; second, proper disposal of feces to prevent soil-pollution; third, the wearing of shoes in infected regions; fourth, the use of pure drinking water.

Trichiniasis is a disease produced by infestation with a kind of roundworms, *trichinæ*, that presents still another life-history and method of infestation. The trichinæ are small worms about a sixteenth of an inch in length that are natural parasites of the rat, but may also infest man and many other animals, including the pig and dog. The danger to man lies principally in the fact that the pig is a rather common host, and it is from that source that he derives his infection. The history is as follows: Rats become infested by eating their infested kindred, and they in turn may be eaten by pigs. The larval worms are encysted in the muscles, as will be described later, and are set free in the pig's stomach, when the muscle and cyst are digested. They then pass to the bowel, where they take up their abode, mature, and bore into or through the intestinal wall to the lymph spaces, where they deposit their numerous progeny, averaging about fifteen hundred in number. These are carried along by lymph or blood-streams, or possibly by their own activity, to various parts of

the body, finally penetrating the muscle fibres and setting up an irritation that results in the formation of little capsules about them. Here they live quiescent, but capable of development, for an indefinite time, occasionally for years. Too often, however, this rest is shortened by the death of the pig and its later consumption by man. If such consumption is preceded by thorough cooking no harm will result, as the heat will destroy the worms; but if, as is sometimes the case, the meat be eaten raw or underdone, the larvæ are set free in the man's stomach as the others were in the pig's, and he in turn develops adults, has larvæ scattered through his muscular system, and suffers from pain, fever, symptoms resembling typhoid, and, possibly, death.

Filaria, a roundworm whose larvæ circulate in the blood, presents a still different history and method of entrance to the body. For at least two varieties, whose development has been pretty well traced, the history is as follows: The young or embryonic forms are imbibed with blood by mosquitoes that bite infected persons. In the blood these embryos, which are about one eightieth of an inch in length, are inclosed in loose individual sheaths or capsules, within which they wriggle about very actively. After reaching the mosquito's stomach the blood containing them clots and they break through their sheaths, and, after about a day, through the stomach-wall. They live in the body of the mosquito, growing and maturing, for two weeks or more, at the end of which time they have increased to four or more times their original length, and have gathered in or about the fleshy part of the mosquito's proboscis. When the mosquito next bites a man they break through a

thin part of the proboscis and pass through his skin by the small wound from which the mosquito is extracting blood. Once in the body, they wander in ways and parts unknown for an indefinite time, until they reach maturity, and, if both sexes be represented, begin to turn out embryos for other mosquitoes to ingest. The adults are very apt to take up their abode in lymph-vessels, and by their residence there they set up irritation that leads to blocking or obliteration of the channels and is thought to be the cause of lymph-scrotum, chyluria (milky urine), elephantiasis, and kindred troubles. Certain *insect larvæ* that bear a resemblance to roundworms occasionally infest man. Blow-flies may deposit their eggs on meat that is left carelessly exposed, and, if they are not destroyed by heat, maggots may hatch from them and appear in the stools of persons eating it. Flies also deposit eggs, and maggots develop, on neglected sores, wounds, or ulcers, which may later be burrowed in many directions.

Screw-worms are the maggots of a small fly found in some of our Southwestern States and elsewhere. This fly deposits its eggs in the nostrils, or, occasionally, on other parts of sleeping persons, especially those afflicted with nasal disease or very foul breath. The maggots hatch out in a few hours and crawl and bore their way in all directions, causing great suffering and, in many cases, death. Among twenty-three cases reported from Arizona in the fly season of 1905, there were four deaths.

Some flies deposit their eggs on or in the skin of animals, and the larvæ develop there, constituting "warbles" in cattle, deer, and wild rabbits. Occasionally man is host, though this is rarely so.

Some insects burrow into or beneath the skin in their adult condition, and there nourish themselves and sometimes deposit their eggs or their young. They cause irritation or disease. Among them may be mentioned the *itch-mite*, *jiggers* or *chiggers*, and many kinds of *ticks*. *Itch* was at one time a very widespread, common, and intractable disease, but is no longer so.

Lice of various kinds are also less common than formerly, but are still seen often enough as human parasites.

Leeches are at times a great pest and cause much loss of blood, particularly in some parts of the Philippines, where they get on troops marching through the jungle, in great numbers. They may get on the body from water or from vegetation. Clothing is the main source of protection from them.

A number of very important and well-known diseases are caused by animal parasites of such extremely small size and so low in the scale of animal life that they may, for our purposes, be regarded as closely allied to the bacteria. In fact, even scientists are not agreed as to whether some of them, for instance the germs of syphilis, yaws, and relapsing fever, are animal or vegetable. A brief consideration of some of the small animal parasites and the diseases they cause may be helpful.

**Minute
Animal
Parasites** *Malaria* is the best known and most common of such diseases, and the parasites causing it are better known than most of the others occurring in man. There are at least three varieties of malaria, each caused by its peculiar type of organism, though these types have much in common. They all live in the blood and

in its red corpuscles, when so small as to be almost invisible with the microscope, and there grow to a considerable size, destroying the blood-cell and converting it into a mere shell and some pigment granules. In twenty-four, forty-eight, or seventy-two hours, depending on the variety of organism, they segment or break up into a number of small new organisms, the number varying from six or eight to fifteen or twenty, and also depending on the type. These young forms are set free in the blood-stream with the breaking up of the shell of the blood-corpuscle in which they were contained, and in turn attach themselves to new red cells, to repeat the performance. Corresponding to the liberation of the swarms of young comes the "ague fit," or the malarial chill, fever, and sweat, and it depends upon the type of organism whether this comes daily, once in two days, or once in three. Various combinations of infections may make the chills appear to come very irregularly, but such irregularities are often explainable if the case be well studied. After a time the body develops powers of resistance that enable it to overcome the effects of the organism, and spontaneous recovery may occur. Or the recovery may be apparent only, and the disease may recur when chill, exposure, or other illness lowers vitality. Malaria causes many deaths, especially in the tropics; but if a single infection is not fatal very soon, the tendency is to recovery, as the parasites cannot maintain their powers indefinitely unless they renew their strength (in some unknown way) by *sexual reproduction*, and that can only occur in *mosquitoes* of certain kinds. Minute differences can be detected among the parasites of any kind of infection, and these are now known to mark sexual differentiation. If the right

kind of mosquito bites the infected individual, these sexual differences become more marked in its stomach, and sexual conjugation there takes place. The impregnated female (this term is not accurate but it conveys the idea) bores into the stomach-wall and undergoes a development which eventually results in the production of hundreds or thousands of young forms that finally find their way to the salivary glands and mouth of the mosquito, and enter, by way of the wound made by the insect, the next man it bites, and the cycle begins again in him. All malarial infections are conveyed in this way, and where there are not the right kinds of mosquitoes the disease cannot originate. So far as known, man is the only animal, and mosquitoes of the sub-family *anophalinx* the only insects, that respectively harbor and transmit malaria, and each derives its infection from the other. Each should therefore be kept away from the other.

Amœbic dysentery, so well known to our army because of the great amount of invalidism it has produced among our soldiers on tropical service, is also due to an animal parasite of microscopic size. This is a minute round mass of living substance endowed with the capacity for motion, by virtue of which it inserts itself between the body-cells or wraps itself about them and digests them; destroying tissue and giving off poison, it causes the formation of ulcers in the bowel, inflammation of the intestinal surface, bleeding, wasting, diarrhœa, and, too often, death. Entering the blood or lymph-streams through the ulcers they make in the bowel, the *amœbas* may reach the liver and there set up the dreaded disease known as liver abscess. The *amœbas* enter the body in polluted water, and probably also on

lettuce, radishes, and other vegetables that are eaten uncooked.

Sleeping sickness, a disease unknown in our possessions but caused by a parasite closely related to that causing the well-known and fatal *surra* of horses in the Philippines, is widespread in central Africa, and threatens to depopulate immense tracts there. The organism and the disease are transmitted by a biting fly.

Syphilis, known everywhere, is caused by an extremely fine organism of spiral figure. This is about 1-2000 of an inch in length, and the thread that is twisted into the spiral shape, the organism itself, is about 1-100,000 of an inch thick. This extreme fineness, and the difficulty with which the germ stains, account for the fact that it was only recently discovered. This organism, which is called a *Treponema*, is not known to occur elsewhere than in syphilitic subjects, but it occurs abundantly in lesions of the disease and is free on the surface of such moist sores as the open chancre and patches in the mouth. If from these it passes to wounds, cracks, or other open and moist surfaces on a healthy person, as may occur in kissing, sexual congress, or other contact, the disease is transmitted.

Yaws, a skin disease that occurs in the tropics, is caused by an organism indistinguishable from that of syphilis.

Many other diseases are produced by animal parasites, but it is unnecessary to consider them here.

Certain diseases, such as *yellow fever* and *dengue*, are produced by unknown organisms that pass through fine filters and are probably so small as to be invisible to our microscopes ; yet the course and history of the

diseases point to an animal rather than a vegetable cause.

The disease-producing vegetable parasites are of three classes, *moulds*, *yeasts*, and *bacteria*, all microscopic in size and all of them germs. The moulds and yeasts are of minor consequence as compared with the bacteria, and cause relatively few diseases, and those principally of the skin. Fatal general infections by germs from both of these classes have, however, been reported. Persons who have served in the tropics recall how difficult it is there to keep shoes, clothing, trunks, and many other articles from moulding or mildewing, especially during the rainy season. Heat and moisture afford such excellent conditions for growth that a pair of shoes will turn green in a very short time. As related vegetable forms are responsible for many of the most common skin-diseases of the tropics, the fact helps us to understand the prevalence of the latter. Not all moulds are harmful, and most of them are mere surface growths. The mistake is occasionally made of rejecting a quite good ham, for instance, because it is mouldy on the outside. Such mould can be trimmed off with the rind, and does not injure the meat. Roquefort and some other cheeses are dependent on moulds for their flavor, and are not good if these be absent.

Bacteria, however, are the best known of the causes of disease, so well known as such, in fact, that two erroneous ideas have arisen in consequence: **Bacteria** one, that all bacteria produce disease, the other that all disease-producing germs or micro-organisms are bacteria, or, to touch a still greater error, bacilli. Bacteria are microscopic, one-celled organisms

of vegetable nature that multiply by simple fission. So far are they from being universally harmful, that life would soon disappear from the face of the earth if all bacteria were killed. This is because of the fundamental difference that exists between bacteria and the larger forms of vegetation. All animal life is maintained directly or indirectly by vegetable life, even the lion and tiger being dependent on the herbivorous deer and similar animals. Most vegetable life draws its substance from the elements and simple compounds of the earth, air, and water. In other words, grass or a tree cannot eat dead grass, meat, or other highly organized substances, until they are first decomposed and broken up into the elements and simple compounds indicated, and returned in those forms to the earth, air, or water. For this purpose bacteria are necessary, and without their action the earth would soon be covered with dead animals and vegetables, the fertility of the soil would be exhausted, and life would be choked out by the accumulation of its products. Familiar and concrete examples of their value are furnished by agriculture, soil that will not produce peas, for instance, being made to raise them of excellent quality by the mere addition of certain bacteria. Their importance being so great, it is evident that their distribution is practically universal. Many trades are partly or entirely dependent on bacteria, yeasts, and moulds. Among them may be mentioned brewing, and wine, vinegar, cheese, and butter-making.

Bacteria are variously classified according to shape, action, and other qualities. Thus some are called putrefactive, others nutrifying or denutrifying, and others pathogenic or disease-producing. The more common

forms are designated as *cocci*, or little balls, *bacilli*, or little rods, and *spirilli* or *vibrios*, or little spirals or pieces of spirals. Cocci in pairs are called *diplococci*; bound together in chains, *streptococci*; gathered into bunches like grapes, *staphylococci*; and, as each organism tends to grow always in the same way, such names become permanently attached as more or less specific. Thus we speak of the streptococcus of erysipelas, the diplococcus of pneumonia, and the yellow staphylococcus of suppuration.

Bacteria may also be *ærobic* or *anærobic*; the former unable to live without oxygen, the latter with it. *Tetanus* or *lockjaw* is caused by an anærobic bacillus, and that probably partly accounts for the fact that it is most apt to follow small deep wounds, such as nail-punctures, to which the air does not have free access.

Reproduction among bacteria is by fission or simple division. One bacillus divides crosswise into two, the two into four, and so on; and so rapidly does the process advance, that under favorable circumstances the increase amounts to uncountable millions or billions in a day. The rapidly multiplying individuals may be bound together in gelatinous masses that are of considerable size, at times forming a layer of skin many yards in extent. When conditions are less favorable, as when all the food-supply is exhausted, the temperature too high, or harmful chemical substances present, the bacteria usually die; but some varieties may preserve themselves under such circumstances by the formation of *spores*. These may be compared to the seeds of larger plants in that they are much more resistant to heat, cold, and other harmful influences, and, after surviving such trials, may again give rise to growth. An-

thrax bacilli, for instance, are readily destroyed by heat, but their spores will withstand boiling for some time. In preparing media for the cultivation of bacteria it is therefore the practice to sterilize them at a temperature higher than that of boiling water, or to use that temperature on three successive days. In the latter event the bacteria develop from the spores after the first sterilization and are killed in their less resistant forms. The amount of heat required to destroy bacteria, what is called the thermal death-point, varies greatly with the species. Some organisms found in water, none of which, fortunately, produce disease, flourish in a high temperature. All disease-producing organisms, except those in spore form, are promptly destroyed by boiling, and many of them by a lower temperature. Few ordinary organisms will survive a temperature of 57° C. for more than a short time, and this fact gives the process of *pasteurization*, as applied to milk, its value. This is the process whereby milk is heated to a point between 65° and 70° C. and maintained at that temperature for ten or fifteen minutes. This kills the disease-producing organisms, without changing the taste and character of the fluid as does boiling. For the same reason, satisfactory results have been obtained in the British service with a camp water-heater that raises the supply to 85° C. for only a very short time.

Moisture is necessary for the growth of bacteria, and drying checks their growth even when it does not kill them, as it does many varieties. Being solid bodies heavier than air, bacteria are not given off from moist surfaces. Occasionally they may be thrown into the air with droplets of water by the bursting of bubbles, but in general a tubfull of bacteria-laden water would not

endanger the purity of the atmosphere. On the other hand, dried or partially dried bacteria may be blown about as dust and do much harm. Cuspidors and water-closets should therefore always contain water. *Sunlight* is a great enemy to most pathogenic bacteria, some of them, for example the cholera germ, being killed by a very brief exposure to it, and all being injured by it. Many chemicals, such as bichloride of mercury and carbolic acid, kill bacteria *when brought into contact with them*. Such mistakes are often made as that of regarding a stool disinfected because some antiseptic solution has been poured over it. As a matter of fact, the antiseptic in such a case is only brought in contact with the surface of the fecal mass and the great bulk of it is quite as dangerous as before.

One variety of bacteria may tend to destroy another. Thus typhoid bacilli will keep alive much longer in sterilized than in contaminated water, the common water-organisms in the last case crowding them out; and fecal matter and its contained bacteria are very quickly made to disappear if mixed with dry earth containing an abundance of nutrifying bacteria, while, if mixed with ashes, which are sterile because burned, they persist much longer. This warfare of bacteria has had a recent exploitation because of the interest attaching to the recommendation that certain milk-souring bacteria be ingested for the purpose of crowding out from the intestines more harmful varieties, and so lessening the danger of poisoning by the products of these. This explains the occasional excellent effects of the use of buttermilk as a diet.

On the other hand, one variety of bacteria may increase the danger or power of another variety. Thus

the tetanus bacillus, being anærobic, cannot grow in the presence of oxygen; but if introduced into the body with a variety of organism that uses a great deal of oxygen, the latter may soon bring about practically anærobic conditions in which the former can flourish.

The virulence of the streptococcus of erysipelas is greatly increased if the organism be grown with *bacillus prodigiosus*, an organism that is itself not pathogenic. Smallpox, though more than a single suppuration, is always associated with that process and with the common pus-producing bacteria. Bacteria in general produce disease by means of the poisons or *toxins* that they elaborate or set free, rather than by their physical presence, though in many instances, such as plague, cholera, typhoid, and tuberculosis, the toxin is only set free when the organisms die and disintegrate, so that their actual presence in the body is necessary for the production of symptoms. Other bacteria, notably the germs of tetanus and diphtheria, produce soluble toxins which can be obtained in germ-free solution, and which, when injected, produce the symptoms of disease, even though the actual germs are not introduced. In nature, however, disease is rarely or never produced in that way, the germs always being present, though it may be only on a surface, such as the lining of the mouth or bowel, the toxin being the only part actually getting into the blood or tissues. It is an interesting fact that living and virulent cholera organisms may be injected beneath the skin by the billions, with no other result than local irritation and a resulting increased resistance to the disease, although the same organisms in the bowel, but actually *outside* of the body-tissues, might cause death.

CHAPTER X

DISEASE-CARRIERS

It has been stated in preceding chapters that diseases are oftentimes diffused by persons, animals, insects, or inanimate substances that act as carriers of the infective matter or germs. It will greatly aid in the maintenance of the health of troops if company officers and the men themselves have a proper realization of such methods of spreading disease, and the better known and more important groups of carriers will therefore be considered.

Man himself is the most important carrier of his own diseases, and most epidemics arise from the presence of one case of disease in man. Isolation and quarantine are designed to limit or prevent disease-transmission in this way, and in the case of measles, scarlet fever, smallpox, and similar diseases the danger is recognized by all. It is not so generally recognized in many other common and serious diseases, such as typhoid and "colds." It is not the purpose, however to discuss here the danger arising from contact with well-marked cases of disease, but rather to consider the unknown or unrecognized carrier. Several diseases may be widely disseminated by persons entirely ignorant of the fact that they are sources of danger. This fact has long been recognized in regard to some infections, is just receiving recognition in regard to some others, and is possibly not recognized at all in regard to still others. For several years it has been a

Human Carriers

matter of scientific demonstration that active and virulent *diphtheria* bacilli may linger for weeks or months, in the mouths, throats, or noses of persons who have recovered from the disease and who appear quite healthy. It has also been demonstrated beyond dispute that some mild sore throats, "colds," and nasal discharges are due to diphtheria germs, even though the subjects do not feel particularly sick and the diseased parts do not present the appearance formerly considered characteristic of diphtheria. Other persons who have no sore throat or other evidence of disease, but who have been in contact with cases, as nurses or otherwise, may harbor the germs. Persons in any of these classes may ignorantly and innocently introduce the disease in new localities, transmit the germs to susceptible people, and so start epidemics. One such person in a crowded and ill-ventilated squad-room in winter may endanger the whole command, and if, in addition he is a person addicted to careless spitting, to the use of other men's pipes, cups, or linen, the danger is greatly increased. The wise and necessary precaution in military life is to examine and reëxamine all men who have had or been exposed to diphtheria, and to keep them isolated until it has been repeatedly demonstrated that they are free from germs, and to isolate cases of sore throat as though they were all diphtheric.

The germs of *pneumonia* and of *cerebro-spinal meningitis* are also occasionally found in healthy mouths and noses, and it is quite possible that those diseases may occasionally be transmitted in a similar manner. *Scarlet fever* often leaves an irritated throat, a nasal discharge, or a running ear, and persons who have otherwise recovered from it may transmit the infection to

others and thus start epidemics. It is only recently that a somewhat similar origin for epidemics of *typhoid* has been recognized; but so many instances are now recorded as to leave no doubt of the possibility, and as to make it seem probable that the disease is often spread in that way. Persons who have recovered from the disease months or years before, and have since seemed in excellent health, have been shown to be passing almost pure cultures of the typhoid bacilli in their stools, and to have caused many epidemics. In no instance did they know that they were carrying the germs or were sources of infection. These carriers will be discussed again in the chapter in Part III dealing with typhoid fever.

The germs of *cholera* and *dysentery* may be passed by persons quite ignorant of the fact that they even harbor them, and may so give rise to epidemics through water-infection, by the use of latrines from which flies may obtain the germs, and in other ways. As stated in the preceding chapter, man is the carrier of the commonest *intestinal worms*, and it is his ignorance or carelessness that is responsible for their spread. Persons who spread *tuberculosis*, *tonsillitis*, and other troubles by their careless spitting usually do it in ignorance of the fact that they are doing wrong. Most consumptives do not realize that they have the disease until after they have been expectorating tubercle bacilli for some time. Other disease-carriers who may be quite ignorant of the rôle they are playing are sufferers from certain eye and skin diseases, such as *trachoma* or granular lids, *gonorrhœal inflammation of the eyes*, *ringworm*, *itch*, *impetigo*, and many other diseases of the skin. They may either be ignorant that they have any disease, or that the diseases they have are contagious, and so, by

careless contact, the use of public towels or brushes, or those belonging to other persons, through barbers' chairs and in other ways, they may infect many persons.

Nurses and doctors may at times carry and transmit infectious diseases, such as *measles*, *smallpox* or *typhoid*, either innocently, as when a diagnosis cannot yet be made and they do not know of the dangerous contact, or culpably, as when they know of the danger but do not take proper precautions to guard against it.

Venereal diseases, both gonorrhœa and syphilis, may be innocently transmitted, quite aside from any question of sexual intercourse. A syphilitic infant may infect its nurse; a girl may get a chancre of the lip from kissing a person who has infective lesions in the mouth and does not know it; or possibly, though it must happen very rarely, by following a syphilitic in the use of a communion cup or other public drinking-vessel; doctors occasionally become infected with syphilis through needle-pricks or slight scratches that become infected during operations. A child may contract gonorrhœa from sleeping with or against its infected mother, or an infant's eyes may be infected during its birth, and others may become infected through handling it. Many men who thought themselves quite free from all venereal disease have infected their innocent wives through sexual intercourse, and every man who has ever had venereal disease should seek the examination and opinion of a competent physician before entering the married state.

The company officer cannot be expected to recognize the various disease-carriers in his company, but he can aid in their detection if he sees that all sick men report promptly to the surgeon, and that they remain on sick report until no longer sources of danger. Occasionally

a company officer takes the unreasonable view that a man who is able to do a bit of window-washing or other light work about the hospital should be sent to his company. Such action might result most seriously both to the company and the man, and the wise company commander will very rarely, if ever, clamor for a sick man. Even diseases not directly transmissible, such as malaria, constitute a menace in barracks, and the victims are, ordinarily, not only better off, but also less dangerous when in hospital. A man infested with worms may be capable of working and may desire to do duty, but he cannot be so well treated elsewhere as in hospital, nor will the disposal of his stools be so well looked after if he be free to go and to defecate where he pleases.

Certain diseases are derived exclusively or almost so from animals. Among them are many very dangerous infections, some of them quite incurable.

**Animal
Carriers**

We will consider some of these.

Hydrophobia is a disease that affects a great many kinds of animals and is almost uniformly fatal. It is ordinarily considered a disease of the dog family, but there is some reason to believe that it is naturally a disease of certain herbivora, especially rabbits, and the dog contracts it from these. So far as man is concerned, however, it may be considered as originating with the biting animals, especially the *dog*, *wolf*, *fox*, *cat*, and *skunk*. By these it may be transmitted to horses, cattle, and other domestic animals, and to man himself. No cure is known for the disease after it has actually begun, but if treatment be begun soon after infection, immunity may be established before the incubation period has ended, and the onset thus prevented. As there are three prevalent errors

regarding this disease, each fraught with the possibility of serious consequences, they will be briefly discussed here.

The first of them is the somewhat widespread belief that the disease does not exist, that its victims really die from fright. This is abundantly disproved by the deaths of people who were not frightened, who for weeks or months had forgotten the incident of the bite; by the ability to infect and kill rabbits, dogs, and other animals, by inoculation from animals dead of rabies but not from those dying of other diseases; by the infection and death of cattle, horses, and men; by the bites of rabid animals, but of no others; and in other ways. *Hydrophobia exists.*

The second error relates to the diagnosis of the disease in the dog. Not every dog that slobbers and bites is rabid, and it is senseless, as well as cruel, to start a howling mob after it and pound it to death on suspicion. The disease usually begins in the dog with a change of disposition, usually shown by restlessness and irritability; the animal is easily startled, or crouches in fear. The appetite may fail or there may be difficulty in chewing or swallowing food. Drinking is less interfered with, and the name hydrophobia (fear of water) is not accurately descriptive. Change in the bark or absence of barking is often noted. Later, the animal may show a tendency to tear and swallow foreign bodies, and to snap at things in its vicinity or at phantoms of its disordered brain. This may be followed by a furious stage, when the dog "runs mad." Always the disease ends in paralysis, usually first showing in the muscles concerned in swallowing, later of the hind-quarters, the jaw, and the entire body, death coming in from four to seven days

after the onset. The disease probably begins with the invasion of the dog's brain by the parasites, and the saliva is not always dangerous at the beginning. The diagnosis can be properly made only by observation of the whole course of the disease, or by examination of the animal's brain. Therefore, the correct course of procedure is either to *capture the dog and confine it* so that the disease may be observed, or to *kill it and at once cut off its head and send that to a competent pathologist for examination.*

The third error relates to treatment of bitten persons. It is computed that about one sixth of the persons bitten by rabid animals develop hydrophobia, the rest escaping because the saliva was not at the time infectious, because the clothing wiped all saliva from the biting teeth, and for other reasons. Nevertheless, the sixth that do develop the disease always die, and proper preventive treatment is essential. This consists in the cleansing of the wound with heat or antiseptics, the promotion of bleeding, proper dressing, and the Pasteur treatment for the establishment of immunity. "Mad-stones" and charms are quite useless.

Glanders is well known as a disease of *horses* and *mules*, less so as a cause of human illness. Nevertheless, human beings do contract it, usually with fatal results. Its contagious and fatal characters are so well recognized in the army, that its presence constitutes proper grounds for the destruction of animals, even though their usefulness is not yet greatly impaired. The persons who most often contract it are those concerned in the care and treatment of horses. As the disease in its early stages may not present characteristic or well-marked symptoms, it may be mistaken for a "cold,"

quinsy, or other less dangerous malady, and the animal be handled or treated without the observation of proper precautions to prevent infection. Special care should always be exercised to prevent any discharge from the nose or from sores on horses coming in contact with wounds, scratches, or the respiratory tract, and to keep the animal from blowing its nasal discharge or saliva into the face of one examining it. All suspected animals should be isolated and handled as though known to be glandered until the diagnosis is settled. The disease may also be contracted from handling dead animals during their removal, or in the course of post-mortem examination, and proper precautions as to cleanliness, avoidance of wounds, and disinfection should be observed. *Glanders* and *farcy* are different forms of the same disease, are due to the same organism, and must be treated with equal respect.

Plague or *the pest* is one of the most serious and important diseases for which man owes a debt of hatred and warfare to his animal neighbors. It has not as yet gained an extensive footing in our own country, but it has maintained a slight hold in San Francisco for several years, has appeared in a few other Pacific coast cities, and has worked some havoc among such near neighbors as Peru, Venezuela, the Argentine, Brazil, Mexico, and the Azores; so that we may see much of it in a few years if the greatest and most unremitting care is not exercised. It has prevailed extensively in the Philippines, though it now seems under good control there, while in India, China, and other oriental countries it has caused many millions of deaths. Its prevalence and extension in India are such at the present time as to be causing the abandonment of large tracts of country and of whole

villages, and, if reports be true, to threaten the very existence of the government. The investigations of the British commission appointed to study the disease in India show clearly that epidemics of human plague arise after and in consequence of epizootics of *rat plague*. The infection is transmitted from *rats* to man by *fleas*, usually those of the rat, and the relation of the epidemics to the epizootic is beautifully shown as follows: There are two principal species of rat concerned there, the brown or Norway rat and the black or house rat, the former also known as the ship and the sewer rat. It is among that class that the epizootic arises, usually, and in about 7.2 days later it is also prevailing among the house rats. This 7.2 days is accounted for by the fact that it is also the average experimental time elapsing between the biting of the black rat by an infected flea and the death of the animal. (The rise of the epizootic is measured by the numbers of rats found dead or dying.) The black rat epizootic precedes the height of the human epidemic by from ten to fourteen days, a period accounted for as follows: The rat flea does not readily attack man until starved for three days. The incubation period of human plague is about three days. The duration of the fatal human plague averages four and a half days. Average total, eleven and a half days.

The relationship between rats and human plague is also proved in other ways, but they cannot be discussed here. The fact is established, and it constitutes the most important of the many reasons why man should be at enmity with, and try to exterminate, rats.

It must be borne in mind, however, that other animals may be infected, and recently wild squirrels have been shown to have the disease in California. They

and gophers, prairie dogs, and similar animals may yet play an important part in disseminating it. While dirt and unsanitary conditions cannot give rise to plague, their influence in promoting the presence and increase of both fleas and rats is so well recognized that it is well to mention the matter as showing how they may strongly predispose to disease that they cannot cause directly.

Tuberculosis is the most prevalent disease affecting man, and is responsible for more deaths than any other one cause. It is also a common disease of cattle and of some other animals, and it may be transmitted to man from them. Tuberculous meat, if not sufficiently cooked to sterilize it, may be the medium of infection, but milk is probably even more commonly the carrying agent. A cow that has tuberculous disease of the udder may give off immense numbers of bacilli, and these may be ingested in a virulent condition in milk, cream, butter, or cheese. This method of infection is so common that some authorities who have devoted great study to the subject contend that it is the usual, if not the exclusive method, and they relegate infection by inhalation to a place of very minor importance. Such a view is extreme and probably not correct, as shown, for instance, by the fact that tuberculosis is very prevalent in Japan and other far eastern countries where neither milk nor beef is a common article of food. Nevertheless, infection by means of these food-articles can and does occur, and the precautions against the sale of tuberculous meat and milk are wise and salutary. Wild or range cattle are much less subject to the disease than high-bred, stabled cattle, and many other animals that suffer from it in captivity are free from it in their native wild state, thus

showing the predisposing influence of confinement and bad air.

Tuberculosis is rather common also in pigs, dogs, and cats, less so in sheep and goats. The disease may be transmitted by the flesh of pigs. Pet animals, such as dogs and cats, are apt to contract and spread the infection because of their habits of licking up sputum and other discharges to which they may have access. This furnishes another argument for the proper care and destruction of all tuberculous discharges. The urine and fecal discharges of men or animals that have tuberculosis may contain the bacilli and may constitute serious dangers. Thus a cow that is passing the germs in these ways may indirectly infect sound cattle or their milk by means of bacilli blown about the stable with the dust of dried feces or urine.

Fish and birds also have tuberculosis, but it is doubtful if the forms of it that they have can be transferred to man. It is not, however, necessary to go to the extreme lengths that fear may carry one in the efforts to avoid tuberculosis. Patients with the disease need to be taught to take proper precautions as to the disposal of their spit and other excreta, but when they take such precautions they need not be avoided. Tuberculous milk is best avoided, but if the milk is not known to be infected, but is merely of unknown or doubtful origin, it can be rendered safe by boiling or pasteurization. The indiscriminate sale of tuberculous meat is properly forbidden by law; but if an animal is not badly diseased and the visibly injured portions of the carcass are rejected, the rest can be eaten with safety if it is first properly cooked. Such meat is not purchased for the army, however, and the above statement is made for its

general worth and not to encourage the acceptance of inferior stores.

Anthrax is a disease that causes a very great number of deaths in cattle and sheep in various parts of the world. Other animals are also susceptible to it in varying degrees, among them being horses, deer, pigs, mice, and rabbits. Man suffers from it in two or three forms, generally obtaining his infection from the hides of animals dying of it. Wool-sorters and hide-handlers are therefore particularly subject to it, and one common name of the trouble is "wool-sorters' disease." Soldiers have also been known to be infected (though not in our army) through the use of sheepskin-lined coats, or boots made of poorly cured leather. The disease is often fatal, and materials from animals dead of it should not be used. It is caused by one of the spore-forming bacilli, and the spores are very resistant to both heat and chemicals. Human infection takes place in two ways: through cuts or abrasions brought into contact with infected skins, carcasses, or other matters, and by inhalation of hairs, bits of wool, or dust that carry the spores.

Man is the occasional subject of some other diseases of animals, and it is a good general rule to take as much care to prevent infection from a sick animal as from a sick person, unless the disease is known to be one, such as rinderpest, to which persons are not liable.

Domestic animals are also occasional carriers of common human diseases. Dogs and cats have been known to have diphtheria. Cats have been experimentally infected with whooping-cough, and there is reason to suspect that either of these animals might carry the contagium of such diseases as smallpox, scarlet fever, or measles in its hair. The liability of getting ring-

worms and other skin diseases, as well as lice and fleas, from domestic animals has already been mentioned.

Insects of different kinds transmit diseases in various ways. The first and simplest way is by acting as *simple*

Insect *carriers*. Thus germs become attached to
Carriers the legs and bodies of flies alighting on
an infected wound or a yaw's papule, and

these by later alighting on clean wounds or abrasions may leave the germs there and cause infection. Or the transfer may be less direct, as in the case of the fly that goes from a typhoid stool to a food-article and there leaves germs to be later ingested with the food. In other instances the simple deposit of the germs may not insure infection, but the irritation caused by the insects may further it. This is well shown in the case of plague. The flea transmitting that disease may do so by reason of the fact that the germs are adhering to its mouth-parts and are introduced directly by the biting act. On the other hand, the mouth and external parts of the flea may be free from germs, in which case infection may occur as follows: The flea has a habit of defecating while it sucks blood, probably to make room for a larger meal, and in so doing it may deposit plague-germs on the skin, near but not in the wound. If, now, the bite is rubbed or scratched, as is often the case, the germs may in that way get into the wound and infection be assured. More complicated processes are involved in other instances, as in malaria, yellow fever, or filariasis, in which the parasites must undergo a cycle of development in the insect, and the latter cannot transmit the disease until a sufficient time has elapsed to allow that to occur. Some insects can transmit germs, and the capacity to convey infection to man or other animals,

to their offspring. This fact has long been known regarding the tick that transmits Texas fever to cattle, and it has recently been proven true as to those infecting man with the spotted or tick fever of Montana and the relapsing or tick fever of Africa. These diseases may be transmitted by the bites of ticks raised from the eggs of those biting infected persons, but which have not themselves ever bitten people or had any other chance than through heredity to obtain the infection. It is regarded as possible that malaria-bearing mosquitoes may possess the same characteristic, and it is probably true of some other insects. Among proved or probable insect-carriers of general diseases may be mentioned *flies* of various kinds, several species of *mosquitoes*, *fleas*, *lice*, *bedbugs*, *ticks*, *mites*, and *roaches*.

Among inanimate carriers of disease, *food*, *water*, and *milk* have already been mentioned several times. They may serve as the conveying media for practically all infections that can occur through the alimentary canal, including such important ones as tuberculosis, typhoid, cholera, dysentery, diphtheria, scarlet fever, and many others. These articles, and milk especially, offer such opportunities for the multiplication of bacteria that they may convey to the consumer many thousand times as many germs as originally had entrance to them. Thus a can of milk, becoming infected with a few hundred typhoid bacilli from the hands of a farmer-carrier, may, when consumed in a city, ten or twelve hours later, have billions of germs in it, and cause a great many cases of the disease. It is, therefore, important to prevent the access of even *one* disease-germ to such articles,

**Inanimate
Carriers**

and carelessness in their handling should not be condoned. Fortunately, thorough cooking will destroy disease-germs in all of these substances, a fact borne out by the relative immunity to cholera shown by the Chinese in the Philippines. They suffer in much less proportion from that disease than do Filipinos, Europeans, or even Americans, a fact accounted for by the Chief of the Bureau of Health by the statement that they eat and drink practically nothing uncooked. In nearly all other respects their sanitary environment is bad.

Clothing, books, carpets, and other personal or household goods that have been used or handled by persons ill with such diseases as smallpox, measles, and scarlet fever may convey infection, and for that reason the disinfection or destruction of such articles is resorted to as a means of preventing the spread of contagion. Such articles as can be boiled or thoroughly steamed can be quite well disinfected; but if facilities are not at hand for sterilizing such large articles as mattresses, they may need to be destroyed. Boots, hats, and certain other classes of articles will not stand steaming, and will have to be disinfected with chemicals or destroyed.

Such articles of personal belongings are also common carriers of the organisms of suppuration. So widespread are such germs that almost everything in common use has them on it; and if brought into contact with a wound they will leave the article and start infection. This alone makes the sterilized and sealed first-aid packet of so much greater value than articles of clothing and handkerchiefs as first dressings for wounds. The packets, if not contaminated by handling, practi-

cally never cause infection ; the other articles almost always do so. Nails, bullets, knives, and other missiles or weapons that have not been sterilized may infect the wounds they cause ; or, if sterile themselves, may carry infection from the skin or clothing they traverse.



PART III

THE PREVENTION AND CONTROL OF EPIDEMICS

*"Thou shalt not be afraid . . . for the pestilence that walketh in
darkness : nor for the destruction that wasteth at noonday."*

PSALM 91 : 56



CHAPTER XI

THE DEFENSES AGAINST DISEASE IN GENERAL

FORTUNATELY for the human race, the contracting and development of disease are somewhat difficult matters, and only follow the coincident occurrence of certain trains of circumstances. They may be compared with the defeat of an army, for which purpose it is necessary that the army be weakened by loss of men, hard marching, lack of food, exposure, or other cause; or that the enemy has been able to bring a superior force on the field, that his supplies are better or are more promptly delivered; that one side improves its opportunities for offense or defense and the other does not; or that other good and sufficient reasons are present. The mere existence or contact of two armies does not explain that a certain one of them is defeated. So in the matter of sickness, the mere existence or contact of a man and typhoid organisms is not in itself sufficient to insure that the man has or will get typhoid fever. For that purpose it is necessary that the germs shall be alive, virulent, and capable of development, that they shall gain entrance to the body of the man in sufficient numbers, that the man's body shall be unable to destroy them before they have time to develop and liberate their poisons, and that his tissues shall unite with such poisons after they are liberated. It is relatively rare that these conditions coexist, and typhoid fever is therefore not more common than it is. Our defense against sickness then rests on our ability

to break or destroy the chain of disease-favoring circumstances. If we can entirely destroy it, we are sure to remain well, but if we cannot do that, the destruction of one link may save us. It is well, therefore, that we should first consider general plans of defense or aggression, and later pass to the discussion of special campaigns or battles.

The body is defended from disease by many natural factors, and we should try to preserve these intact. The *skin* presents an impassable barrier to most organisms, and very few infections can enter through it as long as it remains unbroken. It has weak points, however, and infections may occur in the hair-follicles or in the skin-glands. Boils are the commonest results of such invasions. The organisms of typhoid, tuberculosis, and many other diseases could be put on the sound skin with impunity, if they did not later gain entrance to the nose, mouth, wounds, or other openings.

The *mucous membranes* are all warm and moist, and, in that way, offer favorable conditions for bacterial growth; but they are all covered with *mucus* to which the germs may adhere, and with which they may be passed out of the body by spitting or otherwise. The cells lining some mucous surfaces have little, moving, hair-like projections, and by means of these are able to pass along and expel small foreign bodies getting on them.

The *gastric juice* is sufficiently acid to destroy many varieties of micro-organisms, and digestive disturbances that lessen that acidity increase the liability to infection. The fluid also has the power of neutralizing or digesting some poisonous products, but not all.

The *urine*, by its acidity and its irrigating and cleansing action, doubtless has an effect in keeping down the number of cases of venereal diseases.

The prompt *reaction to irritation* is often a disease-preventing factor. A dose of arsenic or a meal of tainted meat may produce such prompt vomiting as to cause the expulsion of all poison, and so prevent further symptoms. The inflammatory reaction following the infection of wounds is usually a conservative process that limits the infection to the locality of the wound, and prevents its diffusion through the body.

As stated before, however, a man may possess an *immunity* to certain diseases even though the germs gain entrance to his body in the numbers and of the virulence ordinarily producing such diseases. This may be hereditary or “*natural*” immunity, when the man is born without susceptibility to the disease. It is *acquired* when due to something occurring after birth. Acquired immunity is usually due to an attack of the disease, and is well recognized in such affections as smallpox, measles, scarlatina, and typhoid. It is common knowledge that one attack of any of these usually protects against a second. This is also known as *active* immunity, because the body is active in maintaining it. *Passive* immunity is due to the introduction into the body of ready-made immunizing substances, such as diphtheria antitoxin, that is contained in the blood-serum of an immunized horse. Active acquired immunity may also be induced by *vaccination*, the procedure whereby disease-producing organisms in an attenuated or weakened state are introduced into the body and set up a very mild form of disease, but one sufficient to set the body on the de-

**Immu-
nity?**

fensive and cause it to produce immunizing substances. This procedure is best known in its relation to smallpox, and as that was the disease for whose prevention it was first used, the name vaccination in the broad sense, that is, the introduction of disease-producing germs to produce immunity, is taken from that instance. It has also been used, though, to produce immunity to cholera, plague, dysentery, typhoid fever, hydrophobia, and several diseases of animals, in some instances with considerable success.

Immunity may also be relative or absolute, temporary or permanent. Man's lack of susceptibility to the pleuro-pneumonia of cattle, for instance, is both *permanent* and *absolute*. An attack of smallpox may produce an immunity that is either *permanent* or *temporary*. In the latter instance the sufferer might have a second attack after some years. A *relative* immunity to typhoid might protect against a small number of bacteria, but not against a very large number.

Immunity is due to several factors, not all of which are understood. In regard to some diseases, such as diphtheria and tetanus, it is due to the development of antitoxin, a substance that acts as a direct antidote to the poison of the disease and neutralizes its effects. In other instances, as in the case of plague, cholera, and typhoid, it is largely due to the presence or development of substances that kill or dissolve the bacteria. In many instances it is partly or entirely due to the activity of certain cells, phagocytes, that take up and destroy the germs. Absolute hereditary immunity may be due to none of these, but to lack on the part of the body cells of chemical groups that enable the poisons to enter into combination with them.

Relative immunity is maintained by those conditions that keep up the general health and maintain the resistance of the body. It is lowered by causes that depress these, such as starvation, exhaustion, injuries, shocks, worry, and grief.

General Health

It is therefore important that the man and his environment be constantly kept at their best, that abundant and proper food, pure water, good air, and right clothing be provided, that sufficient exercise to keep the respiration, circulation, digestion, and excretion active be taken; that overwork, improper food, alcoholic and venereal excesses, and the entire list of things predisposing to disease be avoided.

All of these measures tend to strengthen the man and put him in good condition and position for the combat with disease; and, as with armies, so in this instance, the maintenance and strengthening of our own forces is not less important than the weakening or hindering of the enemy; and as the possession of a large and well-trained army is, in these days, often spoken of as a "national insurance policy" and a guarantee of peace, so may the well-trained and sound body be regarded as insurance against disease. As the combat with germs is, however, one that never ceases, we must also try to injure the enemy as much as possible, giving no quarter, as we may expect none. We should try to eradicate all disease-producing organisms, to reduce their numbers where we cannot do that, and to weaken them by every means available. Many of the measures designed to accomplish these ends have been mentioned or discussed in preceding chapters or will be taken up more fully in later ones, but some of them may be considered at length here and not again described. They

may be classed in two categories : measures designed to prevent contact with the germs, and those aimed at their destruction.

Among the former, avoidance of the sick may be considered one of the most important measures. This

**Avoid-
ance of
Sick**

recommendation cannot be taken as countenancing neglect of the sick or lack of all proper care and attention, but it refers to unnecessary contact with them, that due to carelessness, curiosity, or sentimentality. As stated before, the principal agent in the dissemination of human diseases is man himself, and nearly every case of infectious disease is derived from another. Therefore, unnecessary visits to hospitals and sick-rooms should be discouraged, and in military life it should be axiomatic that sick men should be removed from squad-rooms, unless good reasons to the contrary exist in special cases. Hospitals are wisely provided for the reception of such soldiers, and if many sick are treated "in quarters," these large and expensive plants are kept from performing one of their important functions.

Even though the sick man goes into a general ward in the hospital, he is less apt to infect others there for

**Ventila-
tion**

several reasons. One of these is that air-space and ventilation are more liberal there.

In the squad-room, where but six hundred cubic feet of space are allowed each person, a sick man may seriously contaminate the air. In hospital the same amount of contagium is dilated two to four times, because the space per man is twelve to twenty-four hundred cubic feet. Naturally an exposed person breathes in only one third as many organisms.

In the more readily transmissible diseases isolation

is resorted to as a means of insuring lack of contact. It is necessary in the case of such "catching" sicknesses as smallpox, scarlet fever, and measles, and its value is being more appreciated in typhoid fever. The degree and character of the isolation vary with the method and ease of transmission of the different diseases. If proper care is taken in the disposal of excreta, bath water, and other waste material, and in the exclusion of flies and vermin, typhoid fever can be safely isolated in one room of a house or hospital. On the other hand, smallpox is so highly contagious that it needs to be treated in a building or tent placed at a considerable distance from habitation. In yellow fever, malaria, and dengue, the degree of isolation is not necessarily greater than that afforded by the screening necessary to exclude all mosquitoes from the sick.

Infectious diseases of various kinds have different *incubation periods*. These represent the time elapsing between the occurrence of infection, the entrance of the germs into the body, and the appearance of the disease. In some cases of cholera, diphtheria, and a few other diseases, the incubation period may be as short as one day or a few hours; while in hydrophobia it may extend many weeks and possibly months. Diseases also vary as to length of time during which the subjects remain sources of danger after their apparent recovery. Yellow fever, for instance, ceases to be dangerous, to others than the patient, after the third day of its existence; while a typhoid patient may continue to give off immense numbers of virulent organisms and constitute a source of danger to the community for many years.

The term quarantine, though used in several senses,

Quarantine

the oldest meaning a forty-day period of isolation for persons thought to be possible subjects of disease, is now most commonly used to indicate the isolation of persons thought to belong to the two classes just discussed, and of others who have been exposed to disease and may belong to one or the other. It is usually understood to mean the isolation of healthy persons as well as the sick. Employed rationally and under sound medical advice, it is often a valuable measure and may prevent epidemics. But it is always attended with hardship to its subjects, and when controlled by laymen actuated by panic, fear, or malice, it may become an instrument of cruelty and oppression. Only trained medical officers should direct or control systems of quarantine, and their training should be relied upon to suit the system to the circumstances. A fixed isolation period, say of one month, for yellow fever and scarlet fever, would be wrong in both instances, but in the one the patient would be unjustly treated, in the other, the public.

In addition to the matters of barrack, camp, and kitchen police already discussed, larger schemes of policing or engineering are important in the prevention of disease. These embrace such diverse measures as the construction of filtering plants and water-systems to furnish an abundance of pure water, the installation of sewerage-systems and crematories for the removal or destruction of waste, thorough policing and cleaning of streets and neighborhoods to keep down dust or reduce vermin, or the draining, filling, or oiling of marshes to do away with mosquitoes.

The best known and most commonly employed measures for killing germs are usually spoken of as disin-

fection and sterilization. The terms are not synonymous, though often used as if they were. When an article is sterile in the bacteriological sense, it is free from all germs. Disinfection may free it from all germs, but does not necessarily do so. It means the destruction of infectious organisms. A urinal or bedpan contaminated by use by a typhoid patient might be disinfected by simply rinsing in hot water, as the typhoid germs are killed in a few minutes by a temperature of 70° C. It would certainly not be sterile.

Disinfection and Sterilization

Heat is the most reliable of the means of sterilization of small inanimate objects. It may be applied moist or dry, the former effective at much lower degrees than the latter. Boiling water or streaming steam, both of which have a temperature of 100 C., will destroy all kinds of growing germs in a short time, while dry heat of 200° C. is necessary to accomplish the same purpose. As this temperature will burn fabrics and is dangerous, it is not much used. Boiling is simple and is easily applied to small articles, so that it is commonly resorted to where special appliances are not available. In hospitals steam sterilization is more often resorted to, as being more convenient and not necessitating so much handling and drying of articles. Large institutions also have steam-chests of sufficient size to receive such bulky articles as mattresses, carpets, and other household furnishings, which may thus be quickly and thoroughly disinfected. Gases and solutions of chemicals are also used for disinfecting; the former for rooms, furnishings, or clothing that would be injured by heat, and to penetrate into cracks and crevices not otherwise accessible. As many of these substances possess disagreeable odors,

or have the property of destroying smells, the lay public at times displays a tendency to confuse *deodorization* and *fumigation* with disinfection. The three may coexist, but they are not the same, and the words should not be used loosely, as such usage may lead to confusion of the processes and so do harm. Charcoal is a good deodorant, and burning tobacco or pyrethrum may furnish efficient fumigation in ridding a house of mosquitoes, but none of these is disinfectant. The principal gaseous disinfectants are chlorine, sulphur fumes, and formaldehyde. All of them are very irritant and unsuitable for use in efficient concentration in rooms or places where they will be inhaled.

Chlorine is not commonly used in its gaseous state, and when "chloride of lime" is sprinkled about it is liberated in such small amounts as to be inefficient. In watery solution it is valuable in disinfecting surfaces to which it can be applied. It is a powerful bleaching agent, and cannot be used on fabrics.

Burning sulphur gives off fumes which unite with water to form sulphurous acid, a very efficient disinfectant. The dry fumes have very little virtue of the kind, but are very useful in destroying vermin, such as flies, mosquitoes, and rats. When it is desired, however, to disinfect a room that has been occupied by a case of contagious disease, such as scarlet fever, by means of sulphurous fumes, the walls, floors, and other surfaces should first be moistened by wiping them with a damp rag, or in some other way. As sulphurous acid also bleaches or fades fabrics, it is frequently not to be used except at great expense.

Formaldehyde gas, or its forty per cent solution in water, known as formal or formalin, is now more often

used than any other gaseous disinfectant. The gas may be generated in the room to be disinfected by burning wood alcohol in special burners known as formaldehyde generators, or it may be liberated by the evaporation of formalin. The latter is a very simple procedure, and while its effectiveness cannot be guaranteed, the same may be said of other gaseous disinfection, and it is often the best available means. Many methods of formaldehyde disinfection have been used, but only the use of formalin will be described here. For disinfecting the contents of trunks and boxes, it may be sprinkled over them freely, or placed on absorbent cotton or blotting-paper and the boxes then closed tightly for twenty-four hours. For the disinfection of rooms, from four ounces to a pint of formalin should be used for each thousand feet of cubic contents. The gas is speedily liberated if six ounces of permanganate of potash be added to the pint of formalin in a large bowl, or if unslaked lime be added, or if the fluid be poured on hot bricks. A still simpler and quite efficient method, however, is to close hermetically all cracks and crevices about the room, stretch ropes or strings across it and on them hang sheets. The formalin is then poured upon the sheets to saturate them, and the room tightly closed and kept so for twenty-four hours or more, after which it should be well opened and aired, to rid it of all fumes.

None of the gaseous disinfectants can be depended upon to penetrate thoroughly into large and thick articles such as mattresses. Their action is apt to be superficial, and it is therefore important that the articles in a room to be disinfected be disposed in such a manner as to expose as much surface as possible. Bedding should be taken out of piles and hung on ropes or racks, pic-

tures removed from walls, books opened so as to expose all pages if possible, pockets turned inside out, and all boxes opened.

Formaldehyde is less harmful to fabrics and colors than the other gaseous disinfectants, and is altogether the most satisfactory for ordinary use.

Fluid and solid chemicals, the latter in solution, are much used as *germicides*, *disinfectants*, or *antiseptics*. The number of substances so used is large, some of the more important of them being the various *acids*, *permanganate of potash*, *peroxide of hydrogen*, and those to be discussed.

Bichloride of mercury, or *corrosive sublimate*, is probably the most important, and for many purposes the most valuable, antiseptic drug. It is used in watery solution of various strengths. A solution as weak as one part in one million has some value, but it is generally used in strengths varying from one part in five hundred to one in five thousand, and in such dilution is a very powerful antiseptic, killing most germs in a few minutes. It has attained popularity, and is commonly sold in tablets of such size that one added to a pint of water makes a one to one thousand solution. The main objections to bichloride are that it is a powerful and deadly poison, and the tablets in which it is sold may, by mistake, be taken for other substances; and that it corrodes and injures metals with which it may come in contact. It works more efficiently when combined with an acid, and the tablets are usually such combinations.

Carbolic acid, or *phenol*, is quite as well, if not better, known as an antiseptic. It is not so powerful as bichloride, and is used in solution of from one to five per cent strength. Occasionally it is used in what is called its

“pure” form, about ninety-five per cent strength. It is a very deadly poison, but it does not corrode metals as does bichloride, and is therefore better for disinfecting them. Both of these substances irritate and injure the hands if much used.

“*Trikresol*” is a disinfectant that is supplied and much used in our military service. It is less readily soluble than phenol, but is equally as powerful a disinfectant when used in half the strength of the solution of the latter. It also is a poison.

Creolin is a proprietary coal-tar preparation issued by the Quartermaster's Department among veterinary supplies. It makes a milky suspension in water, is less poisonous and less irritating to the skin than carbolic, and is quite as potent as a germicide.

There are numerous other valuable disinfectants on the market, but it is not necessary to discuss them, as those issued to the service answer quite as well as any. It should be stated, however, that *formalin* is also used as a liquid disinfectant, and for some purposes, such as the disinfection of urine and feces, it is probably the most valuable one we have. It should be thoroughly mixed with a stool to be disinfected, lumps of feces broken up, and the two allowed to stand in contact for a time. Formalin is also a valuable deodorizer.

For a few diseases we have remedies that may be termed *specific*, or truly curative. In most infectious diseases, however, we are only able to modify the course of the affection, to relieve symptoms and maintain the strength and vitality until nature, or the body forces, accomplishes a cure; and even the specifics cannot always be given in such a way that they get at and destroy all the organisms of disease.

**Medi-
cines**

Quinine is a specific for malaria, and actually destroys the growing and multiplying organisms in the blood. Yet men occasionally die of malaria in spite of quinine, and many hundreds of army men can testify that quinine does not always effect a prompt and permanent cure of the disease.

Mercury is a specific for syphilis. It kills the organisms causing the disease, and causes the lesions to heal; yet the men who have devoted the most attention and study to the subject agree that treatment must be continued from two to four years in order to assure a cure.

Antitoxin is a specific for diphtheria, neutralizing the poison and making it harmless. Yet, if it be not given early and the toxin has a chance to unite with the tissues, it may fail, and death result in spite of its use.

Altogether, the consideration of specific remedies cannot but convince us that the prevention of disease is of much more importance and value than its treatment, and that the tendency, often manifested, to regard the medical officer as one whose main office is the treatment of men already sick, one who should have nothing to do with the general administration of a post or an army, is a very grave mistake. Specifics are very valuable, however, in the prevention of disease. Mosquitoes are not apt to obtain infection from the blood of men taking quinine, nor are they apt to convey it to men who are so doing. The man who is taking mercury is less apt either to give or to receive syphilis than he who is not. Prophylactic injections of diphtheria antitoxin offer great protection to persons exposed to that disease, and the amount of "fourth of July tetanus" has been very greatly reduced by the use of tetanus antitoxin before the disease shows.

CHAPTER XII

TYPHOID AND OTHER WATER-BORNE DISEASES

WE now know that typhoid fever is transmitted in a great variety of ways, and there is reason to believe that cholera, dysentery, "camp diarrhœa," and parasitic infestations may also be. This entire group of diseases has, however, so long been spoken of as water-borne, and water does so often play the part of principal medium in their transmission, that it is convenient to discuss them under this head. For the military hygienist they constitute the most important group of diseases, and the army that escapes them is truly fortunate. In our own army they have given and still give much cause for anxiety. During the Civil War they caused more sickness than all other diseases, and so lately as 1898 they were responsible for the well-known and terrible conditions prevailing in the camps of concentration.

Speaking of typhoid fever alone, the Board on the Origin and Spread of Typhoid Fever in United States Military Camps during the Spanish War of 1898 said : "About one fifth of the soldiers in the national encampments in the United States in 1898 developed typhoid fever. Among 107,973 officers and men in 92 regiments, the records of which we have carefully studied, the number of cases of typhoid fever, according to our estimates, was 20,738." As no other epidemic on such a large scale has been studied as that one was ; as the lesson is drawn from our own service and the facts are more or less familiar to officers in that service to-

day ; and as the Board's report advanced conclusions of vast importance, frequent reference will be made to it in this chapter, and, to save time and space, it will be spoken of as the Reed Board, because Major Walter Reed of the Medical Corps, U. S. Army, was at its head. Furthermore, typhoid fever will be the subject discussed in nearly all of this chapter : first, because it is always with us, in spring, summer, fall, and winter, in Manila, Cuba, home stations, or Alaska ; and, secondly, because it is transmitted in so many ways and is such an insidious enemy that absolute protection against it will probably insure against the other " water-borne " diseases.

The exciting causes of these diseases are as follows : Typhoid fever is caused by a bacillus that is found in the bowels, gall-bladder, blood, and internal organs of those having the disease. It is an actively mobile organism, does not form spores, is easily killed by heat, and does not, under natural conditions, produce disease in other animals than man. Cholera is caused by a vibrio or spirillum, a spiral bacterium that occurs in the intestinal discharges and contents of persons having the disease. It is actively mobile, multiplies with great rapidity, and in so doing the spirals break into comma-shaped fragments, whence the name *comma bacillus*. It does not form spores, is easily killed by heat and acids, and succumbs to the action of sunlight more promptly than most other organisms. Dysentery is a name applied to a group of symptoms rather than to one specific disease. Consequently it is due to several causes. One class of cases is caused by a group of closely related bacilli that present some differences, another to an amœba, an animal parasite, and still others to other microscopic animal parasites.

Intestinal infestations with worms have already been discussed.

Every case of any of these diseases is derived from some other case. We may at times have great difficulty in tracing the origin of given cases, but the more we learn about the diseases the more evident the truth of the preceding sentence becomes. Diet and unsanitary conditions

**Origin
of Epi-
demics**

promote the spread of, but cannot originate, typhoid. The germs must be introduced by man and are so introduced. The man who first brings infection to a camp may have the fever and not know what is the matter; he may be recovering from it and still not know, or he may have had it and recovered long before, but be a "carrier" and continue excreting bacilli for years. Some of the conclusions of the Reed Board are as follows:—

"(1.) During the Spanish War of 1898 every regiment constituting the First, Second, Third, Fourth, Fifth, and Seventh Army Corps developed typhoid fever.

"(2.) More than ninety per cent of the volunteer regiments developed typhoid fever within eight weeks after going into camp.

"(3.) Typhoid fever developed also in certain regular regiments within three to five weeks after going into camp.

"(4.) Typhoid fever became epidemic both in the small encampments of not more than one regiment and in the larger ones consisting of one or more corps.

"(5.) Typhoid fever became epidemic in camps located in the Northern as well as in those located in the Southern States.

"(6.) Typhoid fever is so widely distributed in this country that one or more cases are likely to appear in any regiment within eight weeks after assembly.

"(7.) Typhoid fever usually appears in military expeditions within eight weeks after assembly."

"(43.) In addition to the recognized cases of typhoid fever, there were many short or abortive attacks of this disease which were generally diagnosed as some forms of malarial fever."

The above conclusions were reached before there was any recognition of chronic typhoidal carriers, and the

Carriers Board knew nothing of their existence. Since the investigation of the first recognized chronic carrier, many interesting cases have been studied and their histories worked out, and, as they illustrate the dangers from this source, a few of them will be detailed. The first case reported was that of the female proprietor of a bakery in Strassburg. All of her employees would sooner or later get typhoid, and the cause could not be located. It was at length ascertained that the woman, who had had typhoid ten years previously, was still passing very large numbers of typhoid bacilli in her stools. In a British reformatory there had been intermittent outbreaks of typhoid, three, four, or five cases occurring at a time. Inquiry pointed to milk as the cause, but inspection of the dairy and its surroundings showed good sanitary conditions. Nevertheless the disease continued to appear. Search among those handling the milk in the institution led to the examination of a girl who had had typhoid six years previously. Two examinations of her stools were negative, but the third showed that she was passing typhoid bacilli. Her removal from the kitchen ended the epidemics.

The first instance thoroughly worked up in America was a very instructive one. The investigation began with the efforts to trace a small epidemic, six persons in a household of eleven being attacked with the disease. Thorough investigation at the time of this outbreak and for some time subsequently failed to disclose the source of infection in the food, drink, or general sanitary conditions. At length suspicion was directed to a cook who had worked in the house for a few weeks, beginning about three weeks before the appearance of the disease. With great difficulty Dr. George A. Soper was able to trace part of her history during a period of ten years, in which she was known to have lived in eight families where inquiries could be made. In seven of these she had been associated with typhoid outbreaks, always escaping the disease herself. In the seven families there were twenty-six cases of typhoid, with one death. Owing to the cook's refusal to tell anything about herself, and the fragmentary character of the history obtained, it is probable that she had been associated with other cases. On the information obtained, however, the New York Department of Health caused her removal to the Detention Hospital, where she was examined, in spite of her objections and resistance, and was found to be passing great numbers of typhoid germs in her feces.

Another striking case is reported from Prussia, where typhoid had been epidemic on a large estate for fourteen years. During that time there had been thirty-two cases of the disease among one hundred and eighty persons on the estate. Investigations focused attention on the dairy, all of the persons involved having used milk from it. A woman employed there had had typhoid seventeen years before, and the bacteria were found in

her stools almost to the exclusion of other varieties. She had worked in the dairy fourteen years, and had been spreading the disease during that time. In a recent outbreak in Georgetown, D. C., the epidemic was traced to milk unknowingly contaminated by a woman who had had the disease eighteen years before, and was still passing the germs in her stools.

Such instances have multiplied of late, and many more could be cited, but enough has been said to indicate the very great importance of cleanliness on the part of cooks and other persons handling foods, and the necessity of investigating the kitchen in the case of company outbreaks. It is also worth repeating that good men should be detailed on kitchen police, and rewarded for efficient and cleanly performance of the duty, rather than that poor men should be detailed as a punishment, possibly for drunkenness, uncleanness, or other evidence of their unfitness. As showing that the danger from "carriers" is not slight, it may be stated that of 386 cases of typhoid occurring in one sanitary district in Germany, in which the source of infection was carefully sought, no less than seventy-seven could be traced to them.

Different series of examinations of the stools of persons who have had typhoid have resulted in estimates varying from 1.7 per cent to 5 per cent as the proportion of such who continued for indefinite periods to give off bacilli. One case of twenty-nine and another of fifty-six years' duration have been reported. In most instances the infection lingers in the gall-bladder, and it may give rise to the formation of gall-stones. In addition to the above-described class of chronic carriers, very many more cases give off bacilli in the stools for a

short time, and from 20 to 25 per cent pass them in urine for periods varying from weeks to months or, in rare cases, years. Rarely there are found carriers who give no history of typhoid, the disease apparently having been localized in the gall-bladder from the beginning.

Carrier cases are also known to occur from cholera and dysentery, and in the latter instance it is well known that the organisms, and, in fact, the disease itself, may be present for years.

The Reed Board said: "(48.) When a command is thoroughly saturated with typhoid fever, it is probable that one fourth to one third of the men will be found susceptible to this disease." It is probable that the susceptibility to cholera and dysentery among American troops will be even higher, as they have not had such opportunities of acquiring immunity at home as they have in the case of typhoid.

Suscep- tibility

The Reed Board reported: "(47.) The percentage of deaths among cases of typhoid fever was 7.6%" [a low figure]. — "(54.) The deaths from typhoid fever were 86.24% of the total deaths." — "(56.) The mortality from typhoid fever per 1000 of mean strength was 14.63."

Deaths

Typhoid fever is known to be transmitted in all of the ways to be considered, and it must be regarded as possible that cholera and dysentery can be spread in most or all of them.

Methods of Trans- mission

Water was not found by the Reed Board to be an important factor in the spread of typhoid in the national encampments in 1898; but it has so often been demonstrated as the most impor-

Water

tant factor in other epidemics, that all authorities agree in considering it one of the first subjects to be investigated in case of any outbreak. This method of transmission is so well recognized and so generally known that it is not necessary to cite any of the very numerous instances showing it. Prior to 1898 practically all of the great epidemics that had been studied were traced to infected drinking-supplies, and the disease was regarded as mainly, if not entirely, water-borne. This fact helps to account for the size and extent of the great camp epidemics of that year. Cholera and dysentery have also usually been traced to infected water. There is, however, the possibility that both may be transmitted in the other ways to be considered.

The germs may be ingested directly in polluted drinking water, and such is the usual method in large epidemics due to water. The danger in such instances of course increases with the dose of germs, that is, with increase in the degree of pollution or in the length of time the water is used. It may also occur in less obvious ways. Thus, water that is used in clothes-washing or bathing may become infected from the clothing or persons of sick men or carriers, and may later infect the hands or persons of the people handling or using it, or the vessels in which it was contained. Or men may contract the disease by bathing or swimming in infected streams or pools, and inadvertently taking water into their mouths, noses, or, possibly, their eyes.

Milk-borne epidemics of typhoid are also well recognized, and many of them have been thoroughly studied.

Milk The milk is, in the great majority of instances, contaminated from the hands of milkers, dealers, or other handlers who are suffering

from beginning typhoid and have not yet ceased work, who are convalescing from the same disease, or who have had it formerly and are still "carriers." The contamination may occur in other ways, such as the use of infected water for washing cans or diluting milk. It may also linger long in cans once infected, if they are not properly washed and sterilized, as the germs will multiply in milk. As the same thing is true of the germs of cholera and dysentery, they may also be milk-borne.

One of the latest and largest milk-borne epidemics of typhoid fever occurred in the suburbs of Boston in the spring of 1908. Four hundred and ten cases occurred, the infection being traced to a milk-dealer who continued his work for two weeks after the onset of the typhoid fever from which he later died.

Lettuce, radishes, and other vegetables or fruits that are eaten uncooked may be contaminated by unclean handling, by washing in polluted water, by contact with typhoid feces or urine in manure, by flies, or by infected dust blown and deposited on them. Bread, cool meats, and other food-articles may also be contaminated in some of these ways after they are cooked. Any article so polluted might cause the disease if it were eaten.

Vegetables and other Foods

Many epidemics of typhoid fever have been traced to the consumption of oysters and other shellfish gathered from sewerage-infected beds. For twenty-five years the city of Belfast had had the greatest mortality from typhoid of all the cities in the United Kingdom, in spite of a good water-supply and good general sanitation. A commission appointed in

Shellfish

1907 to investigate the matter showed that the endemic prevalence of the disease was due to the unrestricted gathering of cockles, mussels, and other shellfish from the "slob-lands" of Belfast Lough, which are laden with the city's sewage. Hundreds of acres are exposed at low tide, and the shellfish are gathered by the poorer classes, who generally eat them raw. The Jews, who do not eat them, were exempt from the disease. It is probable that cholera and dysentery may be transmitted in the same way.

Living and virulent typhoid organisms have been found in the mouths of persons sick with the disease, and the vomit of cholera may contain the germs of that ailment. It may therefore be understood that the common use of drinking-cups, clinical thermometers, and other articles that are put to the mouth, may lead to infection. In all of the diseases discussed in this chapter the organisms occur in the intestines, and careless use of rectal syringes may spread them. Such an accident is, of course, more apt to occur in hospitals.

None of these diseases is air-borne in the sense that smallpox and scarlet fever are, but feces, urine, or other material containing the causative organisms may be scattered as dust when well or partially dried and pulverized, and may be inhaled in that form, or may light on articles of food or drink and be ingested with them. The Reed Board reported: "(31.) It is probable that the infection was disseminated to some extent through the air in the form of dust."

"(28.) Flies undoubtedly served as carriers of the infection. Flies swarmed over infected fecal matter in the pits, and then visited and fed upon food prepared

for the soldiers at the mess tents. In some instances where lime had recently been sprinkled over the contents of the pits, flies with their feet whitened with lime were seen walking over the food. It is possible for the fly to carry the typhoid bacillus in two ways. In the first place, fecal matter containing the typhoid germ may adhere to the fly and be mechanically transported. In the second place, it is possible that the typhoid bacillus may be carried in the digestive organs of the fly and be deposited with its excrement.”

**Flies
and
other
Vermin**

So far as known, the above remarks describe equally well the relation of flies to cholera and dysentery infections, and infestations with intestinal parasites. The writer has seen larval roundworms that could not be differentiated from some species infesting man, in the proboscis of a fly. So far as known, no epidemics have been directly traced to roaches, yet persons familiar with these insects, and knowing how impartially they run in privies and kitchens, can readily see how they might spread disease, especially in the tropics. Recently the Director of Health and Chief Quarantine Officer for the Philippine Islands has advanced an interesting hypothesis to account for the persistence of cholera in Manila. He says: “The cholera organism is continuously present in some form in the sewer system of Manila, either in its fluid contents or in vermin. . . . It appears probable that the cholera organism is conveyed either directly or indirectly from the sewers to easily contaminated food or drink by the roaches, either through direct mechanical contact with their feet or bodies, or by the deposition of their excretions; thence it is passed to human beings; but flies, water-bugs, and other vermin com-

monly found in sewers are also no doubt partly responsible."

It would be well for the line officer to know the following facts concerning house-flies. They breed in horse-manure, human feces, and other filth, and can readily emerge through six inches of loose earth, but not through earth saturated with water. When breeding in feces and it dries, the larvæ will go into the earth five or six inches, and there find more congenial surroundings. They are naturally most numerous about trench latrines and kitchens, because they find in those places the best breeding and feeding places respectively. The female lays about 120 eggs, and the development of these varies with the temperature, somewhat as follows:—

	HOT WEATHER	COLD WEATHER
	Days	Days
Eggs hatch in	1	2
Maggots change to pupæ in	5	14
Pupæ change to adult flies in	3	5
	<hr/>	<hr/>
Total	Short 9	long 21 or more

Contact with sick persons offers many opportunities for both mediate and immediate infection, and the class of diseases under discussion often infect nurses and attendants. The typhoid patient, as stated before, may give off the germs of the disease from his bowels, bladder, or, occasionally, his mouth. From these sources the bacilli get on towels, bedding, urinals, chamber pots, the patient's skin, and into bath water. Any of these articles may be handled by nurses, associates, or casual visitors, who may thereby infect their own hands or persons, and later take the germs

into the mouth. Persons using the same drinking glasses, linen, or other personal articles, as the sick man, are especially apt to be infected. This method of infection is lately receiving much more recognition than formerly, and it is now prescribed in the army that typhoid fever cases shall be isolated and treated as though contagious. The greatest care in cleanliness and disinfection is necessary on the part of all hospital attendants and others brought in contact with the disease. The danger is of course greater if the nature of the disease is unknown, and the subject pursues his ordinary course of life, eating, drinking, sleeping, defecating, and urinating, with no thought of being a menace to others.

Tent-mates and bed-mates and members of the same family and household are particularly exposed. The Reed Board published the following conclusions that bear on the subject: "(30.) Typhoid fever, as it developed in the regimental organizations, was characterized by a series of company epidemics, each one having more or less perfectly its own individual characteristics. . . . Of 1608 cases of typhoid fever which we have been able to accurately locate in the particular tents in which they occurred, together with the date of the commencement of the attack, the results may be summarized as follows:—

"Directly connectable attacks, 563, or 35.01 per cent.

"Indirectly connectable attacks, 447, or 27.79 per cent.

"Total connectable attacks, 1010, or 62.8 per cent.

. . . "We believe, therefore, that personal contact was a very important factor, probably the most important, in the spread of the disease."

It will be readily understood that the dangers of contact would be greatly increased by crowding, poor ventilation, and uncleanly habits.

House epidemics of cholera are common in the Philippines during epidemic seasons, and owe their origin and spread to the causes discussed above, to the common use of food and drinking utensils, and to infection contracted in cleaning up vomit and watery stools. There is good reason to think that dysentery and diarrhœa may be spread in the same ways, while it is well known that intestinal worms often are, among children.

The above discussion of the dangers of contact with sick persons may be applied equally well to similar relations with carriers, except that they are not known to be sources of danger, and are thereby rendered more harmful. They are more apt than sick men to sleep with other persons, to lend or borrow clothing, to be put on duty in kitchens, or handling foods, or caring for cows, while no precautions are taken in the matters of disinfecting their clothing, stools, or urine.

It does not require lengthy argument to demonstrate that, if urine and feces contain disease-producing germs

**Soiled
Feet**

and are deposited on the surface of the ground about camps or are allowed to soil the edges and vicinity of latrines, men may get their shoes soiled with such matters and the contained germs. Such being the case, the contamination might easily be conveyed to tents, bunks, company streets, and the hands of the men. Needless to say, such infection could be conveyed even though all gross evidence of fecal or urinary soiling were removed.

This term includes such inanimate articles or substances as are thought capable of absorbing, preserving,

and transmitting the contagii of disease, and is used as a general designation for such things as books, clothing, tents, bedding, and baggage. The **Fomites** Reed Board reached the following conclusions bearing on the subject: "(32.) A command badly infected with typhoid fever does not lose the infection by simply changing location. (33.) When a command badly infected with typhoid fever changes its location, it carries the specific agent of the disease in the bodies of the men, in their clothing, bedding, and tentage. (34.) Even an ocean voyage does not relieve an infected command of its infection. (35.) After a command becomes badly infected with typhoid fever, changes of location, together with thorough disinfection of all clothing, bedding, and tentage, is necessary. . . .

"That the floors, furniture, and bedding of barracks may become infected with typhoid fever and may continue to endanger the life and health of its occupants for a long time is abundantly shown in the records of military surgeons. . . .

"Our investigations certainly demonstrate that typhoid fever is not only an infectious but also a contagious disease; that it may be transferred from one person to another by contact, and that the clothing, bedding, and rooms of typhoid patients should be disinfected with as much care as is now given to these matters in cases of diphtheria and scarlet fever."

Clothing that is grossly contaminated, such as sheets or shirts soiled by the diarrhœal discharges of sick men, is, of course, most dangerous, but the danger in such cases is apt to be recognized and provision made for it. On the other hand, bedding soiled by dirty shoes, tents that have been urinated upon, the

clothing of apparently well carriers, and other articles, may be infected and show no signs of it, and so cause great harm. Laundry-workers, hospital attendants, and tent-mates of the sick are naturally more liable than others to infection in this way; yet all may be exposed to it in time of epidemics.

The general measures for the prevention of these diseases are those hygienic precautions that have been discussed in preceding chapters. If cases do begin to appear in a command, it is evident that they have not been sufficient, and more vigorous steps should be taken to protect the men not yet infected. If they show in the neighborhood of the command, or along lines of communication with it, the same active measures should be adopted. The best prospects for the prevention or control of the epidemic will be offered if the following steps be taken.

One of the most important of these is early diagnosis of the dangerous cases. As long as their nature is unknown, they continue to disseminate the disease-germs; and such undiagnosed or wrongly diagnosed cases apparently had much to do with the spread of typhoid fever in the camps in 1898. At that time the methods of diagnosis were not so good as at present, and a positive diagnosis of typhoid could not usually be made until after the patient had been sick a week or more. Laboratory methods now permit of it sooner, but they are not always available, and the safe rule is to regard and handle suspected cases as though the disease were known to be typhoid.

The diagnosis is a matter for the medical officer to worry over, but the line officer can be of assistance if

he instructs his men to go on sick report if they are not well, and if he does not unduly encourage the notion that men only seek sick report to escape work. It is at any rate better that two or three frauds should be allowed to loaf for a time than that one sick man should be allowed to spread disease through the command. The Reed Board made the following findings that bear on the difficulties and importance of early diagnosis:—

“(46.) Army surgeons correctly diagnosed about half the cases of typhoid fever.”

“(14.) A man infected with typhoid fever may scatter the infection in every latrine in a regiment before the disease is recognized in himself.”

Diagnosis is of no value in the prevention of epidemics unless it leads to immediate steps to prevent infection through contact, soil-pollution, water-pollution, and in other ways. It **Isolation** should therefore lead to the isolation of all recognized or suspected cases, and such measures of care as will prevent their spreading the disease in any of the ways discussed in this chapter. Such measures must include screening from flies, disinfection of clothing and linen, sterilization of bath water, dishes and utensils, disinfection of stools, urine, and spit. Only the necessary attendants, and they well-trained ones, should have access to the patients. The isolation should at times include suspects who have been exposed to disease as well as the actually sick. This is particularly true as regards cholera. It is necessary in a less degree for typhoid, and still less for dysentery. Whenever resorted to, such isolation should more than cover the ordinary period of incubation of the disease. Cholera suspects should be isolated for five days, those of typhoid for three weeks

or until well. In the latter disease such strict isolation may not be necessary, but in case of severe epidemics it will be wise, and can be done in isolation camps, where the men can work, drill, and play. At times it may be advisable to quarantine entire organizations as suspects, or to direct the measure against towns or communities.

Isolation cannot be made an effective measure of disease-prevention unless it includes all cases of the

**Compul-
sory Re-
porting
of Cases**

disease in question, and, in many instances, all suspected cases. It is, therefore, essential that all cases and suspects be promptly reported to the central authority, in order that proper steps may be taken. It is only rarely that difficulty will arise in this matter as it regards the military personnel; but cases among civilians and residents of the locality in which the troops are quartered may be willfully concealed. Epidemics "hurt business," and often interfere with transportation and traffic, and for this reason and others it is not uncommon that great efforts are made to hide or deny their existence.

It therefore occasionally becomes necessary to make regular and frequent inspections and examinations of

**Health
Inspec-
tions**

all persons in camp, and less frequently of those in the neighborhood. If the community is under military control, this may be a relatively simple matter, but where such is not the case it may be one of extreme difficulty, and call for the display of much forbearance, good temper, and tact. When such measures are necessary, they should be under the charge and direction of medical officers who will see that they thoroughly accomplish their purpose, but at minimum of inconvenience, and that they are

conducted in such a manner as to gain rather than alienate the sympathy and the coöperation of the civil population and its medical practitioners. If many sick are found it may be necessary, in order to control the epidemic, to establish large isolation camps and special hospitals for civilians. The gathering of large crowds from many places, as in pilgrimages, fairs, and religious festivals, should be particularly forbidden during cholera epidemics. Health-inspections may also be made the means of much sanitary instruction, and may thus do great good in addition to that accruing from a knowledge of the location of cases. Such inspection, examinations, and instructions have been resorted to in times of cholera epidemics, and among peoples living under military control, with great success. They redound to the benefit of the civil as well as the military population. Inspections of troops for the presence of disease are not infrequently resorted to, and often do much good.

Every case of any of these diseases that is reported or discovered should be investigated as thoroughly as possible as to its origin. This investigation may show no results in single cases, but when the findings in many cases are compared they may give important information as to the cause of the epidemic. Thus it may be found that it is confined to persons using a certain water, to those consuming milk sold by a special dealer, or who eat the productions of a certain cook, or who have partaken of shellfish; or that the cases are otherwise connected in some way. The information thus gained may direct attention to the cause of an epidemic, and its removal may put an end to the whole matter. The tracing of epidemics is, however, often a difficult matter,

Investigation of Cases

and the detection of typhoid-carriers is particularly so. If, for instance, a carrier were put on kitchen police for a day and infected the food of a company, a dozen men might contract the disease as a result. They might develop it, however, in from one to three weeks, or possibly more, and meanwhile a large number of men may have worked in the kitchen, and many other factors having an apparent bearing may have come to light, so that all thought is diverted from the man really causing the trouble.

In very many instances the cause of the epidemic cannot be ascertained, or at least not soon enough to permit its removal in time to prevent numerous infections, and protection must be sought in the practice of such general preventive measures as are applicable. These should take into consideration the various methods in which the diseases are transmitted, and will embrace the following.

Cleanliness of persons, tents, kitchens, and camps is of paramount importance, and officers should encourage

General Measures it by precept, example, orders, and, if necessary, punishment. Those associated with the sick should be particularly careful as to their persons, surroundings, and food. They should keep clean tents, wash themselves frequently, disinfect their hands after handling the sick, their bedding, clothing, or other personal belongings, should be careful not to use the same drinking or eating utensils, and in other ways should avoid the dangers of contact. *Disinfection* of clothing, bedding, tents, rooms, barracks, urinals, latrines, and all the things used by the sick man, should follow his removal from the barracks, tent, camp, or temporary hospital. Articles that cannot be well disinfected, such as hats and leather gloves, should be

destroyed if probably infected. Latrines that are infected should be disinfected as well as possible, and if pits or other cheap arrangements, closed. *General good hygiene* should prevail. Sunshine and fresh air should be provided in abundance to destroy and dilute the poison; temperance, order and regularity of habits encouraged, overwork and idleness alike avoided, and cheerful and rational living enjoyed. The ground should be kept so clean as to leave no opportunity for fecal or urinary contamination of shoes, no breeding-places or food-supplies for flies or roaches, and no infected dust to blow about. The effect of not doing these things is seen more promptly in cholera than in typhoid, because the incubation period in the former is shorter and effects become manifested before causes are forgotten. In that instance, lack of sunshine, intemperance in food and drink, fear, and unhygienic surroundings are regarded as important predisposing causes of the disease. Increased care and attention should be given to the *disposal of feces, urine, vomit*, and other wastes, and to the *elimination of flies* and other insects and protection from them. *Cremation* of waste materials should be resorted to if practicable. Reed troughs, safe water-carriage, or, better still, incinerators, should be used to dispose of feces and urine. Men should be instructed to use only the latrines and urinals of their own companies so far as it is possible to do so, and under no circumstances to urinate or defecate on the ground. The Reed Board reported: "(15.) Camp pollution was the greatest sin committed by the troops in 1898."

Cleanliness of camp-sites and *good general police* are therefore of very great importance, both as preventing

accumulations of infective material and as keeping down flies and other vermin. The Reed Board published two important and pertinent conclusions bearing on this:—

“(36.) Except in case of the most urgent military necessity, one command should not be located upon the site recently vacated by another.

“(37.) The fact that a command expects to change its location does not justify neglect of the proper policing of the ground occupied. . . . A camp-site should be thoroughly policed up to the moment of vacating it. This should be insisted upon as a matter of military discipline, and camp commanders should regard proper attention to the sanitation of the site occupied by their troops as one of their highest duties, and its neglect as a crime.”

“(21.) Greater authority should be given medical officers in questions relating to the hygiene of camps.”

Recent experiments show that typhoid bacilli can live and retain their virulence in soil for several weeks or months.

When, because of want of foresight, neglect, or for any other reason, a command is located on a polluted site, a change may be desirable. As stated
Change of Site before, this cannot be depended upon to rid the command of infection and disinfection: isolation and the other measures herein advocated must continue to be used, but the change may at any rate reduce the intensity of the virus and the number of channels of infection. The dust blown about camp would be less dangerous, the number of flies smaller, and the opportunities for fecal contaminations fewer, in the new camp. The Reed Board considered the failure

to change from polluted or bad sites a factor in the spread of the epidemics in 1898.

“(16.) Some commands were unwisely located.

“(17.) In some instances the space allotted the regiments was inadequate.

“(18.) Many commands were allowed to remain on one site too long.

“(19.) Requests for change in location made by medical officers were not always granted.”

“(39.) In some of the encampments the tents were too much crowded. The danger of camp-pollution would naturally be increased by crowding.”

The water used for drinking, dishwashing, and similar purposes can, of course, be sterilized by boiling; but as many men seek other water to drink, it is desirable that only a pure supply be available, and if one purified by large filtration plants or other means is to be obtained, it should be. At times it may be necessary to take steps to guard the purity of a supply known to be good. Reservoirs, filters, or watersheds may need to be patrolled to prevent their contamination. Bathing in drinking-supplies should of course be prevented. The rule to be always followed in time of epidemics is, however, to drink no water not purified by heat. No matter how good a general supply is to-day, the possibilities of infection are so numerous, and so hard to guard against, that certainty as to the purity of to-morrow's supply cannot be entertained.

Water-Supply

Ice, unless made from distilled water and handled with all possible care, should not be allowed to come in contact with articles of food or drink during the prevalence of epidemics. It may contain germs frozen

in it, or they may have been deposited on the surface by improper handling.

The safest rule as to foods is the same as that just applied to water, — to take none that is not sterilized.

Food-Supply

This rule is pretty generally applied by Americans in the Philippines, especially during cholera epidemics, but it has not had such general application in our own country or as a measure of defense against typhoid. Heat is the means of sterilization for most foods, and cooking the method of applying it. However, such articles as bananas, mangoes, apples, cucumbers, melons, that are protected by a thick skin or rind, may be washed in bichloride or other antiseptic solution, and, later, peeled. The use of pies, soft drinks, milk, and similar articles sold by peddlers, is in general to be deprecated, and it may be advisable to stop such sales. Green vegetables may also have to be banned, particularly in cholera times and where dysentery and the practice of manuring with human feces coexist. If either typhoid or cholera appears in the person or family of a dealer in food-stuff or milk, his business should be stopped at once, or allowed to continue only under such rigid rules as will destroy all chance of dissemination of the infection. If the person cannot be controlled by military authority, the soldiers should be forbidden to deal with him.

Typhoid, cholera, and dysentery are all diseases in which more or less protective value has been attributed

Vaccination

to vaccination. The degree of protection against cholera that is afforded by this measure is considerable, and its very extensive use in India, Japan, and other countries has given results that assure it a prominent place in the

suppression of epidemics. The methods of so-called vaccination vary considerably, but all have the common feature that the germ-substance, either living, dead, or extracted, is injected beneath the skin. This gives rise to varying degrees of local reaction, and some fever and discomfort that lasts from a few hours to a few days, but it results in the formation in the blood of anti-bacterial and immunizing substances that in most instances fully protect against the disease. Anti-typhoid inoculation or vaccination has had a less extensive use, and such as it has had has been almost entirely in the British army. The results there, however, have been sufficiently favorable to cause an extension of the practice, so that it is now very generally used. The accounts thus far published indicate that the method is of great value, and that it may yet come to be the practice in our own country to vaccinate new troops before sending them on campaign or to great camps of concentration. The method at present in use consists in the injection of a large number — 1,000,000,000 to 2,000,000,000 — of germs that have been killed by exposure to a temperature not in excess of 60° C. This, as in the case of cholera, results in the formation in the body of anti-bacterial and immunizing substances; but if the injection is made during the incubation period of the disease, or if infection takes place very soon after, the fever is both more certain and more severe in its development.¹

¹ S. O. 279, Dec. 1, 1908, WAR DEPT.

(Extract.)

12. A board of medical officers to consist of Brig. Gen. Robert M. O'Reilly, Surg. Gen.; 1st Lieut. Victor C. Vaughan, M. R. C.; 1st Lieut. William T. Councilman, M. R. C.; 1st Lieut. John H. Musser, M. R. C.; 1st Lieut. Alexander Lambert, M. R. C.; 1st Lieut. Simon Flexner, M. R. C.; 1st Lieut. William S. Thayer, M. R. C.; Capt. Frederick F.

Anti-dysenteric vaccination has been said to be of value, but it has thus far not had sufficient trial, or given sufficiently satisfactory results to make its use advisable.

Russell, M. C., recorder, is appointed to meet in Washington on Dec. 7, 1908, for the purpose of investigating the question of the advisability of immunizing the Regular and, in case of war, the Volunteer Army against typhoid fever.

The board met at 2 P. M., all the members being present. It proceeded to review the history of vaccination as a method of protecting troops against typhoid fever. The experience of both the English and the German armies was considered.

The theory on which the practice is based as well as many details involved in carrying out the procedure were considered, as was also the probable effect on the medical profession and the public at large in the United States.

As a result of several hours' discussion the board adopted, without any dissenting voices, the following conclusions: —

I. The board is convinced that the practice of anti-typhoid vaccination is both useful and harmless, and that it offers a practicable means of diminishing the amount of typhoid fever in the Army, both in times of peace and war.

II. It finds that the experience to date with anti-typhoid vaccination justifies it in recommending the introduction of the practice in the Regular and Volunteer Armies in time of war.

III. It recommends the immediate introduction of the practice of voluntary vaccination against typhoid in the Hospital Corps, the Army Nurse Corps, and in any expedition of troops from the Regular Army which is ordered to take the field for active operations; and, further, that an opportunity be given to Volunteers from the Army as a whole to be protected by vaccinations against typhoid.

There being no further business before it, the board adjourned *sine die*.

R. M. O'REILLY, *Surgeon-General, U. S. A., President.*

CHAPTER XIII

AIR-BORNE AND FOMITES-BORNE DISEASES

THE terms "air-borne" and "fomites-borne" do not tell the whole story of the methods of transmission of the diseases here to be considered, just as "water-borne" failed to tell it of those discussed in the preceding chapter. Most of those now to be treated of are also known as contagious diseases, though the term is elastic and may be used in different senses. As ordinarily applied to those diseases, it is understood to mean that the subject of one of them is surrounded by a zone of atmosphere in which the poison or contagium is present and which may convey the disease to persons entering it.

Contagious Diseases

The principal diseases in this group are measles, scarlet fever, German measles, smallpox, chicken-pox, influenza or gripe, mumps, whooping-cough, and diphtheria. Many other diseases may be borne by fomites and by dust-laden air, among them tuberculosis, typhoid and cholera, tonsillitis, and possibly pneumonia and meningitis; but the bulk of this chapter will be devoted to the contagious group, especially measles, scarlet fever, and smallpox.

Some of the diseases mentioned above—influenza, tuberculosis, diphtheria, meningitis, and tonsillitis—are known to be caused by bacteria, and the organisms can be isolated and identified. Smallpox is probably due to a minute animal parasite, but the relationship is not

Causative Organisms

generally acknowledged. The causes of measles, German measles, scarlet fever, and whooping-cough are still unknown, and it is possible, though not certain, that some of them are due to organisms too small to be seen by the microscope. On the other hand, different investigators have at various times described what they thought to be the causative organisms. The chances are rather against great advances being made in the management of these diseases until their causes are known, though such is not necessarily the case. Vaccination was found to be preventive of both smallpox and hydrophobia, though the causative organisms were unknown; yellow fever has been brought under control despite the fact that we do not know its cause, and typhus has been all but exterminated and we are still ignorant as to what produces it. Nevertheless, vaccination, mosquito destruction, and good general hygienic measures have failed to do away with measles, scarlet fever, and influenza.

Certain factors have an important influence in predisposing to some of these diseases, and enable us to exercise some measure of control over them.

**Predis-
posing
Causes**

Most of them are so prevalent in the *very young* as to be known as "the diseases of childhood." By this phrase we usually speak of the group including mumps, measles, scarlet fever, German measles, chicken-pox, and diphtheria. Before the era of vaccination, smallpox was also principally a disease of childhood, and produced its greatest mortality among those under five years of age. Thus in "Geneva from 1580 to 1760 there were 25,349 deaths from smallpox, and of these 21,078 were under five years old and 961 per 1000 were under ten years old. In

Edinburgh in 1764-83, the proportion under ten years old was 993 per 1000. In the Chester epidemic of 1774 there were 202 deaths, all among children under ten years old." All of these diseases do attack adults, however, though less frequently and usually less severely than children.

Season exercises some influence. The cold, raw weather of the late winter and early spring particularly favor most of them, while they are also common in winter and rather rare in summer.

Crowding favors epidemics, quite apart from the question of ventilation, and it also predisposes to severe attacks. Many cases of a very severe type of measles developed in the concentration camps in 1898. School epidemics of measles, German measles, scarlet fever, mumps, and diphtheria are rather common.

Poor hygienic conditions, such as dirty surroundings, the breathing of effluvia from decomposing matter, and bad ventilation, are all thought to predispose to them.

Unknown atmospheric conditions apparently influence the rise and fall of some of them. Influenza and diphtheria increase and decrease in ways thus far not fully explained.

A previous attack of one of these diseases usually protects from a second, but such is not the case in all of the group. One attack of diphtheria or influenza, for instance, seems to predispose to a second rather than to protect against it. Second attacks of smallpox, scarlet fever, and mumps are rare, those of measles less so.

These diseases are not all transmitted in exactly the same way, but the transfer of no one of them is so limited as the title of the chapter would indicate; so the various

**Methods
of Trans-
mission**

methods will be considered and the exceptions noted as we progress.

Some of these affections are truly contagious, as the word was defined earlier in the chapter. Mere approach

Air

to the patient and breathing of the air surrounding him in a greater or smaller zone seems sufficient to allow infection by smallpox, measles, scarlet fever, typhus, German measles, mumps, whooping-cough, and oftentimes influenza. In some of these instances, and in diphtheria, meningitis, and some other diseases, the infection may occur through the inhalation of dust or of droplets of sputum expelled in coughing, and which contain the organisms. In measles, mumps, smallpox, and scarlet fever this is not always the case, and the only explanation that will account for their high degree of contagiousness is that the virus is floating in the air in an impalpable and unidentified form. It may even be carried to a considerable distance at times, and smallpox hospitals, in particular, are apt to be a menace to their neighborhoods. It is not understood how other aerial conveyance than by droplet or dust infection can well occur in influenza, yet the rapidity with which the epidemic of 1889-90 spread over the whole world makes it seem possible that such exists. The periods during which these diseases are most contagious vary considerably and will be discussed later.

The contagious principles of most of the group here discussed are well known to cling to such inanimate

Fomites

articles as books, clothing, bedding, and carpets, and they may sometimes be preserved in these things for considerable periods. Toys may serve as the transferring medium, and measles, German measles, diphtheria, and scarlet fever have

often been conveyed by them. Handkerchiefs and clothing soiled by the nasal discharges, expectoration, or skin effluvia of the sick are particularly dangerous. It is often extremely difficult to draw hard and fast lines separating pure contagion (in the above defined sense), dust and droplet contagion, fomites-conveyance, and contagion by contact. Where there has been a possibility of infection in one of these ways, there have often been other opportunities for it.

Contact-infection we know to be possible in all of them, though, and kissing or sleeping with a person with any of the diseases under discussion is particularly dangerous. They are dangerous **Contact** cases to nurse or to treat, and nurses and doctors often contract them, and would do so much oftener except for the protecting influences of previous attacks, age, and constant sanitary precautions.

The virus of most of these affections is given off in the spit and nasal secretions of persons suffering from them. Such is certainly the case in diphtheria, scarlet fever, measles, mumps, and **Sputum** sometimes in meningitis, and it may be true of all the others. Such relations as bring healthy persons into contact with these substances are therefore particularly dangerous, and the handling or using of towels, handkerchiefs, and similar articles used by the sick is to be carefully avoided. Droplets of sputum or mucus expelled by coughing or speaking, and the inhalation of them, are frequently a cause of disease. One case of diphtheria or measles in the early stage, and showing no symptoms except cough or "cold," may infect a whole school of children, or a barrack full of soldiers, before its nature is recognized.

It is quite probable that the virus of some or all of these diseases may also be contained in other natural secretions of the patient, or in pus or eruptions that sometimes occur. It is highly probable that the ears may be infectious in measles, and discharges from the ears and nose are certainly so in scarlet fever. Other parts of the body than the throat and nose may be infected with diphtheria, and discharges from them would be highly contagious. The disease may attack the eyes, the genitals, the rectum, wounds, and, rarely, the sound skin and stomach.

Other Secretions In smallpox, measles, scarlet fever, German measles, and typhus, it is either certain or probable that the virus is given off from the skin, and that contact with that may give the disease, whether it be direct, as in handling the patient, or indirect, as in handling clothing, bedding, or animals touched by him. In several of these diseases there is a marked shedding of the upper layers of the skin as convalescence begins. The part shed may be in the form of thick scabs, as in smallpox, strips, sheets, or large scales of parchment-like skin, as in scarlet fever, or fine branny scales, as in measles or German measles. Whatever their form, these products are dangerous and are to be avoided.

Skin In most or all of the diseases here considered the virus is at some time contained in the blood, and contact with that, as in dressing wounds or cleaning up spilled blood, would probably offer chances of infection.

Blood The introduction of infectious matter into wounds serves to inoculate most of these affections. It is much resorted to in efforts to infect animals for experi-

mental or other purposes, and in pre-vaccination times inoculation with smallpox was sometimes done as tending to produce a milder attack of the disease than resulted from natural infection.

Inoculation

Infection through the medium of utensils, such as cups, spoons, thermometers, bedpans, or pipes, may readily occur, but is an evidence of mediate contact rather than of contagion by virus contained in the air. Such articles can only rarely convey the disease in question, unless they have been soiled by contact with the mouth, skin, or other portions of a patient.

Utensils

Animals may convey these diseases in various ways. Their hair or fur may absorb the poison from the air of the contaminated zone surrounding the patient and convey it as a window-curtain or similar article might; or they may gather up infected dust, sputum, or flakes of skin by lying on or near the sick-bed, and, by shaking themselves, liberate it elsewhere. It is possible that they might lick up moist sputum or other secretions, and retain the germs alive in their mouths and transmit them through fondling or fawning, even though they do not themselves have, or are not subject to, the disease. In other instances they are subject to the diseases, and may contract them through contact from eating spit, or in other ways, and breed and give off germs just as do human patients. Cats, dogs, and many other animals are subject to diphtheria, and may aid in spreading it. Cows suffer from a disease of the teats which, it has been said, can by means of milk set up scarlet fever in the human subject. Cattle suffer from smallpox in the mild and al-

Animals

tered form known as vaccinia or cowpox, which, when in turn inoculated into man, affords protection against the first, or more severe disease. Whooping-cough has been transmitted to a cat by feeding it the mucus and vomit from patients having that disease.

Epidemics of scarlet fever and diphtheria have many times been traced to contaminated milk, and it is not

Milk

impossible that several of the other diseases we are discussing may likewise be so conveyed. The milk may, in exceptional instances and in the two diseases just named, be infected by the animal giving it, but the virus is usually derived from some human carrier of the organisms. Milk is a good culture medium for most disease-producing bacteria, and we have already seen how important it may be in the diffusion of typhoid fever and how readily it may be infected with the germs of that disease. It is therefore easily conceivable how milk-handlers suffering from influenza, mild diphtheria, mild scarlet fever, or others of the diseases we have been discussing, might, through coughing, tasting the milk, careless handling of vessels, or in other ways introduce germs into it. Milk epidemics of scarlet fever, like those of typhoid, are apt to show a somewhat "explosive" character, that is, many cases develop at or near the same time. Measles is not thought to be milk-borne.

The virus of any disease having gained entrance to a susceptible body, some time is required for its multi-

Incuba- tion

Periods

plication and development, and for it to manifest its effects on the body. This time between the entrance of infection and the appearance of the first symptoms of disease is known as the incubation period, and is more or less

constant for each disease. As it is important to take this into consideration in attempting to trace, control, or prevent epidemics, the following may be considered as representing it with tolerable accuracy in the diseases named:—

DISEASES	INCUBATION PERIODS
Smallpox	8 to 20 days, oftenest 12.
Typhus fever	12 days or less.
Chicken-pox	10 to 15 days.
Scarlet fever	1 to 7, oftenest 2 to 4 days.
Measles	7 to 18, usually 14 days.
German measles	14 days or more.
Mumps	14 to 21 days.
Whooping-cough	7 to 10 days.
Influenza	1 to 4 days.
Cerebro-spinal meningitis	Unknown.
Diphtheria	1 to 7 days, usually 2.
Tonsillitis	1 to 3 or 4 days, usually 2.

The incubation period is of value in enabling us to approximate the time and place of infection and to judge correctly of the length of time that suspects should be isolated. After the disease has developed, however, the character and length of the isolation period for the sick will be controlled rather by the degree of the contagiousness and the time at which it is greatest or during which it persists. It will therefore be well to summarize briefly our knowledge on those points.

**Degree
and
Time of
Contagious-
ness**

Smallpox is one of the most contagious of diseases, and prior to the introduction of vaccination almost everybody had an attack, usually in childhood, as natural

immunity is very rare. It is probably contagious from an early stage, and it continues so until the patient has recovered from the disease and has ceased to desquamate or shed scabs. The contagium is very persistent, and clings for long periods to infected places or articles. Varioloid is a mild form of smallpox, usually seen in vaccinated persons, and may give rise to the most virulent type of the disease in those who have not been vaccinated.

Typhus fever was, during its height, considered one of the most contagious of diseases during all its course. There is at present a tendency to attribute its spread in former times to insect transmission, bedbugs and fleas being accused, but the facts are not determined.

Chicken-pox is probably contagious from its onset until after the completion of desquamation of the scabs.

Scarlet fever is most contagious when the fever is highest, but it is so even after the subsidence of all fever and for some weeks after the disease has apparently ended. This long persisting contagiousness may be due in part to virus contained in the desquamating skin, but it is probably more often due to a focus of inflammation persisting in the nose, throat, or ear. Infected places or articles may retain the virus for a long time, even for many months.

Measles is even more contagious than scarlet fever and smallpox, but fortunately the period during which it is so is relatively short. It is most marked during the period of onset of the disease, when the patient shows no symptoms but those of a cold. The contagium is more evanescent than in the case of scarlatina, and neither places nor things retain it so long.

The contagium of *German measles* is probably like

that of measles, active for a short time, but not persistent.

Mumps is contagious for about two weeks, or a little more, from the time of its onset.

Whooping-cough is probably contagious during the entire period of its existence, including the early stage before the development of the whoop.

Influenza, likewise, is probably contagious from the onset. It often passes into a chronic stage, wherein the active symptoms have subsided, but a slight cough or some similar reminder persists. As the bacilli remain and may be expelled by the cough, there is reason to think that such cases may remain sources of infection for indefinite periods.

Cerebro-spinal meningitis being due to an organism that may be found in healthy noses and throats, but one which possesses but feeble resisting powers, it is impossible to say how long it may be contagious. The degree of contagiousness is not high.

Diphtheria is contagious from its onset until such time as the bacilli have disappeared from the breathing passages, a period exceeding the apparent duration of the disease by days, weeks, or months. Different strains of the organisms present decided differences of virulence, and this partly accounts for the varying grades of severity shown by epidemics. The virus attaches itself to the clothing, the bedding, and the room in which the patient has lived, and has in many instances displayed great tenacity. "It has been found to live in blood serum for one hundred and fifty-five days, in gelatine for eighteen months, dried on silk threads for one hundred and seventy-two days, on a child's plaything, which had been kept in a dark place, for five months, and in

bits of dried membrane for from fourteen to twenty weeks. . . . Forbes isolated diphtheria bacilli from a vessel which was regarded as the cause of the disease in twenty-four families. The bacilli grow readily in milk without changing its appearance."

Tonsillitis is an inflammation of the tonsils due to any cause. It may, therefore, be diphtheritic, but as generally used and as intended here, the term signifies a tonsillar inflammation due to other organisms than that of diphtheria, and applies to the common "sore-throat" of winter-time. Though neither so dangerous nor so highly contagious as diphtheria, it occasions much discomfort and may spread through a barracks to a considerable extent, especially if the ventilation is poor.

The prevention and control of epidemics of these contagious diseases naturally rests on two classes of measures,—those of general hygiene and cleanliness, and those directed at special features characteristic of the affections.

Prevention and Control

The general measures should include good *policing* and *cleanliness* of camps or barracks and of persons. It cannot be said that any disease in

the list under discussion, with the exception of typhus, depends very directly upon the presence of dirt or filth; yet the other disease-favoring conditions that accompany dirt, such as easier contaminating contact, poor ventilation, and vermin, do favor their spread. Good policing is one indication of good general sanitation.

General Measures

Ventilation should at all times be abundant, and it should be superabundant in the neighborhood of the sick, as it not only promotes comfort and health, but

dilutes the poison or contagium, and so reduces the dose of it that those in the neighborhood may inhale.

Water is of only minor importance in the diffusion of these diseases, but the supply should be good and special care should be used in disposing of that which has been used in washing the sick or their effects. It should be disinfected by chemicals or heat.

Foods, other than milk, are seldom responsible for the spread of the diseases considered; but food that has been in contact with the sick should be carefully disposed of. The diphtheria bacillus, for instance, might grow on meat, bread, potatoes, or other food-articles.

Persons having any of these diseases should be excluded from kitchens, dining-rooms, and all contact with any food except that for their own use and consumption. Supplies from bakeries, meat-shops, or other food-distributing places that harbor persons sick of these diseases, are best avoided. It is conceivable that they may occasionally be transmitted by means of food that has been exposed for sale in places where it might be contaminated by infected dust or by droplets of spit expelled by coughing persons.

Milk is particularly dangerous as a means of dissemination of scarlet fever and diphtheria, and it may possibly act in a similar capacity for others of these diseases. It is certainly the part of wisdom to exclude persons having, or recovering from, or in contact with, any of these diseases, from handling milk intended for the use of others. In case of epidemic the milk-supply should be investigated, and steps be taken to prevent infection through it. Pasteurization or boiling will make the milk safe if it is protected from subsequent contamination.

Vermin should be guarded against as possible disseminators. The spread of typhus has been credited to both fleas and bedbugs, and it is conceivable that either might carry the virus of smallpox, scarlet fever, or other diseases, from one person to another. Flies that pass from infected sputum to food-articles or to well persons might also carry it.

Animals, such as dogs, cats, and other pets, should be excluded from access to the sick for reasons set forth earlier in the chapter. The principal measures of specific protection against these diseases are as follows:—

Early diagnosis is absolutely essential for the control of most of them, and is helpful in all. Measles is highly contagious before the eruption appears, and may be widely spread by persons apparently suffering from “colds.”

Early Diagnosis All suspicious cases should therefore be reported to the surgeon for examination as soon as any symptoms present themselves. As stated before, company officers should encourage their men to go on sick report at the first appearance of disease.

Notification of cases is the next step in control. It follows diagnosis naturally in the case of soldiers reporting to the surgeon; but occasionally officers or their families, or more rarely enlisted men, consult civilian physicians. In such instances the presence of contagious disease may not be notified and may escape official recognition, with the consequence that preventive measures may not be adopted. Line officers, enlisted men, and civilian physicians, all should realize that the highest function of the medical officer is the prevention of disease, and that

they are preventing his performance of it, as well as violating orders, when they do not promptly report to him any cases of contagious disease in the post with which they may be concerned. Unless agreeing to act thus fairly, civilian physicians should not be allowed to practice on a military reservation, and they cannot properly do so; and officers and men who through carelessness or design are responsible for the presence of such diseases on the post being unknown, should be punished.

Notification of the presence of any of the diseases in this group should at once lead to an investigation of its source, with the object of removing this if possible. The nature of the investigation will be indicated by what has been said as to methods of infection and incubation periods.

Tracing Cases

The sick should be isolated as soon as the diagnosis is made or suspected. The character and duration of the isolation will vary with the disease. It should in all cases be sufficiently strict to prevent all unnecessary contact with the sick, and should last until the dangerous period is safely past. Smallpox should be isolated at some distance from dwelling or gathering places, even at the expense of some inconvenience. In military life tents make suitable isolation hospitals unless the weather is unusually severe. The possibility of apparently well persons acting as carriers must be borne in mind. Diphtheria patients and their attendants must not be released from isolation until their respiratory passages are free from germs, scarlet fever patients until all discharges from the throat, nose, and ears have ceased, and the individual has been well a month or more. In all instances the character

Isolation

and duration of isolation are matters to be determined by the medical officer. Men breaking isolation bounds should be severely punished, as they may expose their comrades to grave danger. Persons known to be exposed to these diseases may properly be isolated, or at least kept under strict surveillance, for a time equal to the maximum period of incubation. Those suspected of having any of them should be isolated until the diagnosis is definitely established in the negative.

Everything that has been contaminated by the use, association, or proximity of the sick should be disinfected before being allowed to pass again to its ordinary place.

Disinfection

Persons and *animals* should be well scrubbed with soap and hot water, and later with weak carbolic or bichloride solution, special attention being given to the hairy parts and the hair itself. Sterile or new clothing should then be put on. *Clothing* may be disinfected by boiling, steaming, soaking in antiseptic solution, or by formalin. The other methods are more reliable than the last.

Houses are best made safe by washing or spraying with warm antiseptic solution, making sure that all surfaces are covered and all cracks penetrated, and then fumigated with sulphur or formaldehyde. After that they should be opened and well dried and aired. Fresh calimine or paint may then be applied.

Utensils may be boiled, steamed, or washed in antiseptics. Letters, books, and similar articles may be sterilized by heat or by formalin, or by a combination of the two. The *discharges* of the patient should be mixed with formalin or carbolic solution and allowed to stand an hour before being thrown out. The *sputum*

should receive particular care, as it causes the contagium of most if not all of these diseases. The patient should only spit into vessels containing antiseptic solution, and even then it is safer if the spit be burned. Handkerchiefs and cloths used to wipe the mouth or nose should be burned or boiled. Surgical dressings should be burned.

The most efficient measure for the prevention of *smallpox* is the induction of acquired immunity through *vaccination*. The government very properly demands that every recruit shall be vaccinated, but occasionally the inoculation does not "take" on a susceptible man, and through a combination of circumstances he may escape its repetition. In civil life there is a considerable agitation against the use of vaccination, the agitators alleging that it does not protect against smallpox, that it is an invasion of personal rights, and indulging in several other incorrect statements. It is not purposed to discuss the matter at length here, but it may be stated that the evidence that vaccination does protect against smallpox is overwhelming, and a belief to the contrary, however honest and earnest it may be, cannot justify any one in endangering the efficiency and safety of an army by its neglect, any more than can a belief that every man is born with a right to smallpox if he wants it. Any officer or man who in any way interferes with the complete and thorough vaccination of the army makes a great mistake and commits a grave offense.

Induced Immu- nity

The spread of *diphtheria* may be partly controlled by the use of *antitoxin*. This may be and is very generally used in the treatment of those sick with the disease, but it is also of great value when used as a

preventive measure. It is then injected in relatively small doses into persons brought into contact with the disease. So used, it greatly lessens the liability to infection, and to severity in case infection does occur. Vaccinations and the use of preventive injections of antitoxin have not yet proved of value in protecting from the other members of this group, though they may later be found applicable in some instances.

CHAPTER XIV

INSECT-BORNE DISEASES

THE principal human diseases that are now known, or thought, to be insect-borne are malaria, yellow fever, filariasis, and dengue, carried by mosquitoes, sleeping sickness by the tsetse fly in Africa, plague by fleas, spotted fever of Montana and relapsing fever of Africa by ticks. Kala-azar of India, typhus and relapsing fever of Europe and America, and leprosy have all been accredited to the agency of the bedbug, and in the first two instances there is much probability that the parasite is the actual carrier. The proof of its relation to leprosy is, however, very slight as yet. These diseases will be discussed in groups made to conform to the insects transmitting them.

MOSQUITO-BORNE DISEASES

The most widespread and common of these is malaria. It prevails in most tropical and sub-tropical countries, and in some of them it produces a great mortality, while it causes much suffering, weakness, and loss of time, even where it is less fatal. It is an important cause of sickness in our army in many parts of our own country, as well as in Cuba, Porto Rico, the Philippines, and the Isthmus of Panama.

Since the beginning of the Spanish-American War the knowledge of the methods of control of this disease has increased so rapidly that it has now become

possible for us to make healthy communities out of places formerly notorious for the certainty and speed with which newcomers sickened or died from malaria. As stated in a previous chapter, there are at least three varieties of this disease which differ because the parasites causing them do so, and which manifest themselves in paroxysms of chill, fever, and sweat, that come at twenty-four, forty-eight, or seventy-two-hour intervals. This regularity of manifestation is only shown, however, in cases of uncomplicated single infections. Owing to multiple infections or complicating circumstances, any one of these types of malaria may show extreme irregularity and may be mistaken for other diseases, the diagnosis being made possible only by examination of the blood, and, at times, with great difficulty. The fever may be continuous and the general symptoms such as to make the disease appear much like typhoid. Or it may be manifested as a neuralgia, as a sudden attack resembling apoplexy, as anæmia, or in other obscure ways. This disease also exhibits periods of latency, during which the parasite is present in the body, but in such numbers or under such control by the body forces, that no signs or symptoms are present. When, however, the vital forces are reduced, as by exposure, overwork, alcoholism, or other disease, the parasites gain the ascendancy and the disease may manifest itself in a severe manner. Many natives of highly malarious regions may be partly immune and show no symptoms even when harboring very severe infections.

Whatever the character of its manifestation, however, the infection is always received through the bite of a mosquito. Many officers from time to time express views to the contrary, usually basing them on their own

faulty observations. Occasionally there is reason to suspect that the views are exploited to account for the presence of malaria when the efforts to prevent mosquito-transmission entail an inconvenient amount of trouble. The fact is that malaria has been investigated with a thoroughness that has been accorded to few other diseases, and the method of mosquito-transmission has been abundantly proved, and no other appears even probable. "If, after they have imbibed malarial blood, certain species of mosquitoes be dissected at serial intervals, the evolution of the malaria parasite can be followed in their tissues, until, finally, the germs of the parasite can be tracked into the cells and secretions of the salivary glands of the insect.

"If after a week, or thereabouts, a similarly fed mosquito bites a hitherto uninfected man, in many instances, after a few days, that man will exhibit the clinical phenomena of malaria infection and the characteristic parasite in his blood.

"A non-immune, if effectually protected against mosquito bite, will not contract malarial disease, however long he may live in highly malarious localities.

"Most observers are now of the opinion that the malaria parasite, under natural conditions, can be acquired by man only through the bite of the mosquito; that the mosquito can acquire the parasite only by ingesting the blood of a malaria-infected man, or possibly other mammal; that there is no extra-corporeal life other than that described; that there is no authentic instance of malaria being acquired in uninhabited places; that in the case of malaria in connection with soil disturbances, it depends on the creation during digging operation of puddles of water in which mosquitoes breed; and that

its epidemic occurrence under these circumstances is owing to unhygienic conditions such as usually prevail when large bodies of men, some of whom may bring the infection with them, are brought together on public works attended with extensive earth-cutting, as in railway, road, or canal-making."

Several species of mosquitoes are capable of transmitting the disease, but they all, so far as now known, belong to the sub-family *anophalinæ*. The features by means of which these various species may be identified cannot be discussed here, and positive identification is a labor necessitating some time and effort even on the part of the specialist; but a popular rough method of identifying *anophalinæ* mosquitoes in general is that the adults, when at rest, stand with their bodies almost or quite at a right angle with the surface on which they are resting, while commoner species rest with the body more nearly parallel to the supporting surface. The larvæ, or "wigglers," of the former class, on the other hand, rest with their bodies like floating sticks, parallel to the surface of the water in which they are swimming. The more common varieties have theirs more nearly at a right angle to it. These mosquitoes bite principally at night.

This is an acute and very fatal disease that occurs principally in tropical and sub-tropical America, though epidemics also occur on the west coast of Africa, where the disease was possibly introduced by slave-traders. It has appeared in epidemic form as far north as Philadelphia.

Until after the American occupation of Cuba and the subsequent work of the board of army medical offi-

cers, under the presidency of Major Walter Reed, the method of infection was unknown, and water, contact, fomites, poor police, and atmospheric conditions were all credited as means. That Board cleaned up all obscure questions as to the manner of transmission, and established beyond dispute the fact that yellow fever is transmitted by the bite of infected mosquitoes of the species *Stegomyia fasciata*. It cannot be transmitted by water, food, fomites, air, or contact, nor is its spread influenced by poor policing, except as the latter promotes the breeding of mosquitoes. The beautiful reasoning and experiments which served to establish our knowledge of the means of transmission also cleared up many hitherto obscure points about the disease, and, better still, led to the use of anti-mosquito measures that resulted in its eradication from Havana, and later from New Orleans, the Canal Zone, and other regions. This constituted one of the greatest sanitary advances that the world has ever seen, one making possible the rejuvenation of Cuba, the construction of the Panama Canal, and, it is hoped, the civilization and development of all tropical America; and it is well that army officers should keep in mind that all of these things were made possible by the fact that there happened to be in command, in Havana, a line officer who had a knowledge and an appreciation of the importance of military hygiene.

**Method
of Trans-
mission**

The Board of which Major Reed was the head showed that contact, fecal contamination, handling of fomites, breathing the close air breathed by yellow-fever patients in poorly ventilated places, and all such things, were incapable of transmitting the disease, but that the bite of some infected mosquitoes would be followed

by the fever in from three to five days, or occasionally longer. This is the ordinary incubation-period in man, and accounts for the fact that non-immunes may develop the disease so short a time after arrival in places where the disease prevails. But there is also an incubation-period in the mosquito, as shown by the fact that it cannot transmit the disease until twelve days after it has imbibed the infection with the blood of a sick person. This accounts for the half-month elapsing between the arrival in an uninfected place of a case of yellow fever and the appearance of the first case of the subsequent epidemic. This double incubation period, three days for a case in an already infected district, and fifteen or more days for secondary cases in previously non-infected districts, is thus clearly explained.

Not nearly all mosquitoes biting yellow-fever patients are capable of transmitting the disease even after twelve days, and the Board showed that this was because of the short period during which the yellow-fever patient is infectious. Unless it bites him during his first three days of sickness, the insect obtains no infection and can transmit none. It was also demonstrated that the cause of the disease is in the blood, that it passes through very fine filters, and is probably too small to be seen with our microscopes, and that it may persist in infected mosquitoes for weeks or months.

“These experiments fully explain, 1st, the impunity with which a yellow-fever patient can be visited by a non-immune if outside the endemic area: the mosquitoes in the vicinity are not infective. 2d, the danger of visiting the endemic area, especially at night: the mosquitoes there are infective and active. 3d, the discrep-

ancy between the incubation-period, three to five days, of the disease, and the incubation-period, fifteen days and over, of an epidemic: the necessary evolution of the germ in the mosquitoes infected by the original introducing patient demanding the space of time indicated by the difference between these two periods. 4th, the clinging of yellow-fever infection to ships, buildings, and localities: the persistence of the germ in infected mosquitoes which are known to be capable of surviving for five months, and probably longer, after feeding on blood. 5th, the high atmospheric temperature required for epidemic extension of yellow fever: such temperature favors the activities and propagation of the mosquito, and is probably necessary for the evolution of the germ also in the mosquito."

The mosquito that transmits yellow fever is a very common one in all those parts of America that show the disease. A closely allied species is abundant in the Philippine Islands. Whether or not this variety is capable of transmitting the disease is not positively known, but if it can do so the introduction of the disease into those islands, a thing quite possible after the opening of the Panama Canal, might prove a disaster of the first magnitude and a great addition to their already numerous afflictions. The mosquito is a small black one, with white or silvery lines and bands on the thorax and legs. There is a peculiar lyre-shaped figure on the back of the thorax which, together with the white bands on its legs, makes a rough diagnosis of species possible. The insects particularly prefer low-lying shores, but may be found in other places. They are essentially house-lovers, are active biters, and are

**The
Yellow-
Fever
Mos-
quito**

very quick, so that they are harder to catch than many other varieties. They breed in small artificial collections of water, such as barrels, puddles, cisterns, and partially filled tin cans. "The nature of the water appears to be a matter of indifference: it is found equally in rain or waste water, but more frequently in water discolored by decaying vegetable matter."

Dengue is a disease of very low fatality, but one which may occasion much suffering and anxiety, and which sometimes occurs in widespread epidemics.

Dengue The disease causes, on an average, about one week of absence from duty, and if the epidemic involves thirty or forty per cent of a command, a not unusual figure, the strength of the force will be much impaired. It prevails in the West Indies, the Philippines, our Southern States, and in most other parts of the tropical and subtropical world. Like yellow fever, it is a disease of hot weather and low lands, that has prevailed as far north as Philadelphia in summer-time. It is transmitted by a very common variety of mosquito, *Culex fatigans*, and possibly by other varieties, but not by contact, food, drink, air, or fomites. The incubation-period is from three to seven days, usually about four; the causative organism is contained in the blood, passes through fine filters, and is probably too small to be seen with the microscope. The disease resembles yellow fever in many respects, and has often been confused with it in America.

Filariasis is infestation with any one of several varieties of filaria, or hair-worm, whose young circulate in the blood. It is thought to be related to **Filariasis** elephantiasis, but was sufficiently discussed in a preceding chapter. The parasites are introduced

through the bites of mosquitoes, one of the most common and widespread species, *Culex fatigans*, acting as a carrier for at least two varieties. The insect bites both by day and by night.

The prevention of these mosquito-borne diseases rests entirely upon our ability to protect against the bites of infected insects, except in the case of malaria, in which we can derive aid from the administration of quinine. Our success then depends on anti-mosquito measures, and it is necessary to discuss these briefly. Mosquitoes breed only in water, the young being as much dependent on that fluid as are fish or tadpoles. The *Culex*, or common mosquito, which transmits dengue and filaria, and the *Stegomyia*, or striped mosquito, which carries yellow fever, generally breed in vessels of water, cisterns, or drains. The *Anopheles*, or malarial mosquitoes, breed "mostly in shallow puddles on the ground, in small ponds, in slow and small runnels of water, in pools of rain water, on rocks, on the margin of slow-running streams, and sometimes, but rarely, in vessels of water."

The larvæ or wigglers of all kinds of mosquitoes require at least six or seven days of life in water before they can develop into winged insects, and the time is usually longer, especially if the weather be cool or conditions otherwise unfavorable. Drying kills them and also destroys mosquito eggs. On the other hand, natural degrees of heat and cold do not necessarily destroy them if they remain in water, and in northern latitudes both larvæ and adults may survive several freezings during a winter and again become active when warm weather returns. Usually only female mosquitoes suck blood, and they do so because the rich nutriment hastens the

Anti- Mosquito Measures

ripening of their eggs, and these are generally deposited only after such a meal. As the mosquito fills herself to a degree that greatly increases her bulk and weight, and makes flight laborious and slow, she usually seeks a place near at hand in which to deposit her eggs. Partly for this reason it follows that most infested houses or localities breed their own mosquitoes.

“Mosquitoes, especially *Anopheles*, love thick undergrowth.” The facts above stated indicate the steps to be taken in warring against them, and of these, **Destruction of Breeding-Places** is probably the most important. The measures designed to attain this end are numerous. *Draining and reclamation* of swamp lands, the *filling* of pools, puddles, and depressions, the *building* of dikes or embankments to prevent flooding, and the clearing or narrowing of streams to give them a more rapid flow, are some of the larger measures that can be profitably employed in the vicinity of towns or permanent garrisons.

But efforts must not be confined to such large undertakings, because the chances are that less ambitious projects, carried out in or about the dwelling, may accomplish almost as much good, and it will be manifested more promptly. Such local measures will include the filling of small hollows in the yard or about the house, or cutting such outlets from them that they will empty quickly after rains, filling or preventing the formation of pockets or pools at the outlets of drains or under faucets, filling, emptying, or obliterating hoof-tracks, wagon-ruts, and similar small depressions, seeing that fire-buckets, fountains, drip-pans under ice-chests

and similar water containers are emptied twice a week, and all larvæ contained in them killed. Empty or partly empty bottles, boxes, tin cans, tubs, and flower-pots must be removed or placed in such positions or conditions that they will not catch and hold water. Useless joints or sections of bamboo must be removed, and pieces that are in use or that form parts of buildings should, if open, be bored at the lowest level of the joint so as to let all water run out. The water in tins or saucers under table-legs or elsewhere as a protection from ants should be kept clean and free from larvæ. It would be well to have such things filled with antiseptic solution rather than plain water. Sagging eave-troughs and drain-pipes and catch-basins, that do not empty promptly, may afford breeding-places, and should therefore be made right. Banana plants afford quite sufficiently large receptacles for breeding water at the points where the leaves diverge from the main stalk, and trees may provide them in knot-holes. All such places must be investigated and faults corrected. Irrigating ditches should run freely and with a good current when in use, and should drain dry at other times. They should not present stagnant pools at any time. Thick underbrush and high grass should be cut and cleared away, as it not only shelters and harbors the grown mosquitoes, but hides and preserves small collections of water and keeps them from drying. Cisterns, shallow wells, rain-barrels, and other large useful containers of water should be screened or tightly covered so that mosquitoes cannot gain access to them. Privies and cesspools should likewise be protected by netting or tight covers, so as to keep them free from the insects, and drain outlets should empty on smooth rocks, or be otherwise so arranged that pools

and puddles are not formed by their discharge. Other precautions may be necessary and the circumstances may suggest them. The object to be kept in mind is the doing away with all stagnant water.

Many of the measures outlined above, such as draining pools and emptying fire-buckets, are destructive of larvæ in that they allow these to dry. In some instances, however, it is not possible to do those things, and other methods of destroying larvæ must be resorted to. Mosquito larvæ, like all other living creatures, have their natural enemies, and these may be utilized. Many varieties of small *fish* devour them, and some of these, for instance the "millions" of Bermuda, are said to do the work so thoroughly as to quite prevent mosquito development in ponds stocked with them. Many varieties of fish are useful for this purpose, and ponds, streams, and marshes that create mosquito nuisances can be profitably stocked with them. Other water animals, such as *tadpoles* and some kinds of *beetles*, also eat them, and their growth may be encouraged for the purpose.

Moulds and other germs may also attack the larvæ and cause their death through disease. Such factors often make the task of artificially rearing mosquitoes for experimental work one of some difficulty, but, so far as known, they have not been used to reduce their natural occurrence. That the natural enemies are not alone sufficient to destroy all larvæ is evident from the fact that the latter are occasionally numerous in waters abounding with fish and tadpoles.

Chemical poisons may also be used, but they are dangerous and can only be applied to waters which are not

needed and which should, therefore, be drained away or otherwise disposed of.

Oil, preferably *crude petroleum*, is therefore much used as a means of destroying larvæ. It acts mechanically. The larvæ, though living in water, are air-breathers, and must come to the surface at intervals of a minute or so for that purpose. They breathe through a small pipe or tube which rises above the back near the tail, and which is projected from the surface of the water when air is to be taken in. If the water be covered with a film of oil, this little tube becomes clogged with the latter as it passes through it, and respiration is hindered or prevented so that the wriggler dies of suffocation. Other oils will answer as well, but crude petroleum, because of its cheapness, harmlessness in the amounts used, and its slow evaporation, is preferred. Refined kerosene spreads more rapidly and evenly, but it also evaporates more readily. The oil may be applied by sprinkling, spraying, or "painting" it over the surface. In the last-named method it is applied by means of a saturated rag on a pole that is rubbed or brushed about on top of the water. So long as the surface is well covered the method of application does not matter. The applications should be renewed as necessary, and the frequency of this will depend on several factors that must be taken into consideration. A slight current or a prevailing wind may drive all of the oil to one side of a pond or may remove it in a short time. In such cases the renewal must be more frequent, and must be on the side from which the oil flows. Light oils of course require more frequent renewal than the heavier ones that do not evaporate so readily. The surface of the water should not be left uncovered for more than three or four

days. If the water be drawn off through a pipe running well under the surface, oiling does not injure it for household uses and is therefore applicable to cisterns, rain-barrels, and other domestic supplies.

Small collections of larvæ-containing water can be easily emptied by sweeping or other means, and the larvæ promptly die. It is said that the common prickly pear cactus of the tropics can also be used in destroying larvæ. "The thick, fleshy leaves contain a mucilage which exudes when the leaves are chopped and added to water, the mucilage rising to the surface and forming a practically air-tight covering similar to that formed by oil."

This is a difficult and unprofitable method of getting rid of a mosquito nuisance, but it is sometimes necessarily resorted to for the purpose of ridding houses of infected insects, those, for instance, that have had opportunities to bite yellow-fever cases. They may then be killed by hand, or caught under bottles containing a bit of chloroform or other volatile poison; or, more effective still, the house may be tightly closed and fumigated with burning sulphur, tobacco, or insect powder. The insects are killed or stupefied by the fumes, and must be swept up and burned after the house is opened.

Certain measures have value as tending to discourage adult mosquitoes and to cause their abandonment of a neighborhood, even though not killing them. The most important of these are the removal of stagnant water in which they may lay their eggs and the cutting and clearing of underbrush, high grass, and vines on or in which they take shelter and rest when not busy. Smoking and anointing the hands and face

with certain volatile oils are advocated as measures of personal protection for the same purpose, but they are not to be relied upon. A hungry mosquito will bite through a layer of oil, and the most enthusiastic smoker cannot produce such clouds as to protect his ankles, hands, the back of his neck, and other parts.

The more effective method of protection from adults is in the use of screening materials either of wire or of soft fabrics. Houses with wire-screened windows and doors are in general use throughout our country, and add enormously to the comfort and safety of life in the hot season. They are still great rarities in the Philippines, however, and the fact that we have been there for over ten years and have suffered for that time from malaria, dengue, and endless vexation of body and spirit, but still keep our soldiers in unscreened barracks and quarters, is much to our discredit. We know better, but do not live up to our lights, and therefore, as a people, we deserve what we suffer, except that those who suffer most are not responsible.

**Protection from
Adults**

Bed-nets are in general use throughout all barracks in the tropics, and, if in good condition and properly cared for, they offer adequate protection during sleeping hours, unless the sleeper rolls or tosses in such a way as to bring part of his person against the net, in which event the insects feed through it. It is neither pleasant nor practicable to spend all of the day, or even most of it in bed, and bed-nets are therefore inadequate.

Head-nets are at times of value for night use in campaign, and rarely so for common use in the daytime. Most mosquitoes, though, do not bite out of doors in the daytime, especially persons who are in motion.

Adequate clothing is of course important in protecting from mosquito bites, and should be used for that purpose. In spite of all precautions an occasional mosquito will manage to bite the most careful man. It is therefore important to see that the insects do not become infected with disease, and yellow-fever, dengue, and malaria patients, particularly the first-named, should be kept screened or doubly screened during the entire time that they are capable of giving infection. For this purpose it is essential that all cases be known, and early diagnosis and prompt notification are therefore important, especially in yellow fever, which is so deadly and yet is infectious only in its early stages. In addition to screening, the malarial patient should be taking quinine to free his blood from germs. As natives of malarious regions are very apt to harbor the parasite in their blood and to keep the mosquitoes in their neighborhoods infected, it is important to keep them away from barracks and healthy households, and it is wise to build dwellings or barracks at some distance from native towns or habitations. Natives working about soldiers' quarters should be required to take quinine regularly and to observe the general rules of hygiene.

Quinine has been much and successfully used as a preventive of malaria. The Italian government has made the most extensive use of it, holding a monopoly on the drug and selling it very cheaply or giving it to the poor. The value of the measure is so well established that there need be no hesitation about prescribing it by order in very malarious regions and in tropical cam-

Protection of Mosquitoes from Infection

Quinine Prophylaxis

paigns. The drug should, for this purpose, be given daily in doses of from two to five grains. Thus administered after meals, it will produce no symptoms and will prevent the disease in many instances.

Experience indicates that the work of mosquito "extermination" is more effectively done if regular forces are kept at it to the exclusion of other work, and every tropical or mosquito-infected post should have such a "mosquito brigade." Men kept on the work soon learn the locations and habits of the insects and become expert in destruction or prevention. When the work is left to householders or to general fatigue parties, it is usually neglected and the plague continues unabated.

**Mos-
quito
Brigades**

FLEA-BORNE DISEASES

Bubonic plague is the most important disease transmitted to man by fleas, and it will be discussed here only briefly, as mention has already been made of it. The disease is due to a small bacillus that is possessed of great vitality, so that the virus may persist for a long time. Extensive investigations in different parts of the world, but principally in India, have clearly demonstrated that the disease is usually transmitted in the manner here indicated, though it may occasionally be communicated in other ways. The relation of rats to plague epidemics has already been discussed. A Russian observer states that bedbugs also may transmit it. The germs are excreted by plague patients in the spit, urine, the pus from ulcers and buboes, and possibly the feces, so that dust-infection and contact may account for some cases. Nevertheless the rat and the rat flea are responsible for the

Plague

majority of cases and probably for all epidemics, and the measures of prevention and control are aimed at these two. The following facts, among others, constitute the reasons and justification for such measures.

Rats abound particularly in houses and neighborhoods which, because of poor policing or other sanitary neglect, afford them food, shelter, and breeding-places. They are very susceptible to plague and are probably the natural hosts of the disease. When severely infected they may have the germs in their blood in enormous numbers, as many as 100,000,000 in a cubic centimetre of blood. All rats have fleas, which remain on them until death, when they leave them and seek other hosts, usually rats, but in their absence men or other animals. Fleas breed in the nests of their hosts, live on their blood, and in biting may readily ingest as much blood as may contain, in the case of plague septicemia, 5,000 plague bacilli. These bacilli may remain alive and virulent in the flea's stomach for as long as fifteen days, and then be ejected when the flea bites again and gain entrance to the wound made by the insect, thus producing infection.

While the rat flea is the insect principally concerned in carrying the disease, dog, cat, and human fleas may act the part. The following is the ordinary life history of the rat flea in Bombay. The eggs are laid at all seasons, from one to five at a time, and hatch in about two days. The larvæ are wormlike and resemble some small fly and weevil larvæ. They live on almost any kind of refuse, and, after varying periods of not less than one week, spin cocoons of fine, silky thread, which become

Life History of Rat Flea

covered with dust and rubbish and are difficult to distinguish. The adult fleas escape from these cocoons after one or two weeks and seek a host, as they can only live on fluid food, especially blood. The most favorable temperature for breeding is from 50° to 85° F., and when the mean is above or below this range, breeding stops or is delayed, a fact which partly accounts for the seasonal variations of epidemics. Each species of flea has its own host, which it prefers to all others, but in the absence of that host, and after a period of starvation, it will seek others. The rat flea, for instance, will not usually bite man until three days after the death of its rat host.

The measures to be taken for the prevention and suppression of epidemics are indicated by the facts recorded above. They must embrace numerous and costly expedients unless the fight is begun in earnest at the first appearance of cases, and the disease eradicated before it has had an opportunity to spread. Failure to take vigorous action early resulted in the spread of plague through San Francisco, so that the work and expense of fighting it have had to be continued for eight years. Similar failure of early action has been followed by the spread of the disease through India to such an extent that from November 24, 1906, to November 9, 1907, there were in that country 1,394,947 cases, of which 1,276,576 were fatal, and the end is not yet in sight.

Prevention and Suppression of Epidemics

General hygiene is of great importance as lessening the numbers of both rats and fleas, both of which abound in the presence of filth. Such measures must include the removal or destruction of rubbish and other

material in which the rats hide or make their nests, the stopping of their holes or other means of access to walls, floors, and other hiding-places, the prevention of their access to garbage and other food-supplies. They must also include such cleanliness of persons, domestic animals, houses, streets, and communities, as will make the presence of fleas, bedbugs, and lice improbable.

**General
Hygiene**

Special measures should also be adopted against rats, such as their destruction by poisoning, trapping, and disease. The last-named measure is employed by scattering bait contaminated with cultures of a bacillus that causes epidemics among rats, but does not harm man or his

**Measures
against
Rats**

domestic animals. The keeping of cats has been advocated as a useful anti-plague measure, and it is probably of value. Old and rat-riddled buildings may need to be destroyed. Good buildings should be made rat-proof by means of concrete, cement, and sheet-iron. Stables and warehouses will require particular attention, and their floors, walls, and roofs should be made as rat-proof as possible. Because of the frequency with which rats are transported in ships, all wharves and docks must be freed from them as far as possible, ships from infected ports should be required to anchor offshore, and all cables or chains leading to shore or to other boats should be provided with shields that rats cannot pass. Sewers should be well flushed to keep them clean, and should be screened or barred so that rats may not pass from them to houses or other buildings. Houses or boats known to be infested should be tightly closed and fumigated with sulphur fumes or other poisonous gas that will destroy the rats.

The most important measures against fleas are probably good general policing, the maintenance of clean floors, and the absence of litter and waste in which they may breed; but other measures are also of value, and among them may be mentioned the use of insect-powder and antiseptic soaps on domestic animals, the use of foot-covering and other adequate clothing, and the use of petroleum on floors and streets, and of putty, paint, and antiseptic solutions in cracks of floors and walls. The use of petroleum is fairly common in the tropics now, as a protection from ants principally, but it will probably be as valuable against fleas. As previously suggested, the use of crude oil in road-making, as is now much practiced in this country, will probably prove of value in lessening the number of fleas. Fleas may be collected in rooms or houses known to be infected by turning guinea pigs loose in them. The fleas attack the animals, and may then be stupefied with chloroform, combed from the hair, and burned.

**Measures
against
Fleas**

Every case of plague in man or rat should be promptly reported, so that proper measures of isolation, fumigation, disinfection, etc., may be instituted early. All rats found dead, as well as those trapped or otherwise captured, should be tagged with a statement of the time and place, and then examined for the presence of plague. If it is found that they have the disease, active measures must be begun at once in the neighborhood from which they came. All persons afflicted should be isolated and their discharges cared for. Disinfection of houses and effects should follow.

**Notifi-
cation of
Cases**

Anti-plague vaccination has been considerably used

in India for some time now, and the following statements as to results have been published:

Vaccination "1. In a native of India, who is more susceptible to the disease than Africans, Europeans and some other races, the inoculation now in force reduces the liability to attack to less than one third of what it is in a non-inoculated Indian. 2. In the one third of cases which still occur, the recovery rate is at least double that in the non-inoculated attacked. The ultimate result is a reduction of plague mortality by some eighty-five per cent. 3. In an inoculated European an attack of plague, if it subsequently occurs, has so far always ended in recovery. 4. The inoculation is applicable to persons already infected and incubating plague, and prevents the appearance of symptoms or else mitigates the attack." These claims are made by the originator of the method, and his opinion may possibly be unduly optimistic.

FLY-BORNE DISEASES

In addition to typhoid and other diseases discussed in preceding parts of this book, in which the fly acts as a simple carrier of infection but does not actually introduce it into the body, the sleeping sickness of Africa is a very important disease. It is introduced by a biting fly, and is caused by a small animal parasite very much like that causing surra in horses in the Philippines and other places. Professor Koch also thinks that the disease may be transmitted by sexual intercourse. The flies concerned are found only in thick underbrush near water, and Koch has also suggested that they obtain the organisms in the first place from crocodiles. The disease is

almost uniformly fatal, is spreading rapidly, and is being thoroughly investigated. The only means of prevention yet known are the avoidance of fly-bites, the cleaning of undergrowth from near streams, and the avoidance of infected localities. It is improbable that our army will ever have any experience with this disease, but other affections due to biting flies may be discovered and the possibility should be kept in mind.

TICK-BORNE DISEASES

The most important tick-borne diseases that have been well studied are the African relapsing fever, due to a spirillum, and the spotted or tick fever of western Montana, due to an unknown organism. The latter disease may affect American troops, as Fort Missoula is in the country in which it occurs. No case has yet appeared in a soldier. Both of these diseases may be transmitted by ticks that have bitten sick persons, or by the young of such ticks. The methods of prevention thus far known to be of value are the avoidance of infected districts and of tick bites. In the case of spotted fever the season is short and the danger is slight, except from the first of March to the middle or end of July. If infected regions must be visited, the use of adequate clothing and of head-nets and bed-nets will be of value, but probably not absolutely protective. Wood-ticks, such as convey spotted fever, disappear as forests are cleared and the land cultivated. The ticks that convey the African fever abound principally in rest-houses and old camps, and cleanliness, avoidance of old huts and old camp-sites, and the use of nets, would seem the best measures of protection.

DISEASES BORNE BY BEDBUGS

As stated above, a number of diseases have been credited to the agency of these pests, and the evidence incriminating them is sufficiently strong to make their avoidance and destruction a necessity, as well as an æsthetic measure. They are so generally recognized as vermin and indicators of uncleanness that their presence constitutes a cause of shame in a well-regulated house or barracks. The avoidance of diseases transmitted by them depends on scrupulous cleanliness of houses, beds, and bedding, and the avoidance of those that are not clean. Soldiers, though living in clean surroundings, may expose themselves to bites in seeking sexual indulgence or drink, and they should be taught by precept and example not to do so.

CHAPTER XV

VENEREAL DISEASES

THE term venereal is usually applied to the diseases resulting from illicit and impure sexual intercourse. The principal members of the group are syphilis, chancroid or soft chancre, and gonorrhœa. In a general sense the adjective is truly descriptive, but, as stated before, it is not always so, and all of these troubles may be contracted by persons absolutely innocent of illicit connections or even of proper intercourse. It is, however, so generally true that in soldiers, at least, the diseases result from improper conduct, that they are assumed to be "not in line of duty," and a report to the contrary requires special explanation.

These diseases are particularly frequent in youth and early manhood, and it is stated that "about 95 per cent of all cases of gonorrhœa occur between the ages of sixteen and thirty."

General Causes

The reasons for this are numerous, and some of them may be summarized as follows:—

1. This is the period at which sexual desire and power are at their maximum. After the age of thirty some decline in both has usually begun.

2. It is also a period in which judgment is not matured, self-control not well developed, and ignorance not dissipated, so that the future wife and family exercise no claims and venereal diseases are regarded as trifling.

3. Manhood and virility are synonymous terms, in

common usage as well as in the dictionaries, and youth delights in demonstrating its manhood.

4. Most men are married by the time they reach the age of thirty, and are less apt to be exposed to infection thereafter. The soldier is probably more apt to contract venereal disease than the young civilian, because his associates are practically all males; his topics of conversation are largely such as are only handled in "stag" gatherings; he is removed from the restraints of the family and of the public opinion that can most influence him, that of his friends and acquaintances at home; he may at times find it difficult to obtain access to other female society than that of prostitutes, and these are always to be found. He may also drink a bit to demonstrate his manliness, to relieve his loneliness, to be companionable, or simply because he is "a young man, void of understanding"; and then, with judgment perverted and desires inflamed by the alcohol, he forgets danger and seeks intercourse where he can most readily obtain it.

These things are enough to account for the fact that venereal diseases are prevalent, but they do not account for the other humiliating fact that our
Prevalence in our Army army shows a greater amount of them than any of the European services. Many explanations of this have been attempted, but none of them seems adequate. One that may partly account for it is that American prudery results in an ignorance of sexual matters and of venereal diseases such as does not obtain elsewhere. According to the last annual report of the Surgeon-General of the Army, venereal disease kept an average of 668.65 men on sick report during the entire fiscal year of 1908.

When to this is added the time lost because of troubles indirectly due to venereal diseases, and the duties poorly performed because of them, it is probably no exaggeration to say that it was almost as though one regiment had been eliminated from the service.

The company officer cannot expect to work a revolution in the morals of his men, or to make chastity a general characteristic of them, but he can exercise some influence in that direction by his personal conduct and example, by choosing non-commissioned officers who will probably do the same, by making the soldier's surroundings pleasant and attractive, and encouraging him to find his amusement and recreation in the post, by providing good reading-matter and helping to make the most of the post exchange as a place of recreation and amusement, and by giving his men instruction as to the dangers of venereal disease and the sanity, healthfulness, and safety of sexual control and continence. Recruiting officers can do much by careful selection of men, as it is a well-known fact that the great number of admissions to sick report for venereal disease is made up by a relatively small number of men, the majority of whom are of the less desirable class of soldiers. Drink, poor home-training, ignorance, and lack of self-respect lead these men to places and practices that their more decent comrades avoid.

**General
Measures
of Pre-
vention**

As man is the only carrier of these diseases, as they are transmitted in nearly all instances by direct physical contact, and as early diagnosis, isolation, and treatment enable us to do much in the control of other transmissible diseases, it seems reasonable to believe that similar measures, if generally enforced, would

produce great results in these. There can hardly be any doubt that if all cases of venereal disease were promptly reported and then isolated and kept from sexual intercourse until well, the diseases would be exterminated. These are such chronic troubles, however, and require such prolonged and careful treatment to insure a cure, that such a plan is quite impracticable. Many efforts have been made to approach it though, by the regular examination of prostitutes and their isolation and treatment until free from contagious lesions, and similar measures have been adopted in regard to soldiers in our own service. But it cannot be said that the diseases have been brought under control by such measures, and the reasons are fairly obvious. It is practically impossible for an examiner to know positively that a given person is free from venereal disease. Chancroids are easy of detection, but both syphilis and gonorrhœa may linger, capable of transmission, without giving any indications of their presence that can be detected by the inspection that the examiner of large numbers can give. This is true in the case of soldiers, and much more so in that of prostitutes, who, moreover, are adept in the removal of slight signs that may be present, and whose interest lies in escaping isolation. Most prostitutes are such for one of two reasons, either for the pleasure and excitement they find in the life or for the money they can make out of it. In either case their object is defeated for the time being if they are isolated. Furthermore, there is always and everywhere a large amount of clandestine prostitution, that is, that practiced by women not openly professing to do so; and as registration of the prostitute is a necessary preliminary of her regular examination,

a large amount of venereal disease thus escapes detection. Naturally, too, the amount of this clandestine prostitution increases with the severity and strictness of the measures of control over that which is licensed. As examinations are not usually made more than once a week, and in many instances not so often, a woman may develop a contagious discharge or sore the day after examination, and transmit disease to fifty men before being examined again. A prostitute living near a Western post stated that she usually had intercourse with from fifteen to twenty-five men on the night following payment of the troops, and with from one to five or six men on other nights.

It must be borne in mind that American sentiment has not favored and possibly never will favor the state control and licensing of prostitution. Without going into a discussion of the right and wrong of the matter, we are brought face to face with the following condition as regards the army. It consists of aggregations of men, mostly young, mostly unmarried, and largely ignorant, who have abundant opportunity to gratify their sexual instincts, but with women who are under little or no control, many of whom are diseased, and all of whom are interested in avoiding treatment that involves isolation. The obvious courses of procedure are, 1st, the provision of wholesome surroundings and amusements to keep the men away from dives and brothels ; 2d, their instruction as to the nature and dangers of the venereal diseases, so that the large amount of these due to ignorance may be eliminated ; 3d, advice as to the best methods of avoiding disease in case they are so persistently foolish as to expose themselves to it. The instruction as to the nature and danger of the

venereal diseases may be derived from the following consideration of them.

Gonorrhœa

Gonorrhœa or "clap" is a disease of such great antiquity that it may be as old as prostitution itself, and the first half of the fifteenth chapter of Leviticus indicates that Moses knew it and also knew that it was conveyed by contact, and that he provided rules to prevent its spread.

The disease is due to a bacterium known as the gonococcus. This organism occurs constantly in the gonorrhœal discharge, and the disease can be produced by artificial inoculation with it. It is very hard to cultivate artificially, and is readily destroyed by drying, heat, and antiseptic drugs. It may live for some time on towels or linen, or in warm water. It grows on mucous membranes, and those of the genital and urinary passages, of the eyes, mouth, and rectum may be affected.

The young are more susceptible to its action than the old, and children are particularly easy of infection.

Age The disease once started in hospitals for children or in orphan asylums, especially among little girls, is apt to become epidemic.

Transmission is usually by sexual contact, but it may occur through mediate transfer, as by towels, soiled hands, penis syringes or urethral sounds. Such methods are uncommon, however, and little credence can be placed in the story of the man who thinks he got it from a water-closet seat. Strain, as from heavy lifting, can never cause the disease, though often alleged to do so.

The incubation period varies considerably, and the disease may manifest itself within a day of the impure connection or it may not do so for ten or twelve days, or for some intervening period. It usually begins with slight burning on urination, which leads the subject to examine his penis, when he may find the margins of the urinary opening red and somewhat puffy. Or the lips of the opening may be stuck together when he awakes in the morning, and a drop of clear discharge, like white of egg, can be expressed from it. The symptoms increase rapidly, and within another day or two the disease is at its height. The discharge is then more profuse, is creamy in color and consistency, the penis is sore, and urination painful. Erections may be frequent and painful, especially when the patient is in bed and when the inflammation has extended out into the body of the penis. These may cause intense pain, and the organ may be curved. This condition is known as chordee. The inflammation may extend to the bladder, causing very frequent and painful urination; to the testicles, causing swelling, pain, and sometimes abscess formation, and often leading to stoppage of the duct leading from the testicle to the seminal bladder, and so rendering the former useless as a procreative organ. Germs may be carried to the lymphatic glands in the groins, and cause them to become very painful, tender, and swollen, and oftentimes to suppurate. This glandular swelling constitutes a bubo.

Incubation

Symptoms

These various troubles may cause great pain and suffering, but, in the absence of further complications, they lessen in severity after a time, usually from one to six weeks, the swellings subside, the discharge dimin-

ishes or ceases, and the patient may think he is well.

**Persist-
ence**

Or he may recognize that he has gleet. This is a condition not attended by other symptoms than the presence of a small, clear drop of discharge in the early morning, and it is usually regarded as of no importance. It is gonorrhœa, however, and is capable of setting up the disease in others, and many unfortunate brides have been so infected. It may persist for months or years, and other manifestations of the disease may do likewise. Many men have suffered or died from the results of gonorrhœa ten, twenty, or more years after they thought themselves cured of it; or, worse still, they have seen their wives or children sicken, become blind, or die as the result of it, and have had no thought that they were themselves responsible for the calamity.

The complications of gonorrhœa are so numerous, so serious, and some of them apparently so remote, that

**Complica-
tions**

there is no space to discuss them here, but some of them will be mentioned. *Stricture* is one of the most common. It is often present when a gleet persists. It may cause obstruction of urine, great suffering, and serious or fatal kidney disease years after the acute gonorrhœa. The principal symptom is obstruction, usually partial, to the urinary stream. *Inflammation of the prostate gland* may develop and last for years, causing pain in urination and perhaps during defecation, pain deep between the legs, marked nervous symptoms, and general invalidism. *Gonorrhœal rheumatism*, or joint inflammation, is always hard to treat, causes great pain, and tends to last for a long time. It may result in great deformity and permanent crippling. *Tendon inflammation* may also occur and produce symp-

toms much like the joint troubles. Pain and tenderness of the heel or of the great tendon leading upward from it may result from gonorrhœa and prevent marching. *Eye diseases* of at least two kinds result from this disease and always endanger and frequently destroy the sight. Gonorrhœa is one of the great causes of blindness, the disease often developing in children at the time of birth, from infection entering their eyes from their mother's genitals, the mothers often being infected by their husbands who thought themselves cured before marriage. The disease may also be conveyed to the eyes by the hands after urinating or wiping away discharge from the penis. It may be transferred from one person to another through the medium of towels, wash-cloths, medicine-droppers, and other things in common use. *Blood-poisoning*, that is, the diffusion of germs through the blood, may result from gonorrhœa and may cause death. *Heart disease* may be a complication, and often accompanies blood-poisoning. It may cause death early or may be apparently recovered from, and the heart may break down from the damage done it only after the lapse of years. Owing to the levity with which the disease is often treated, this complication may not be detected at the time of its occurrence.

As already stated, the subject of gonorrhœa is a source of danger to his companions and his family, and in military life it is desirable that he should be confined to hospital, at least during the time of most active symptoms, and that all men should be cautioned against using the towels or linen belonging to others. No man worthy the name would deliberately infect another person with his disease, but many men unintentionally do so. All should

**Danger
to Others**

be instructed to abstain from sexual intercourse until entirely cured.

Without condoning or attempting to justify illicit intercourse, we must recognize that many men will indulge in it in spite of instruction, warning, and example, and it is the part of economy and makes for the efficiency of the army if those men can be protected from disease in spite of their folly and misconduct. No infallible means of protection from venereal contamination is known except chastity, but the following measures may offer some degree of it. *Urination*, if practicable immediately after intercourse, tends to wash out germs that may have entered the urethra. *Irrigation* of the urethra with a mild antiseptic solution, such as permanganate of potash solution in strength of one part to three thousand, has a similar action. *Moderation* in intercourse is advisable. Repeated intercourse in the space of a few hours brings about a congestion of the genitals which is slow in subsiding and which predisposes to infection. *Condoms*, impermeable rubber coverings for the penis, are sold and used for the prevention of disease, and they should protect against gonorrhœa.

After the disease is once contracted the best way to prevent complications is by rest in bed and proper treatment. Soldiers should, therefore, be taught to report early for treatment, and to continue at it until entirely cured, to be cleanly in their habits, and by frequent hand-washing and care of towels and linen to avoid infecting the eyes, and (very important) to avoid consultation with advertising quacks or the use of advertised medicines, both of which are fraudulent.

Prevention

Preventions of Complications

Chancroid

Chancroid, soft chancre or soft sore, is the least important of the diseases here discussed, as it is a local infection and does not do the general damage or assume the manifold and insidious forms that often characterize gonorrhœa or syphilis.

The typical and usual soft chancre is due to a small bacillus which bears some resemblance to that causing plague, but ulcers may be produced on the penis, as elsewhere, by other organisms.

Cause

The incubation period varies from one to ten days, and its length probably depends on a number of factors, such as the degree of infection in the woman, the presence and size of scratches or abrasions on the penis, and the violence, length and frequency of the infective intercourse.

Incubation

The disease begins as a small ulcer, or sore, which may be no larger than the head of a pin. The favorite location is in the depression or groove just back of the head of the penis, though it may be anywhere on that organ. Many

Symptoms

sores may show at once, or one may appear at first and others develop later. The ulcer enlarges and deepens, the surrounding parts are red, swollen, and tender, and the surface of the sore more so. A thin and usually moderate discharge is present, and is highly infectious, and its presence causes the development of the secondary sores. Under proper treatment the ulcers can usually be healed in a few days or weeks, but if neglected or improperly treated or in inaccessible positions, they may spread and produce considerable loss of tissue, half or more of the head of the penis being some-

times destroyed. The base of the ulcer, if not irritated, is usually soft as compared with the base of the syphilitic chancre, hence the name soft chancre. Occasionally this sore may appear on the bag, on the lower hairy part of the abdomen, or anywhere, in fact, that the virus has entered the skin.

Complications The most frequent complication is bubo, which occurs in about one fourth of all cases. It is very painful, causes lameness, fever, and may require operation.

The methods mentioned as tending to protect from gonorrhœa are equally applicable here, except that **Prevention** irrigation would not be particularly valuable, as the sore usually develops externally rather than in the urethra. Washing the genitals in an antiseptic solution, such as one to one thousand bichloride, will offer a certain amount of protection. Early and proper treatment is required to produce speedy healing and lessen the tendency to the formation of buboes, and soldiers should report promptly for it.

Syphilis

There has been much speculation and discussion as to whether syphilis, or pock, originated in the old world or the new, whether the Spaniards brought it to America, or took it from here. At any rate, the disease spread throughout Europe in the century following Columbus's voyages, and it has prevailed throughout the world until the present time, and is now seen in all ranks of society and all walks of life. It is the most dreadful of the diseases considered in this chapter because of its chronicity, the long time required

for its treatment, the great variety of its manifestations, the number and seriousness of the conditions resulting from it, and the fact that it is transmissible by heredity. The disease cannot be described or adequately discussed in the space allotted here, but a very brief outline of it will be given, in order to enable the line officer to impart some degree of information about it to his men.

Although the disease has been known and industriously studied for centuries, it is only lately that its cause has been known. This is the very fine and delicate spiral organism mentioned in a preceding chapter as a *treponema*.

Cause

The commonest method of transmission of syphilis is through sexual intercourse, and this is especially true as regards soldiers, to whom many of the other avenues of infection are not open. But there are such other avenues and they are numerous; and innocent victims of this disease are therefore more numerous than those of gonorrhœa or chancroid. Among such methods we may mention the following:—

**Trans-
mission**

Heredity. The disease may be inherited from either parent, and may infect the child in the womb and cause its death there, cause it to be born diseased, or to be born apparently well and develop the disease soon after. Or, in rare cases, it may be born apparently well and continue fairly so for years, and only show the disease in recognized form when it reaches the age of puberty.

Contact, especially such as kissing, is a fairly common method of transmission. Ulcers or sores in the mouth are a common manifestation of the disease, and in such cases the organisms are present and sometimes numer-

ous there. The saliva or discharge from such a mouth is highly infectious, and, if brought in contact with a slight scratch or sore, will give rise to the disease. Children and girls have many times been infected by kissing persons so affected, and syphilitic children may by this means infect the nipples of their wet nurses. For this reason, also, the spoons, cups, and other eating utensils and pipes or cigarette-butts that have been used by syphilitics, are dangerous. The blood and discharges from sores on syphilitic persons are infectious, and nurses and doctors are occasionally infected in dressing or operating upon such cases. Several instances of the kind have occurred in the army. Barbers may receive infection if they get wounds or scratches contaminated by the saliva or blood of their patrons, and they may also transmit it through scratches or cuts. Dentists and their implements, if not clean, may transmit it.

The incubation period of syphilis is long. It is never less than ten days, and it is said that it may be as long as ninety. As a rule, however, it is from three weeks to one month, after which time the primary sore or chancre develops at the point of entrance of the virus. This fact may be of value in determining the manner of infection. If the chancre be on the genitals, it is usually a result of sexual intercourse; if elsewhere, the case may be different. It may be on any part of the body, as on the top of the head or the ear, where it has been seen to result from bites received in fighting; inside the rectum as a result of unnatural practices; on the breast of a wet nurse; or on the arm of a child vaccinated with virus from a syphilitic.

Incubation

The chancre constitutes the first sign of the disease. It appears first as a small firm lump resembling a pimple. This may enlarge considerably or not at all, depending somewhat on its situation. It usually ulcerates slightly at its top, and

**Primary
Stage**

shows a slight discharge, which may be thin and clear, or purulent. The chancre is nearly always single, and secondary ones rarely develop. Its base is usually hard and firm and feels like cartilage or even like bone, hence this is called the hard chancre. It may be so small as not to be noticed, and, as the patient may continue feeling well while he has it, he may not know that he is diseased until the appearance of secondary symptoms after another period of six to twelve weeks. It is not rare to have soldiers appear with well-

**Second-
ary
Stage**

marked secondary signs of the disease and no chancre, and denying that they have had any. The secondary symptoms are usually plainly marked and lead the patient to seek treatment. They are referred particularly to the skin and mucous membranes, and the most common of them are skin eruptions, sores in the mouth and throat and about the anus, falling of hair, slight fever, pains in the bones or joints, anæmia, and pallor, and various eye troubles. The skin lesions of syphilis are so numerous and of such varied character that it has been said that they can simulate all other skin diseases. The arteries, heart, and internal organs may also be affected in the secondary stage. In fact, no other known disease manifests itself in so many ways as this one, and it simulates the greatest variety of affections, including smallpox and other eruptive diseases, malaria, liver abscess, tuberculosis of lungs, bones, or joints, epilepsy, and very many more. It is,

therefore, obviously out of the question even to mention most of the symptoms here.

This stage may not manifest itself if the case is well treated in the primary and secondary stages. Otherwise it may merge with the secondary, or it may appear only after the lapse of years.

Tertiary Stage

It is characterized by deeper skin lesions, that show a tendency to ulcerate and form large and deep sores, by destruction of bone, the formation of swellings or tumors, and the degeneration of internal organs. Tertiary syphilis is more often mistaken for tuberculosis, cancer, and other chronic ailments, than for more acute ones. The syphilitic tumors may form in any part of the body, and those in the brain, heart, spinal cord, and other vital parts may be rapidly fatal. Syphilis is the most common cause of aneurism.

The disease is usually spoken of as consisting of three stages, but some writers consider a group of severe and incurable nervous troubles, especially locomotor ataxia and general paralysis of the insane, to constitute a so-called

Fourth Stage

fourth stage. Whether or not this is true cannot be positively stated; but syphilis undoubtedly exercises a strong predisposing influence in regard to those diseases.

Syphilis is transmissible in any of its stages; highly so in the first, more so in the second, and but slightly in the third. The time of greatest danger is that during which there are sores, with open moist surfaces, in the mouth, the nose, on the genitals, and about the anus; and it is advisable, in military service, to confine in hospital men who present such lesions. At other

Period of Contagiousness

times if they are kept under observation by a medical officer and continue their treatment, they may safely mingle with their fellows and do full duty.

There is some difference of opinion as to the length of time required to effect a cure of syphilis, some authorities giving two years as the minimum period, and others saying that at least four are required. It is probable that two years of faithful and proper treatment will suffice to cure the majority of cases of syphilis, but a small minority require all of four years or more. This cannot be too strongly impressed on all syphilitics, and their own welfare and that of their families demands that they should take proper and continued treatment under the supervision of a competent physician for at least two years, and longer if the physician advises it. No syphilitic should marry until under observation and treatment for at least two years *from the last appearance of symptoms*.

**Length
of Treat-
ment**

The special preventive measures that may be employed by men exposing themselves to the dangers of polluted intercourse are the same as were mentioned for gonorrhœa and chancre. A small amount of experimental and clinical evidence also indicates that protection may be conferred, in at least some instances, by the use of a calomel ointment. This is thirty per cent calomel in lanolin, and should be well rubbed into the penis and exposed parts for from five to twenty minutes, at the conclusion of the intercourse and after the use of the irrigation and wash mentioned as preventive of gonorrhœa and chancre.

**Preven-
tion**

Mixed Infections

One may be simultaneously infected with two or all of the venereal diseases, and in such instances the more quickly developing gonorrhœa or chancreoid masks the presence of the chancre, and great surprise may result when the secondary symptoms of syphilis become manifest. Occasionally the chancre develops inside the urethra and occasions a slight discharge which may be mistaken for gonorrhœa, or the latter disease may co-exist. In either event the chancre may be overlooked. More frequently, however, the confusion arises from the coexistence of soft and hard chancres. It is generally believed by men having venereal sores that the soft sore is never syphilitic, and that it always arises within ten days, while the hard chancre is always syphilitic, and never appears within ten days. These are usually true as related to pure single infections. But the viruses of soft chancre and of syphilis may both enter the skin at the same point, in which case a soft and ulcerating sore may develop early and be followed by a large suppurating bubo, a thing not occurring in pure syphilis, and may heal under local treatment, leaving no induration, or only a slight one that is ascribed to inflammation. The syphilitic virus nevertheless continues its work and in due time the secondary symptoms develop. Some soft sores, especially those burned with acids, have a base of inflammatory tissue that may greatly resemble the hard base of the syphilitic sore.

So difficult is the differentiation of syphilitic from non-syphilitic venereal sores, by the appearance alone, that most careful practitioners do not definitely decide that a given sore is syphilitic, and therefore do not

begin constitutional treatment until secondary symptoms occur. Laboratory methods now permit of positive early diagnosis in a considerable number of cases.

In some cases of syphilis the secondary stage is very lightly marked, and may be overlooked, so that men are occasionally seen suffering with tertiary lesions, and stoutly and in good faith denying that they ever had either a hard chancre or secondary symptoms.



INDEX

ABDOMEN, the, 6.
 Abstinence, total, encouragement of, 28.
 Accidents, condition of mouth in, 25.
 Acid, boric, as food preservative, 39.
 Acid, carbolic, as antiseptic, 210; as vermin killer, 72; use of, in flush closets, 63; for camp urinals, 98
 Acids, organic, food-value of, 35, 122.
 Actinomyces, transmission of, 37.
 Age, as a factor in efficiency of the recruit, 4; diseases and, 143.
 Age limit of recruits, 3.
 "Ague-fit," 171.
 Air, arctic, 131; as predisposing factor in disease, 149; effect of, on epidemics, 242; fresh, value of, 20; moist, supply of, in rooms, 69.
 Air-space, quantity of, per man, 56.
 Albuminuria, due to faulty diet, 34.
 Alcohol, 28, 31, 151; on marches, 106; use of, in arctics, 133; use of, in tropics, 123.
 Alcoholism, indicated by condition of nose, 9.
 Alkalies, as cause of disease, 160.
 Ambulance kitchens, 110.
 Amœba, as cause of liver abscess, 172.
 Anemia, influence of neglected mouth on, 25.
 Animal poisons, 162.
 Animals, disinfection of, 254.
 Anopheles, carriers of malaria, 72, 260, 265.
 Anthrax, transmission of, 37, 191.
 Antiseptics, 210; action of, 178.
 Antitoxin, as specific for diphtheria, 212, 255; as specific for tetanus, 212.
 Anus, cause of irritation of, 64.
 Appearances, as guide to judgment of characters, 18.

Appetite, effect of cooking on the, 39.
 Aprons, liberal use of, by company barber, 61.
 Arch of foot, deformities of, 15.
 Arms, of recruit, 12.
 Assimilation, increase of, by exercise, 21.
 Asthma and barrel-shaped chest, 6.
 Athletes, effect of oxygen on, 104.
 Athletics, encouragement of, 21.
 Atmosphere, state of, as affecting epidemic, 241.
 Auger, post-hole, for digging latrines, 96.
 Auto-intoxication, 161.
 Bacilli, definition, 176.
 Bacillus, anthrax, resistance of spores of, 176.
 Bacillus prodigiosus, harm of, in erysipelas, 179.
 Bacteria, 174; action of sunlight on, 178; aerobic, 176; anaerobic, 176; dried, harm of, 178; mutual destruction of, 178; reproduction of, 176; transmission of, 37.
 Baking-ovens, field, 87.
 Baldness of recruit, 12.
 Barber, company, 61.
 "Barrack odor," avoidance of, 21.
 Barracks, hygiene of, 54.
 Barrels, as night urinals, 98.
 Barrels, use of for drinking water, 81.
 Bathing, by roster if necessary, 62; in causes of disease, 152; in camp, 83.
 Bath-rooms, 62.
 Baths, 23, 127.
 Bath-tubs, 67.
 Batter-cakes, 41.
 Battlefield, hygiene of, 115.
 Bay rum, use of by drunkards, 29.
 Beans, 24; cooking of, in field, 87.

- Bedbugs, 22, 117, 252, 280; as disease-carriers, 193.
- Bedding, airing of, 23; improvised, for camps, 78.
- Bed-nets, necessity of, 57, 271.
- Beef, standard component of meat ration, 35.
- Beets, 42.
- Belly-band, 124.
- Belly, first aid to wounds of, 116.
- Beverages, 47.
- Bichloride, for camp urinals, 98.
- Birds, as carriers of tuberculosis, 190.
- Bitter Root Valley, fever in, 74.
- Bivouac, bedding for, 78; menace of, to inhabitants, 77.
- Bill of fare, weekly, 48-50.
- Binder, abdominal, 124.
- Biscuits, baking of, by company cooks, 40.
- Bladder, cause of irritation of, 107.
- Blankets, frequent airing of, in camps, 77; when to be carried, 111.
- Bleeders, 143.
- Blindness, due to wood alcohol, 29.
- Blood, in causes of epidemics, 244; in feces, surgeon to be consulted, 64; vomiting of, cause for rejecting recruit, 8.
- Blood-poisoning, gonorrhœal, 289.
- Blouses, impediment of, in tropics, 111.
- Blow-flies, 169.
- Board, anti-typhoid, 237.
- Board floors, provision of, in camp, 78.
- Body, types of, 143.
- Body-vermin, spread of, in camps, 78.
- Boils, preventive measures for, 128.
- Bone, deformities of, 6.
- Books, as disease-carriers, 194.
- Bottoms, river, diseases of, 74.
- Bowels, regulation of, 24.
- Bow-legs, 13.
- Bowls, lavatory, as receptacle for improper objects, 65, 67.
- Bread, 40.
- Breeding-places, mosquito, destruction of, 266.
- Breast-bone, depression of, 6; prominence of, 6.
- Bright's disease, relation of weights to deaths from, 5.
- Buboes, 16, 287, 292; cases of, on marches, 105.
- Buildings, 54.
- Bunion, 14.
- Butter, 45; as carrier of tuberculosis, 189.
- Buttermilk, 45.
- Cabins, as substitutes for tents, 79.
- Caisson-disease, 158.
- Cake, 40.
- Camp craft, development of, 87.
- Camp police, 99.
- Camps, and camp life, 73.
- Camp-sites, as factor in causes of disease, 154; in prevention of epidemics, 233.
- Cancer, as effect of tobacco, 30; relation of weights to deaths from, 5.
- Canning, comparative value of, for food preservation, 39.
- Cans, as night urinals, 98; garbage, care of, 52, 61, 88; tin, disposal of, in camps 89; use of, for cooling drinking water, 81.
- Carbohydrates, 34, 122.
- Carbon, monoxide, given off by cast-iron stoves, 69.
- Carpets, as disease-carriers, 194.
- Carriers, chronic, of typhoid, 216.
- Carriers, disease, animal, 184; human, 180; inanimate, 193; insect, 192; of anthrax, 191; of cerebrospinal meningitis, 181; of dysentery, 182, 213; of glanders or farcy, 186; of gonorrhœa, 183; of gonorrhœa of eye, 182; of hydrophobia, 184; of impetigo, 182; of intestinal worms, 182; of itch, 182; of plague, 187; of pneumonia, 181; of ring-worm, 182; of scarlet fever, 181; of smallpox, 183; of syphilis, 183; of tonsillitis, 182; of trachoma, 182; of tuberculosis, 182, 189; of typhoid fever, 182, 213.
- Carriers, of water-borne diseases, 213.
- Casualties, relation of weights to deaths from, 5.
- Caterpillars, as cause of disease, 158.
- Cats, as carriers of diphtheria, 191;

- as carriers of hydrophobia, 184; as carriers of tuberculosis, 190; exclusion of, from camps, 78; not to be allowed in kitchen, 61.
- Cattle, as carriers of tuberculosis, 189.
- Cavalry, most desirable recruits as to size for, 4.
- Cellars, damp, avoidance of, 55.
- Cellular tissue, diseases of, relation of weights to deaths from, 5.
- Cellulose, 34.
- Cereals, underdone, as cause of indigestion, 87.
- Cerebral affections, relation of weights to deaths from, 5.
- Cerebro-spinal meningitis, transmission of, 181.
- Chafing, preventive measure for, 128.
- Chancroid, 291.
- Character, of recruits, 18.
- Cheerfulness, as stimulant, 106.
- Cheese, as carrier of tuberculosis, 189.
- Cheese, Roquefort, moulds of, 174.
- Cheeses, 45.
- Chemicals, as cause of death or disease, 158; as meat preservative, 39; bactericidal action of, 178.
- Chest, asymmetry of, 6; barrel-shaped, 6; certain types of, indicative of disease, 6; freedom of, on marches, 104; long, narrow, predisposing to consumption, 6; measurement and weight, publication of proportions of, 4; mobility of, 5, 6.
- Chicken-pox, incubation period of, 247.
- Chiggers, 170.
- Childhood, diseases of, 240.
- Chlorine, as disinfectant, 208.
- Cholera, isolation of, 229; prevalence of, 121; prevention of, in camp, 80, 108; toxin in, 179; transmission of, 127, 182, 193, 213; vaccination against, 236.
- Chordee, 287.
- Cigarette-smoking, 30.
- Cigar-smoking, 30.
- Circulatory system, relation of weights to deaths from diseases of the, 5.
- Cirrhosis, of liver, relation of weights to deaths from, 5.
- Cleanliness, a primary rule for cooks, 51, 61; as preventative of epidemics, 232, 250; habits as to, revealed by linen worn, 17; in tropics, 129; personal, 22.
- Climate, demands of as to clothing, 30, 152; influencing marches, 101.
- Climates, 120; arctic, 131; cold, diet in, 34.
- Clothing, adjustment of, to variable needs, 30, 111, 124, 152; allowance in field kit, 112; as block to actinism, 125; changing of, 23; dirty, to be excluded from squad-rooms, 58; disinfection of, 254; facilities for washing in camp, 86; for arctics, 134; in causes of disease, 152, 193; quality of, as issued, 24; vermin-infested, 72.
- Clubfoot, 13.
- Coast artillery corps, most desirable recruits as to size for, 4.
- Coats, impediment in tropics, 111.
- Cocci, definition of, 176.
- Cocoa, 47.
- Coffee, 47; drinking of, on the march, 109.
- Cold, as meat preservative, 38.
- "Colds," cause of, 20.
- Cold storage, use of, to preserve foods, 46.
- Colic, gallstone, as cause of rejection of recruit, 8.
- Comma-bacillus, 214.
- Company mess, influence of, 24, 50.
- Components, substitutes of meat ration, 35.
- Condiments, food-value of, 35; use of, in tropics, 123.
- Constipation, causes of, 45.
- Consumption, and type of chest, 6.
- Contact, as cause of epidemic disease, 224, 243.
- Containers, food, handling of, against vermin, 88.
- Contamination, fecal, in field, 88.
- Continence, sexual, 27.
- Cook, as model of cleanliness, 61, 68.
- Cookers, fireless, 87, 110.
- Cooking, aspect of, in field, 86; field facilities for, 87; on transports, 117.

- Cooks, company, common faults of, 39; importance of, 50.
- Corn, 42.
- Cornbread, 41.
- Corns, 14.
- Cots, for permanent camps, 78.
- Cough, in recruit, 10; tobacco, 29.
- Crab-lice, presence of, 16.
- Cream, as carrier of tuberculosis, 189.
- Cremation, as preventative against epidemics, 233.
- Crematory, improvised, 91.
- Creolin, as germicide, 211.
- Crowding, influence of, on epidemics, 241.
- Culex fatigans*, carrier of dengue and filariasis, 264.
- Cuspidors, for day-rooms, 59.
- Darnall's filter, value of, in camp, 81.
- Day-rooms, 58.
- Day urinal, for camps, 98.
- Deaths, from typhoid, 219.
- Deer, as carriers of anthrax, 191.
- Defecation, danger in, 26.
- Defects, if noted by recruit's C. O. to be referred to medical officer, 19.
- Dengue, 173, 264; isolation of, 205; protection against, 111, 129.
- Dentists, scarcity of, 25; to be promptly consulted, 26.
- Deodorants, use of, 64.
- Deodorization, 208.
- Desserts, 43.
- Dhobie itch, preventive measures for, 128.
- Diabetes, relation of weights to deaths from, 5.
- Diagnosis, early, in control of epidemics, 252; in prevention of epidemics, 228.
- Diarrhœa, camp, 76, 110, 213; due to faulty diet, 34.
- Digestion, increased by exercise, 21.
- Digestive system, diseases of the, relation of weights to deaths from, 5.
- Dining-room, 59.
- Diphtheria, 181; antitoxin in, 255; incubation period of, 247; toxin in, 179; transmission of, 26, 44, 191, 193, 251.
- Diplococci, 176.
- Discharges, disinfection of, 254.
- Disease, carried by water-supply, 26, 213; cause of, by leaky plumbing, 67; *echinococcus*, 132; effect of mental state on, 145; epidemic, compulsory registration of, 230, 277; epidemic, investigation of cases of, 231; epidemic, tracing of, 253; general defense against, 199; of Japanese river-bottoms, 75; prevention of, by medical department, 73; *schistosomum*, 152, 162; transmission of, by toilet articles, 22; vermin as carriers of, 22; wool-sorters', 191.
- Disease-carriers, 180, 213. *See, also*, Carriers, disease.
- Disease-centres, selection of campsites in, 76.
- Disease-transmission, prevention of, 180.
- Diseases, air-borne, 239, 240; borne by bedbugs, 280; "catching," isolation of, 205; causes of, 139; contagious, 239; debilitating, recurrence of, 105; exciting causes of, 156; flea-borne, 273; fly-borne, 278; formites-borne, 239; general, indication of, 10; infectious, in company cook, 51; influence of condition of mouth on, 25; inherited, 141; insect-borne, 76, 129, 257; intestinal, prevention of, 87; intestinal, due to vegetables, 122; mosquito-borne, 257; skin, in men or company barber, 62; table of death rates of, as influenced by weight, 4, 5; tick-borne, 279; tropical, nature of, 125; venereal, 281; venereal, during marches, 105; venereal, measures against, 284; water-borne, 213.
- Dishes, as carriers of water-borne diseases, 222.
- Disinfection, 206, 254; against epidemics, 232, 254.
- Dogs, as carriers of diphtheria, 191; as carriers of hydrophobia, 184; as carriers of tuberculosis, 190; exclusion of, from camps, 78; not to be allowed in kitchen, 61.
- Doors, storm, 57.

- Drain openings, clogging of, to be prevented, 62.
- Drainage, town, danger of, to water-supply, 83.
- Dress, discretion in, 30.
- Drills, as aid to change of air in lungs, 20.
- Drugs, as food preservatives, 39.
- Dry-earth system, in disposal of excreta, 97.
- Drying, for preserving of food, 46.
- Dugouts, as substitutes for tents, 79.
- Dust, as disease-carrier, 222.
- Dysentery, amœbic, 172; causes of, 214; isolation of, 229; occurrence of, 76; prevalence of, 121; prevention of, in camp, 80, 108; recurrence of, on marches, 105; transmission of, 182, 193, 213; use of alcohol in, 124; vaccination against, 236.
- Dyspepsia, causes of, 34, 47.
- Ears, examination of, 9.
- Eating, as cause of exhaustion, 107; leisurely, 60; proper manner of, to be taught to soldier, 24.
- Echinococcus, 132.
- Economy, in use of water, 65; measures of, in company mess, 41.
- Effects, personal, disinfection of, 254.
- Eggs, 36.
- Electricity, as cause of disease, 158.
- Emissions, seminal, 27.
- Emphysema, and barrel-shaped chest, 6.
- Emunctories, activity of, 21.
- Endurance, increase of, by oxygen, 104.
- Engineering, works of, against epidemics, 206.
- Environment, affecting disease, 148.
- Epidemics, control and prevention of, 197, 250; danger of, 73; general measures against, 232, 250; methods of transmission, 241; origin of, 215.
- Epilepsy, indications of, 11.
- Epizootic, transmission of, 188.
- Erysipelas, bacterial action in, 179.
- Excitement, sexual, periods of, 27.
- Excreta, disposal of, on the march, 114.
- Exercise, 21; in tropics, 126.
- Exhaustion, due to water, 107; pre-disposed by alcohol, 106.
- Exposure, effect of, on disease, 147.
- Extracts, flavoring, use of, by drinkers, 29.
- Eyes, affections of, transmitted by toilet articles, 22; examination of, 8; gonorrhœal infection of, 289.
- Farcy, transmission of, 186.
- Fare, bills of, advance preparation of, 40.
- Fatigue parties, under medical officer, 99.
- Fats, 34, 122.
- Feces, disposal of, to prevent epidemics, 233; habitual inspection of, 64; human, as field manure, 236; human, disposal of, in camp, 91.
- Feet, care of, 23, 31, 112; of recruit, 12; soiled, as carriers of epidemics, 226; stinking, 14; sweating, 14; weak, men with, excluded from marches, 105.
- Fever, Malta, transmission of, 44; relapsing, transmission of, 75, 193, 279; scarlet, incubation period of, 247; scarlet, isolation of, 205; scarlet, transmission of, 44, 181, 191, 193; Texas, 193; tick, of Montana, 74, 193, 279; yellow, 173, 192, 260; yellow, protection against, 111.
- Filaria, 168, 192.
- Filariasis, 264.
- Filter, Darnall's, use of, in camp, 81.
- Filters, Berkfeld, 81.
- Fingers, deformities of, 12; loss of, 12.
- Fire buckets, as mosquito breeders, 71.
- Fire, kitchen, as crematory, 89; protective measures against, 71.
- Fires, open, as aid to ventilation, 68.
- First-aid packets, 116.
- Fish, 36; as carriers of tuberculosis, 190; use of, in exterminating mosquitoes, 268.
- Flannels, use of, in tropics, 125.

- Flat foot, 15.
 Flavors, food-value of, 35.
 Flea, of rat, history of, 274.
 Flea-bites, as cause of plague, 126, 188, 192, 273.
 Fleas, as carriers of typhus, 252; as disease-carriers, 193; in camp, 79; measures against, to prevent plague, 277; transmission of, by domestic animals, 191.
 Flies, as carriers of general diseases, 193; as cause of epidemics, 223; as transmitters of sleeping sickness, 75, 173; breeding of, in waste, 91, 224; prevention of, in camp, 79, 88; prevention of, in excreta trenches, 94; protection from, 59; protection from, in kitchen, 52; protection from, to prevent epidemics, 233.
 Floor-room, minimum quantity for each man, 56.
 Flukes, 162.
 Fomites, 227, 242.
 Food, and its supply during epidemics, 251; as carrier of disease, 193; cooked, value of, in field, 87; cooking of, during epidemics, 236; exclusion of, from tents, 77; facilities for storing in field, 88; in arctics, 133; in hot countries, 121; on transports, 117; supply of, on the march, 109.
 Food-receptacles, washing of, in camps, 83.
 Foods, etc., 33, 151; absence of, from squad-rooms, 58; canned, 46; odorous, not to be kept in ice chest, 52; raw, washing of, before eating, 83, 236.
 Forbes-Waterhouse sterilizer, 81.
 Formaldehyde, as disinfectant, 208.
 Formalin, as disinfectant, 209, 211; for camp urinals, 98; as food preservative, 39.
 Fowl, 36.
 Fox, as carrier of hydrophobia, 184.
 Freezing, 131, 136; predisposed by alcohol, 106.
 Frost-bite, 131, 136.
 Fruits, dried, 47; unripe, interdiction of, 87; use of, in tropics, 122.
 Fumigation, 208.
 Furnaces, hot-air, 69.
 Gallstone, colic due to, as cause for rejection of recruit, 8.
 Games, encouragement of, 21.
 Gangrene, local, influenced by neglected mouth, 25.
 Garbage, disposal of, in camp, 89.
 Garbage-cans, care of, 52, 61, 88.
 Gases, as cause of death or disease, 158.
 Genitals, of recruit, 15.
 Genito-urinary system, diseases of the; relation of weights to deaths from, 5.
 Germicides, 210.
 Germs, disease-producing, conveyed by meats, 37; non-destruction of, by cold, 38.
 Glanders, transmission of, 186.
 Goats, as carriers of tuberculosis, 190.
 Goggles, use of, in arctics, 137.
 Gonorrhœa, 286; complications of, 288; of eye, transmission of, 182; prevention of, 290; transmission of, 183, 286.
 Gophers, as plague-carriers, 189.
 Gravel, as foundation for camps, 78.
 Grits, occasional serving of, 42.
 Gum trouble, prompt report of, to medical officer, 26.
 Gums, 10.
 Gymnastics, as aid to change of air in lungs, 20.
 Habits, as factor in disease, 146; flexibility of, after certain age, 4.
 Ham, mouldy, rejection of, 174.
 Ham, raw, danger in eating of, 39.
 Ham-bones, use of, in soups, 43.
 Hammer-toe, 14.
 Hands, care of, 23; cleanliness of, at meals, 60; washing of, after visit to toilet, 64.
 Hard-bread, 41.
 Hat, campaign, 113, 125.
 Head, of recruit, 8.
 Head-gear, 113.
 Head-nets, use of, against mosquitoes, 271.
 Health inspections, 230.
 Health, requisites for possession of, 203.

- Hearing, testing of, 8.
 Heart, irregularity of, due to tobacco, 29; organic diseases of the, relation of weights to deaths from, 5; strength of, increased by exercise, 21.
 Heart-disease, due to gonorrhœa, 289.
 Hearts, weak, exclusion from marches of sufferers from, 105.
 Heat, action of, on germs, 207; prickly, prevention of, 128.
 Heat-stroke, effect of alcohol on, 106, 124; predisposed by want of water, 107.
 Heating, 57, 59, 68.
 Hemorrhoids, 7, 8.
 Heredity, 141.
 Hernia, indication of, 7; predisposition to, 7; sites of, 7.
 Hoarseness, in recruit, 10; tobacco, 29.
 Hominy, cooking of, in field, 87; occasional serving of, 42.
 Hood, over kitchen range, 61.
 Hookworm, 126, 165.
 Hope, as stimulant, 106.
 Horseback riding, hindrance of, by venereal cases, 105.
 Horses, as carriers of anthrax, 191; as disease-carriers, 186.
 Hospital treatment, advantages of, 184.
 Hot-water heating, 70.
 Houses, disinfection of, 254.
 Huts, as tent substitutes, 79.
 Hydrocele, 16.
 Hydrogen, peroxide, 210.
 Hydrophobia, transmission of, 184.
 Hygiene, golden rules of, 31.
 Ice, as disease-carrier, 235.
 Ice-chest, special scrutiny of, 52.
 Immunity, 201.
 Immunity, induced, 255.
 Impetigo, transmission of, 182.
 Implements, as carriers of water-borne diseases, 222.
 Incineration, human waste, 92.
 Incubation periods, 205, 246.
 Indigestion, causes of, 87; indication of, by condition of nose, 9; influence on, by neglected mouth, 25; severe, cause for rejecting recruit, 8.
 Infection, intestinal, prevention of, 86; mixed, of venereal, 298.
 Infections, of camps, 76; predisposed by alcohol, 106; preventions against, 126; transmission of, 26.
 Influenza, incubation period of, 247; relation of weights to deaths from, 5.
 Injuries, condition of mouth in, 25; predisposed by alcohol, 106.
 Injury, as affecting disease, 147.
 Inoculation, as preventative of epidemics, 245.
 Insect-borne diseases, 257.
 Insect-larvæ, in cause of disease, 169.
 Insects, disease-carriers, 76, 192; poisonous, protection from, 111, 126, 129.
 Insomnia, due to tobacco, 29.
 Inspection, daily, of field kitchen, 89; house-to-house of civilians, 99, 230; of person, 58.
 Inspections, Saturdays', 58.
 Isolation, of diseased, 205, 229, 253.
 Itch, 22, 170; transmission of, 182.
 Japan, disease of, in certain river-bottoms, 75.
 "Jepson" urinal, 95.
 Jiggers, 170.
 Juice, gastric, bactericidal action of, 200.
 Kerosene, use of, in urinals, 64.
 Kitchen, hygiene of, 50, 60.
 Kitchens, ambulance, 110; camp, 86.
 Kitchen-waste, disposal of, 89.
 Knock-knee, 13.
 Lanterns, to mark night urinals, 98.
 Larvæ, mosquito, destruction of, 268.
 Latrines, water-carriage, 92, 96.
 Laundry-work, facilities for, in barracks, 65; facilities for, in camp, 86.
 Lavatory, location of, 57.
 Lead-poisoning, avoided by cleanli-

- ness, 23; influenced by neglected mouth, 25.
- Leeches, 170; protection from, 126.
- Legs, of recruit, 12.
- Leprosy, indication of, 14.
- Lice, 22, 170 192; as disease-carriers, 193.
- Lighting, interior, 56, 59, 67.
- Lime, chloride, use of, 63.
- Line, coöperation of, with medical department, 73.
- Lime-washing, in camp, 90.
- Liquids, as cause of injury or death, 159; avoidance of large quantities with meals, 24.
- Liver abscess, cause of, 172.
- Liver, cirrhosis of, relation of weights to deaths from, 5; condition of, in tropics, 123.
- Liver disease, indication of, 7.
- Lockjaw, cause of, 176.
- Lumpy-jaw, transmission of cause of, 37.
- Lungs, capacity of, increased by exercise, 21.
- Lymphatics, of recruit, 11, 16.
- Macaroni, 42.
- Maggots, 169.
- Malaria, 170, 257; protection against, 111, 129; recurrence of, on marches, 105; recurrence of, through alcohol, 278; relation of weights to deaths from, 5; transmission of, 183, 192, 193.
- Malnutrition, influenced by neglected mouth, 25.
- Manhood, misconception as to loss of, 27.
- Manners, table, 59.
- Manœuvre-camps, as guides to camp craft, 87.
- Marches, 101.
- Marriage, of syphilitics, 297.
- Marshes, vicinity of, as camp-sites, 76.
- Mastication, thorough, 60.
- Matches, as cause of trouble in plumbing, 67.
- Material, building, choice of, 56.
- McCall incinerator, 92, 97.
- Meals, hot, supply of, on the march, 110; time allowance for, 21.
- Measles, incubation period of, 247; isolation of, 205; transmission of, 183, 191.
- Measures, sanitary, duty as to, not ended by care of self, 101.
- Meat, tuberculous, sale of, 189.
- Meats, 35; fried, faulty digestion as result of, 39.
- Medicines, specific, 211.
- Melancholia, cause of, 129.
- Membranes, mucous, germ-expelling function of, 200.
- Men, condition of the immature, 3.
- Meningitis, cerebro-spinal, 247.
- Mercury, as specific for syphilis, 212; bichloride, as antiseptic, 210.
- Mess, field service, 86; lack of variety in, cause of, 40.
- Mess-kits, washing of, in camp, 83.
- Mess-management, books on, 53.
- Mess-room, 59.
- Mice, as carriers of anthrax, 191; as plague-carriers, 130; prevention of, in camp, 79.
- Micro-organisms, effects of, in "close" rooms, 20; presence of, in tents, 77.
- Milk, 43; as carrier of disease, 193, 220, 246; as carrier of scarlet fever and diphtheria, 193; as carrier of tuberculosis, 189; as cause of typhoid epidemic, 220.
- Milk cans, special scrutiny of, 52.
- Milk-sickness, transmission of cause of, 37, 44.
- Mind, condition of, predisposing to disease, 145.
- Mites, as disease-carriers, 193.
- Moisture, aid of, to bacteria, 177.
- Mosquito, dengue-infected, 264; malarial, 260; yellow fever, 263.
- Mosquito-borne diseases, 257.
- Mosquito brigades, 273.
- Mosquito nets, use of, in tropics, 111.
- Mosquito-netting, use of, in field kitchen, 89.
- Mosquitoes, as carriers of malaria, 170, 193; breeding of, in fire buckets, 71; measures against, 265; protection from, 126, 130, 265; protection of, from infection, 272.
- Moths, as cause of disease, 158.
- Moulds, as destroyers of mosqui-

- toes, 268; as disease-producing parasites, 174.
- Mouth, care of, 25; of recruit, 9.
- Mucus, large amounts of, in feces, 64.
- Muffins, baking of, by company cooks, 40.
- Mules, as disease-carriers, 186.
- Mumps, incubation period of, 247.
- Muscles, wasting, causes of, 13.
- Mush, corn-meal, 41.
- Mutton, 35.
- Nephritis, relation of weights to deaths from, 5.
- Nervous control, increased by exercise, 21.
- Nervous system, relation of weights to deaths from diseases of the, 5.
- Neurasthenia, cause of, 129.
- Newspaper, as cause of trouble in plumbing, 67.
- Night-clothing, desirability of, 21.
- Night urinals, 98.
- Nose, the, 9.
- Nose, indicating alcoholism, 9; indicating syphilis, 9.
- Notification, of epidemic disease, 252, 277.
- Oatmeal, cooking of, in field, 87.
- Occupation, diseases of, 148.
- Officer, line, responsibility of, for pure water-supply, 81; medical, as authority on drinking water, 109; medical, as police officer, 99; medical, responsibility of, for pure water-supply, 81.
- Officers, familiarity of, with sanitation, 73; of the line, coöperation of, with medical officers, 19.
- Oil-cloth, in mess rooms, 59.
- Oil, crude, use of, against flies, 94; use of, against mosquitoes, 269.
- Old age, relation of weights to deaths from, 5.
- Oleomargarine, 45.
- Onions, 42.
- Organisms, disease-producing, destruction of, 177; of air-borne diseases, 239.
- Overweights, ratio of deaths from, 4, 5.
- Oxygen, effect of, on endurance, 104.
- Pail system, for disposal of excreta, 97.
- Paper, tissue, issue of, 64.
- Paralysis, general, relation of weights to deaths from, 5.
- Parasites, large, 162; minute, 170; prevention of infestation of, 108, 127; transmission of infestation of, 213; vegetable, 174.
- Paratyphoid, transmission of, 37.
- Pasteurization, 177.
- Pastries, 40.
- Pease, 42.
- Peculiarity, personal, 144.
- Peddlers, exclusion of, from camp, 91.
- Penis, abnormalities of, 15; puffiness of, 16; redness of, 16; signs of irritation on, 16; sores on, 16; ulcers on, 16.
- Pensions, possible grounds for, care in noting them, 19.
- Personal hygiene, 20.
- Persons, disinfection of, 254.
- Pest, the, see Plague.
- Petroleum, use of, against mosquitoes, 269; use of, in camps, 78.
- Pets, animal, exclusion of, from kitchen, 61; from camps, 78.
- Phenol, as antiseptic, 210.
- Physicians, civilian, practice of, in posts, 252.
- Pickling, as to meats, 38.
- Pigs, as carriers of anthrax, 191; as carriers of tuberculosis, 190.
- Piles, 7, 8.
- Pinworm, 165.
- Pipe-smoking, 30.
- Pits, excreta, care of, 94.
- Plague, infection of, 126, 129, 273; measures against, 275; toxin of, 179; transmission of, 187, 273; vaccination against, 278.
- Pleurisy, chronic, and chest type in, 6.
- Plumbing, 66; absence of, in squad-rooms, 58; company commander's familiarity with, 65; poor, of closets indicated by bad odors, 63.
- Pneumonia, cause of, 20; influenced

- by neglected mouth, 25; relation of weights to deaths from, 5; transmission of, 181.
- Poisoning, metallic, cause of, by canned foods, 46; ptomaine, causes of, 37; toxic, influenced by neglected mouth, 25; wood alcohol, 29; tyrotoxic, 45.
- Poisons, as cause of death or disease, 158; bacterial, 179; vegetable, 160.
- Police, as factor in causes of disease, 153; camp, 99; in prevention of epidemics, 206; in tropics, 129; kitchen, to be rewarded for efficiency, 61; of camps, importance of, 77; of kitchen, as punishment, 52; on the march, 114; officer, to be a medical officer, 99.
- Poncho, when carried, 111.
- Pork, 36.
- Potash, permanganate, use of, with formalin, 209.
- Potatoes, cooking of, in field, 87; cooking and serving of, 41.
- Practice-marches, as guides to camp craft, 87.
- Prairie dogs, as plague-carriers, 189.
- Preservatives, food, 38.
- Pressure, air, as cause of disease, 158.
- Privies, danger of, to water-supply, 83.
- Privileges, as reward for work in kitchen, 53.
- Problems, sanitary, officers' familiarity with, 73.
- Prostate, inflammation of, gonorrhoeal, 288.
- Proteids, 33, 122.
- Provost sergeant, directed by medical officer, 99.
- Ptomaine poisoning, causes of, 37.
- Ptomaines, cause of, 160.
- Pus, in feces, surgeon to be consulted, 64.
- Putty, as aid in exclusion of vermin, 72.
- Quarantine, 205; as part of camp police, 99.
- "Quarters," disadvantage of treatment in, 204.
- Quinine, as specific for malaria, 212, 272.
- Rabbits, as carriers of anthrax, 191.
- Race, as factor in disease, 146.
- Ration, for tropics, 122; full use of, in field, 87; increase of, in Alaska, 133.
- Rations, army, books of reference on, 53.
- Rats, as plague-carriers, 130, 188, 273; measures against, to prevent epidemics, 276; prevention of, in camp, 79; protection from, 52.
- Ray-fungus, transmission of, 37.
- Rays, actinic, protection from, 125.
- Reaction, prophylactic, 201.
- Recreation-rooms, 58.
- Recruit, the, 3; age limit for first enlistment, 3; arms of, 12; baldness of, 12; character of, 18; cough of, 10; condition of, if immature, 3; defects of, if noted by C. O. to be referred to medical officer, 19; disproportional size of, 4; ears of, 9; eyes of, 8; feet of, 12; fixity of habits of, after certain age, 4; genitals of, 15; head of, 8; hoarseness of, 10; legs of, 12; loss of voice in, 10; lymphatics of, 11; mouth of, 9; nose of, 9; presence of vermin on, 11; proportionate weight and chest measurement of, how published, 4; rejection of, on account of tumors, 7; rejection of, on account of gallstone colic, 8; rejection of, on account of severe indigestion, 8; rejection of, on account of vomiting of blood, 8; scalp of, 11; size of, 4; skull of, 17; teeth of, 9, 10; throat of, 9, 10; tongue of, 11.
- Reed trough, 92, 93, 97.
- Reed Board, findings of, as to typhoid, 213, 215, 219, 222, 225, 227, 229, 233.
- "Regulations, Field Service," observation of, 74.
- Reptiles, protection from, 111.
- Respiratory system, relation of weights to deaths from diseases of the, 5.
- Rest, importance of, 20, 102.

- Rheumatism, gonorrhœal, 288.
 Rice, cooking of, in field, 87; cooking and serving of, 41.
 Rickets, deformed breastbone as result of, 6.
 Rinds, ham, use of, in soups, 42.
 Ringworm, 17; prevention of, 128; transmission of, 22, 182, 191.
 Roaches, as disease-carriers, 193; prevention of, in camp, 79; protection from, 52.
 Roads, state of, 103.
 Rooms, 56.
 Roundworms, 165.
- Salads, 43.
 Salivation, mercurial, influence of neglected mouth on, 25.
 Salting, as to meats, 38.
 Salts, inorganic, food-value of, 34; urinary, formation of, in urinals, 64.
 Sanitation, desiderata as to supreme control of, 99.
 Scalp, cleanliness of, 23; of recruit, 11.
 Scarlet fever, incubation period of, 247; isolation of, 205; transmission of, 44, 181, 191, 251.
 Schistosomum disease, 152, 162.
 Scratching, signs of, 17.
 Screening, of camp kitchen, 89.
 Screens, necessity of, 57, 59.
 Screw-worms, 169; protection against cause of, 111.
 Scrotum, enlargement of, 16.
 Scurvy, 133.
 Seasickness, 118.
 Season, influence of, on epidemics, 241.
 Seatworm, 165.
 Secretions, in causes of epidemics, 244.
 Self-poisoning, 161.
 Self-respect, indication of, by cleanliness, 22.
 Service, as a refuge after failure in civil life, 4.
 Sex, self-control of, in tropics, 128.
 Sexual life, 27.
 Sheep, as carriers of tuberculosis, 190.
 Shelter, of camps, 77.
- Shins, crooked, 13.
 Shirts, flannel, use of, in tropics, 111.
 Shoes, 112; use of, in tropics, 126.
 Shower-baths, advantages of, over tubs, 62; improvised, 84.
 Sick, the, avoidance of, 204.
 Sickness, influence of, on marches, 101, 105.
 Sinks, kitchen, care of, 52.
 Site, of camps, 74.
 Size, disproportion of, in the recruit, 4.
 Skin, diseases of the, differentiation by line officers, 17; diseases of the, relation of weights to deaths from, 5; in causes of epidemics, 244; of recruit, 17; the, as a wall against germ invasion, 200.
 Skin diseases, transmission of, by domestic animals, 191; transmission of, by toilet articles, 22.
 Sleep, amount of, 21.
 Sleeping, as factor in cause of disease, 153.
 Sleeping-sickness, transmission of, 75, 173, 278.
 Slop-huckets, care of, 88.
 Smallpox, bacteria in, 179; incubation period of, 247; isolation of, 205; transmission of, 23, 183, 191; vaccination against, 235.
 Smoking, as cause of exhaustion, 107; as meat preservative, 38; effects of, 29; in day-rooms, 58.
 Snake-venoms, 162.
 Snow-blindness, 136.
 Snow-eyes, 137.
 Soils, camp-site, kinds desirable, 75.
 Sore throat, cause of, 20; diphtheria in, 181; influenced by neglected mouth, 25.
 Soup-carts, field use of, 87, 110.
 Spectacles, wooden, for snow-blindness, 137.
 Spices, use of, in tropics, 123.
 Spine, curvature of, 6.
 Spirilli, definition of, 176.
 Spitting, danger of, 26; promiscuous, by chewers of tobacco, 29, 59.
 Spores, bacterial, 176.
 Sports, encouragement of, 21.
 Spotted fever, transmission of, 193, 279.

- Sputum, disinfection of, 254; in causes of epidemics, 243.
- Squad-rooms, 56.
- Squad, sanitary, of camps, 99.
- Squirrels, as carriers of plague, 189.
- Staphylococci, 176.
- Starches and sugars, 34.
- Steam-heating, 70.
- Stegomyia fasciata*, carrier of yellow fever, 261.
- Sterilization, 207.
- Sterilizer, Forbes-Waterhouse, 81.
- Store-rooms, 62.
- Stoves, 68.
- Straw, for use in camps, 78.
- Strength, muscular, dependent on heart, 21.
- Streptococci, 176.
- Sublimate, corrosive, 210.
- Sucking-worms, 162.
- Sugars and starches, 34.
- Suggestion, mental, for nervous disorders, 126.
- Suicides, relation of weights to deaths from, 5.
- Sulphur, as disinfectant, 208.
- Sunlight, effect of, on bacteria, 178.
- Sunlight, strong, exposure to, 31.
- Sun-stroke, predisposed by want of water, 107.
- Sun, consideration to, in placing buildings, 55.
- Suppuration, germs of, 194.
- Surra and sleeping sickness, relation of, 173.
- Susceptibility, to water-borne diseases, 219.
- Swamps, malarial, 76.
- Sweets, 47.
- Swimming, 127.
- Syphilis, 173, 183, 292; fourth stage of, 296; in company cook, 51; indicated by condition of nose, 9; indications of, 9, 10, 11, 12, 14, 17; influenced by neglected mouth, 25; prevention of, 297; primary and secondary, 295; tertiary, 296.
- Syringes, rectal, as disease-carriers, 222.
- Tapeworms, 163; transmission of, 37.
- Tea, 47; on the march, 109.
- Teeth, care of, 25; loss of, 25; of recruit, 9-10.
- Temperature, demands of, as to clothing worn, 30; in causes of disease, 158.
- Tendon, inflammation of, gonorrhoeal, 288.
- Tents, 77.
- Testicles, condition of, 15.
- Tests, for purity of water, 82.
- Tetanus, cause of, 176.
- Throat, of recruit, 9, 10.
- Trough, Reed, 92, 93, 97.
- Thumb, loss of, 12.
- Tick fever, 193.
- Ticks, 170, 193; as carriers of disease, 74, 75, 193, 279; protection against, 111, 126.
- Tin cans, disposal of, in field, 89.
- Toasting, bread, 40.
- Tobacco, 29.
- Toilet articles, exclusive ownership of, 62; transmission of disease by, 22.
- Toilet-paper, camp use of, 94.
- Tongue, of recruit, 11.
- Tonsillitis, incubation period of, 247. transmission of, 182.
- Tonsils, enlargement of, 10.
- Tooth-brush, use of, 26.
- Tooth-trouble, prompt report of, to medical officer, 26.
- Towel, care of, in field, 113.
- Towels, liberal use of, by company barber, 62.
- Toxins, bacterial, 179.
- Traffic, regulated by camp police, 99.
- Training, lack of, as cause of disease, 146.
- Transportation facilities, effect of, on marches, 103.
- Transports, 117.
- Trappings, useless, exclusion of, from squad-rooms, 58.
- Traumatism, 156.
- Trembles, in cattle, transmission of, cause of, 37.
- Tremor, tobacco, 29.
- Trenches, in disposal of wastes, 93, 96.
- Trypanosome, 173.
- Trichina, transmission of, 37.

- Trichiniasis, 167.
 "Trikresol," as disinfectant, 211.
 Troop trains, 118.
 Troops, assemblage of, danger in, 73; moving, hygiene of, 101.
 Tropics, clothing in, 111.
 Tuberculosis, indication of, 14; influenced by neglected mouth, 25; in company cook, 51; relation of weights to deaths from, 5; toxin of, 179; transmission of, 26, 37, 44, 182, 189, 193.
 Tumors, as cause for rejection of recruit, 7.
 Typhoid, as camp disease, 76; cause of, 214; death rate of, 219; destruction of organism of, 80; isolation of, 205, 229; milk-borne epidemics of, 220; transmission of, 22, 23, 26, 37, 44, 51, 98, 108, 127, 182, 183, 193, 213; relation of weights to deaths from, 5; toxin of, 179; vaccination against, 236.
 Typhus, incubation period of, 247; transmission of, 252.
 Tyrotoxicosis, development of, 45.
 Ulcers, in throat of recruit, 10; on penis, 16; on leg of recruit, 13.
 Underclothing, care of, on the march, 113; kind for tropics, 125; not to be slept in, 21.
 Underweights, table of death rate of, 4, 5.
 Urethra, contents of, 15.
 Urinals, 64, 67, 97.
 Urination, careless, danger of, 26; near tents in camp, 79.
 Urine, antiseptic action of, 201; cause of insufficiency of water on the, 107; disposal of, to prevent epidemics, 233; flowing of, 17; human, disposal of, in camp, 91.
 Urine pit, importance of, 95.
 Utensils, as carriers of epidemic disease, 245; disinfection of, 254; barber's, sterilization of, 61; kitchen, cleanliness of, in field, 88.
 Vaccination, against epidemic disease, 236, 255, 278.
 Varicocele, 16.
 Veal, 36.
 Vegetables, 41, 152; as carriers of typhoid, 221; as cause of intestinal diseases, 122; control of sale of, during epidemics, 236; underdone, as cause of indigestion, 87.
 Veins, dilated, of leg, 13; distended abdominal, cause for rejection of recruit, 7; varicose, of abdomen, cause for rejection of recruit, 7; varicose, of leg, 13.
 Venereal, 281.
 Venereal disease, veracity as to, in recruits, 17.
 Ventilation, 56, 59, 68, 78, 204, 250; in arctics, 131; in prevention of epidemics, 204.
 Vermin, as cause of epidemics, 223; destruction of, by sulphur, 208; elimination of, to prevent epidemics, 233, 252; exclusion of, from kitchen, etc., 52, 55, 71, 129; on recruit, 11, 16, 17, 22; spread of, in camps, 78.
 Violence, relation of weight to deaths from, 5.
 Vision, testing of, 8.
 Vitality, influence of weight on, 4, 5.
 Voice, loss of, in recruit, 10.
 Voit, food-standard of, 35.
 Vomit, disposal of, to prevent epidemics, 233.
 Walls, damp, avoidance of, 55.
 "Warbles," 169.
 Warts, venereal, 16.
 Waste, channels of excretion of, affected by exercise, 21; disposal of, in camp, 91.
 Waste water, disposal of, in camp, 85; kitchen, disposal of, in camp, 89.
 Water, boiling, for dilution of milk, 45; use of, as vermin killer, 72.
 Water, food-value of, 35; sterilization of, in camp, 80.
 Water-carriage latrines, 92.
 Water-closets, 57, 63.
 Water-supply, 65, 80, 106, 121, 251; as disease-carrier, 26, 98, 150, 193, 219, 235; in arctics, 132.
 Weather, influence of, on marches, 101; protection from inclemencies of, 80.

- Weight, carried on marches, 103.
Weight and chest measurements for recruits, publication of, 4.
"Wet dreams," 27.
Whipworm, 165.
Whooping-cough, incubation period of, 247; transmission of, 191.
Wind, regard to, in placing buildings, 55.
Windows, storm, 57; variable needs of, 56.
Winds, value of, in camps, 76.
- Wire-gauze, use of, in field kitchen, 89.
Wolf, as carrier of hydrophobia, 184.
Wool-sorters' disease, 191.
Worms, as animal parasites, 162; intestinal, transmission of, 182; in feces, surgeon to be consulted, 64.
Yaws, 173.
Yeasts, as agents of disease, 174.
Yellow fever, 192, 260; isolation of, 205; protection against, 111, 129.





The Riverside Press
CAMBRIDGE . MASSACHUSETTS
U . S . A







